

## Catch – Up Schedule - Test

1. Karen is 2 years old and had HIB dose 2 at age 15 months. When is her next dose due? (Answer – no further doses needed)
2. Tika is 4 yr. 5 months old. He had one dose of OPV in Nepal at age 2months old. He had one dose of IPV at age 3 yrs old. How many doses does he need to complete his series and when are they due? (Answer -he must have a total of 4 doses because he has a mixed vaccine hx (IPV and OPV). Dose 3 is due now. Dose 4 is due 6 months after dose 3.
3. Joey will be 5years old next month. He had 4 doses of PCV 7 (Prevnar™). His last dose was at age 12 months. Is he due for any more pneumococcal vaccine? (explain) (Answer – yes Joey is due for one dose of PCV 13 before his 5<sup>th</sup> birthday. Because his other four doses were PCV 7 he should have one dose of PCV 13 to cover the serotypes he did not receive with PCV 7.)
4. Ahmed is 4 years old and just had dose 4 of DTaP today. Dose 3 was given 8 months ago. When is his next dose of tetanus due? (Answer – Ahmed is now UTD – his next routine dose will be Tdap at his 11 yr old WCC).
5. If Miranda (age 12 years old) had varicella vaccine dose one today, when can she have dose two? (Answer -varicella dose 2 is due in 3 months – minimum interval)
6. If Amir is 10 years old and has never had tetanus, diphtheria or pertussis vaccine how will you administer his series doses? (Answer - Tdap dose one now, Td in 4 weeks, Td 6 months after dose 2). (Tdap can be substituted for one dose in this 3 doses series, but since he has never had pertussis vaccine it would make sense to give this as the initial dose)
7. What is the recommended interval between doses 1 and 2 of HPV vaccine? What is the minimum interval between these doses? (Answer - recommended interval is 2 months – minimum interval is 4 weeks)
8. If an infant turns 8 months old next week, and his first dose of RotaTeq was 5 months ago, when can he have his remaining doses? (Answer – dose 2 must be given by 8months 0 days. No other doses can be given after that)
9. What are the 3 minimum age/interval criteria for administering Hepatitis B dose number three? (Answer – minimum interval of 8 weeks between dose 2 to 3. Minimum interval of 16 weeks between doses 1 and 3. Infant must be minimum of 6 months (24 weeks) at the time of the 3<sup>rd</sup> dose.)
10. Rebecca age 3 years old recently had a cochlear implant. She finished 4 doses of PCV 7 at age 12 months. What kind(s) of pneumococcal vaccine should she have today? (Answer – PCV 13 one dose now and PPSV (pneumovax™) one dose at least 8 weeks after the PCV 13 dose)

NAME: \_\_\_\_\_ date: \_\_\_\_\_