

Table 1C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Demographic Indicators ⁽²⁾

Total Records	Number	%	Source of Data	Number	%
Complete Records	3,555	70.5	WIC	5,046	100.0
Prenatal Records	771	15.3	Maternal Health	0	0.0
Postpartum Records	720	14.3	Other	0	0.0
Total Records Accepted	5,046		Total	5,046	100.0
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%
White, Not Hispanic	4,359	86.4
Black, Not Hispanic	171	3.4
Hispanic	344	6.8
American Indian/Alaska Native	35	0.7
Asian/Pacific Islander	110	2.2
Multiple Races	27	0.5
All Other	0	0.0
Total	5,046	100.0
<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Age Distribution	Number	%
< 15 Years	3	0.1
15 - 17 Years	205	4.1
18 - 19 Years	612	12.1
20 - 29 Years	3,222	63.9
30 - 39 Years	925	18.3
>= 40 Years	76	1.5
Total	5,043	100.0
<i>Excluded Unknown: 3</i>	<i>0.1</i>	

Education Distribution	Number	%
< High School	775	23.7
High School	1,463	44.7
> High School	1,038	31.7
Total	3,276	100.0
<i>Excluded Unknown: 1,770</i>	<i>35.1</i>	

% Poverty Level	Number	%
0 - 50	1,356	31.6
51 - 100	1,300	30.3
101 - 130	678	15.8
131 - 150	339	7.9
151 - 185	418	9.8
186 - 200	34	0.8
> 200	54	1.3
Adjunctive Eligibility	108	2.5
Total	4,287	100.0
<i>Excluded Unknown: 39</i>	<i>0.9</i>	

Program Participation ⁽³⁾	Number	%
WIC	4,326	100.0
SNAP	4,326	33.5
Medicaid	4,326	47.9
TANF	4,326	17.8
<i>Excluded Unknown:</i>		
<i>WIC:</i>	0	0.0
<i>SNAP:</i>	0	0.0
<i>Medicaid:</i>	0	0.0
<i>TANF:</i>	0	0.0

Migrant Status	Number	%
Yes	1	0.0
No	5,045	100.0
Total	5,046	100.0
<i>Excluded Unknown: 0</i>	<i>0.0</i>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Health Indicators

Summary of Health Indicators (2)				
	New Hampshire Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	4,894	5.3	1,226,875	4.5
Normal Weight	4,894	43.3	1,226,875	42.5
Overweight	4,894	23.9	1,226,875	25.9
Obese	4,894	27.4	1,226,875	27.0
Maternal Weight Gain (4)				
Less Than Ideal	4,094	18.1	917,032	21.2
Ideal	4,094	27.3	917,032	30.6
Greater Than Ideal	4,094	54.7	917,032	48.2
Anemia (Low Hb/Hct) (5)				
1st Trimester	1,560	4.9	436,474	7.7
2nd Trimester	1,326	9.5	375,216	12.2
3rd Trimester	734	26.3	189,944	33.9
Postpartum	983	22.8	412,633	28.8
Parity (6)				
0	4,617	33.9	1,108,534	41.3
1	4,617	29.3	1,108,534	26.7
2	4,617	17.9	1,108,534	16.3
≥ 3	4,617	18.9	1,108,534	15.6
Interpregnancy Interval				
< 6 Months	1,794	13.2	614,765	12.0
6 - <18 Months	1,794	28.4	614,765	26.7
≥18 Months	1,794	58.5	614,765	61.3
Gestational Diabetes	4,176	5.9	389,161	5.0
Hypertension During Pregnancy	4,183	10.0	394,368	6.8

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Health Indicators

Summary of Health Indicators (2)				
	New Hampshire Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	3,744	89.8	527,184	82.2
No Care	3,744	2.4	527,184	3.1
WIC Enrollment				
1st Trimester	5,027	37.1	1,213,982	34.1
2nd Trimester	5,027	31.7	1,213,982	34.7
3rd Trimester	5,027	16.9	1,213,982	17.5
Postpartum (3)	5,027	14.3	1,213,982	13.7
Prepregnancy Multivitamin Consumption (4)	5,046	12.7	360,369	18.7
Multivitamin Consumption During Pregnancy (5)	4,322	62.5	372,743	83.5
Smoking				
3 Months Prior to Pregnancy	4,713	41.1	1,040,568	22.9
During Pregnancy	4,024	26.9	895,533	13.8
Last 3 Months of Pregnancy	3,517	25.0	838,781	13.0
Postpartum	3,611	26.9	782,035	15.0
Smoking Changes (6)				
Quit Smoking by First Prenatal Visit	1,674	36.1	206,583	43.1
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	1,145	29.8	156,709	36.7
Smoking in Household				
During Pregnancy	3,967	22.2	650,648	17.1
Postpartum	3,620	16.1	559,995	16.8
Drinking				
3 Months Prior to Pregnancy	4,595	21.8	1,040,654	8.3
Last 3 Months of Pregnancy	3,461	0.3	604,343	1.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

(6) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Health Indicators

Summary of Health Indicators (2)				
	New Hampshire Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	2,505	1.2	1,018,145	1.1
Low	2,505	6.5	1,018,145	7.1
Normal	2,505	83.0	1,018,145	85.1
High	2,505	9.3	1,018,145	6.7
Preterm (4)	4,203	6.9	1,017,333	11.1
Full term Low Birth Weight (5)	2,318	4.3	848,957	3.5
Multiple Births	4,254	3.8	1,095,174	1.4
Breastfeeding				
Ever Breastfed	4,183	73.9	997,136	66.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.



Table 3C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Comparison of Racial and Ethnic Distribution by Contributor to Nation

Comparison of Racial and Ethnic Distribution ⁽²⁾								
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
New Hampshire	5,046	86.4	3.4	6.8	0.7	2.2	0.5	0.0
Nation (Prior Year)	1,304,320	41.6	24.1	28.3	1.1	3.2	1.2	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Comparison of Age Distribution by Contributor to Nation

Comparison of Age Distribution ⁽²⁾								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
New Hampshire	5,046	0.1	4.1	12.1	63.9	18.3	1.5	0.1
Nation (Prior Year)	1,304,320	0.3	5.8	12.3	59.4	20.4	1.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Comparison of Maternal Health Indicators by Contributor to Nation

Comparison of Maternal Health Indicators ⁽²⁾														
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
New Hampshire	4,894	5.3	51.3	4,094	18.1	54.7	734	26.3	983	22.8	4,176	5.9	4,183	10.0
Nation (Prior Year)	1,226,875	4.5	52.9	917,032	21.2	48.2	189,944	33.9	412,633	28.8	389,161	5.0	394,368	6.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Comparison of Maternal Behavioral Indicators by Contributor to Nation

Comparison of Maternal Behavioral Indicators ⁽²⁾									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
New Hampshire	3,744	89.8	2.4	5,027	37.1	5,046	12.7	4,322	62.5
Nation (Prior Year)	527,184	82.2	3.1	1,213,982	34.1	360,369	18.7	372,743	83.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Comparison of Smoking Indicators by Contributor to Nation

Comparison of Smoking Indicators (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	%	Number	%	Number	%	Number	%		
New Hampshire	4,713	41.1	3,517	25.0	1,145	29.8	3,967	22.2	3,620	16.1
Nation (Prior Year)	1,040,568	22.9	838,781	13.0	156,709	36.7	650,648	17.1	559,995	16.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Comparison of Infant Health Indicators by Contributor to Nation

Comparison of Infant Health Indicators ⁽²⁾									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)	High (4)							
	Number	%	%	Number	%	Number	%	Number	%
New Hampshire	2,505	7.7	9.3	4,203	6.9	2,318	4.3	4,183	73.9
Nation (Prior Year)	1,018,145	8.2	6.7	1,017,333	11.1	848,957	3.5	997,136	66.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 9C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Maternal Health Indicators by Race/Ethnicity, Age, or Education

Maternal Health Indicators by Race/Ethnicity, Age, or Education ⁽²⁾

	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes		Hypertension During Pregnancy		
	Number	Under-weight	Over-weight	Number	< Ideal	> Ideal	3rd Trimester		Postpartum		Number	%	Number	%	
		%	%		%	%	%	Number	%	Number					%
Race/Ethnicity															
White, Not Hispanic	4,226	5.4	51.8	3,547	17.6	55.6	646	24.5	874	21.5	3,613	6.0	3,618	10.0	
Black, Not Hispanic	163	2.5	54.0	128	26.6	43.8	22	*	25	*	138	3.6	138	8.7	
Hispanic	335	3.0	50.7	280	19.3	51.4	37	*	54	*	285	4.9	286	9.8	
American Indian/Alaska Native	35	*	*	27	*	*	5	*	8	*	25	*	25	*	
Asian/Pacific Islander	109	14.7	24.8	88	*	*	20	*	21	*	91	*	91	*	
Multiple Races	26	*	*	24	*	*	4	*	1	*	24	*	25	*	
All Other/Unknown	0	*	*	0	*	*	0	*	0	*	0	*	0	*	
Total	4,894	5.3	51.3	4,094	18.1	54.7	734	26.3	983	22.8	4,176	5.9	4,183	10.0	
Age															
< 15 Years	3	*	*	3	*	*	0	*	2	*	3	*	3	*	
15 - 17 Years	197	7.1	35.0	175	11.4	58.9	24	*	39	*	179	1.7	179	11.7	
18 - 19 Years	592	8.3	42.7	502	13.3	63.5	78	*	122	26.2	510	2.5	511	9.2	
20 - 29 Years	3,135	5.5	50.6	2,619	18.7	54.4	482	25.3	610	23.3	2,654	5.2	2,658	9.3	
30 - 39 Years	892	2.6	63.0	734	19.8	49.9	139	27.3	186	18.3	764	11.0	766	12.0	
>= 40 Years	72	*	*	58	*	*	11	*	24	*	63	*	63	*	
Unknown	3	*	*	3	*	*	0	*	0	*	3	*	3	*	
Total	4,894	5.3	51.3	4,094	18.1	54.7	734	26.3	983	22.8	4,176	5.9	4,183	10.0	
Education															
< High School	759	5.7	47.3	629	19.7	53.3	96	*	132	27.3	647	4.2	648	9.7	
High School	1,430	5.8	52.0	1,218	18.0	55.7	232	26.3	276	25.4	1,257	5.9	1,258	9.5	
> High School	1,008	4.4	52.4	874	15.7	56.3	174	19.5	211	19.4	908	6.5	908	9.9	
Unknown	1,697	5.4	52.0	1,373	18.9	53.5	232	22.4	364	21.2	1,364	6.2	1,369	10.5	
Total	4,894	5.3	51.3	4,094	18.1	54.7	734	26.3	983	22.8	4,176	5.9	4,183	10.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 10C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education (2)									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	3,239	90.6	2.2	4,344	37.3	4,359	13.4	3,711	62.7
Black, Not Hispanic	123	79.7	5.7	171	28.1	171	8.8	154	66.2
Hispanic	257	86.4	3.9	340	43.2	344	7.0	309	57.6
American Indian/Alaska Native	22	*	*	35	*	35	*	26	*
Asian/Pacific Islander	79	*	*	110	25.5	110	9.1	98	*
Multiple Races	24	*	*	27	*	27	*	24	*
All Other/Unknown	0	*	*	0	*	0	*	0	*
Total	3,744	89.8	2.4	5,027	37.1	5,046	12.7	4,322	62.5
Age									
< 15 Years	3	*	*	3	*	3	*	2	*
15 - 17 Years	161	80.7	5.0	204	43.6	205	6.3	182	62.6
18 - 19 Years	457	86.7	4.2	609	41.2	612	5.7	550	54.0
20 - 29 Years	2,409	90.5	2.3	3,211	37.2	3,222	12.8	2,760	62.6
30 - 39 Years	659	91.5	1.2	921	33.3	925	17.9	764	67.5
>= 40 Years	52	*	*	76	*	76	*	61	*
Unknown	3	*	*	3	*	3	*	3	*
Total	3,744	89.8	2.4	5,027	37.1	5,046	12.7	4,322	62.5
Education									
< High School	566	84.6	4.8	770	44.9	775	5.9	732	61.1
High School	1,125	90.0	1.7	1,460	38.5	1,463	9.7	1,320	65.6
> High School	827	93.5	1.7	1,034	36.7	1,038	20.7	899	74.7
Unknown	1,226	89.4	2.5	1,763	32.8	1,770	13.4	1,371	52.2
Total	3,744	89.8	2.4	5,027	37.1	5,046	12.7	4,322	62.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 11C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Smoking Indicators by Race/Ethnicity, Age, or Education

Smoking Indicators by Race/Ethnicity, Age, or Education ⁽²⁾										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity										
White, Not Hispanic	4,060	44.8	3,022	27.7	1,078	29.2	3,394	24.2	3,119	17.3
Black, Not Hispanic	164	21.3	125	10.4	19	*	140	6.4	122	6.6
Hispanic	325	17.5	243	6.6	32	*	296	11.1	251	9.2
American Indian/Alaska Native	32	*	23	*	8	*	24	*	24	*
Asian/Pacific Islander	105	3.8	83	*	3	*	89	*	82	*
Multiple Races	27	*	21	*	5	*	24	*	22	*
All Other/Unknown	0	*	0	*	0	*	0	*	0	*
Total	4,713	41.1	3,517	25.0	1,145	29.8	3,967	22.2	3,620	16.1
Age										
< 15 Years	3	*	3	*	1	*	2	*	3	*
15 - 17 Years	193	38.3	151	17.2	45	*	169	35.5	158	23.4
18 - 19 Years	575	51.3	443	30.7	191	34.0	494	33.0	453	22.5
20 - 29 Years	3,007	42.5	2,226	26.1	752	29.4	2,535	21.2	2,298	16.0
30 - 39 Years	863	31.4	645	19.4	147	21.8	709	15.8	657	10.8
>= 40 Years	69	*	47	*	7	*	55	*	49	*
Unknown	3	*	2	*	2	*	3	*	2	*
Total	4,713	41.1	3,517	25.0	1,145	29.8	3,967	22.2	3,620	16.1
Education										
< High School	763	49.4	596	32.2	248	23.0	708	28.7	605	21.7
High School	1,437	44.3	1,110	27.5	416	29.3	1,287	22.0	1,133	17.3
> High School	1,009	29.8	792	15.7	191	37.2	869	16.6	814	10.7
Unknown	1,504	41.4	1,019	25.3	290	31.4	1,103	22.6	1,068	15.9
Total	4,713	41.1	3,517	25.0	1,145	29.8	3,967	22.2	3,620	16.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 12C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Infant Health Indicators by Race/Ethnicity, Age, or Education

Infant Health Indicators by Race/Ethnicity, Age, or Education (2)									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)						
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	2,164	7.3	9.9	3,635	6.7	2,012	4.0	3,621	72.5
Black, Not Hispanic	66	*	*	136	5.9	60	*	132	85.6
Hispanic	178	11.8	5.1	287	10.1	155	7.7	286	81.5
American Indian/Alaska Native	15	*	*	27	*	15	*	27	*
Asian/Pacific Islander	67	*	*	93	*	63	*	92	*
Multiple Races	15	*	*	25	*	13	*	25	*
All Other/Unknown	0	*	*	0	*	0	*	0	*
Total	2,505	7.7	9.3	4,203	6.9	2,318	4.3	4,183	73.9
Age									
< 15 Years	1	*	*	2	*	1	*	2	*
15 - 17 Years	109	4.6	6.4	182	7.7	99	*	182	65.9
18 - 19 Years	316	7.6	6.6	516	6.6	293	4.4	514	66.9
20 - 29 Years	1,600	6.3	9.4	2,677	6.6	1,488	4.2	2,668	74.7
30 - 39 Years	444	7.0	12.4	761	7.8	406	4.9	752	77.0
>= 40 Years	34	*	*	62	*	30	*	62	*
Unknown	1	*	*	3	*	1	*	3	*
Total	2,505	7.7	9.3	4,203	6.9	2,318	4.3	4,183	73.9
Education									
< High School	373	11.3	7.2	642	8.9	336	6.5	641	64.3
High School	690	6.7	10.7	1,251	8.2	631	4.1	1,243	71.6
> High School	460	4.8	8.9	901	5.5	432	3.7	898	85.6
Unknown	982	5.4	9.4	1,409	5.8	919	3.8	1,401	72.7
Total	2,505	7.7	9.3	4,203	6.9	2,318	4.3	4,183	73.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 13C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Maternal Weight Gain and Birth Outcomes by Select Health Indicators

Maternal Weight Gain and Birth Outcomes by Select Health Indicators (2)										
	Weight Gain (3)			Birth Outcomes						
	< Ideal	> Ideal		Birthweight			Preterm (6)	Full Term LBW (7)		
		Number	%	%	Low (4)	High (5)				
	Number	%	%	Number	%	%	Number	%	Number	%
Prepregnancy BMI										
Underweight	214	22.0	36.0	127	7.9	2.4	211	7.1	118	5.1
Normal Weight	1,767	18.7	48.7	1,057	8.6	7.6	1,758	7.5	970	4.8
Overweight	991	12.2	67.2	604	6.0	10.1	988	6.2	569	3.2
Obese	1,122	21.5	56.6	665	6.9	13.4	1,119	7.4	609	3.0
Weight Gain										
< Ideal	-	-	-	415	13.5	5.8	704	9.1	375	7.7
Ideal	-	-	-	650	6.5	6.0	1,102	7.4	603	3.6
> Ideal	-	-	-	1,360	6.1	12.2	2,222	6.2	1,263	2.9
Smoking During Pregnancy										
Smokers	868	21.2	50.2	552	10.1	6.7	870	8.7	501	5.4
Non-Smokers	2,407	16.1	57.1	1,460	5.9	10.9	2,423	8.0	1,337	2.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Racial and Ethnic Distribution

Summary of Trends in Racial and Ethnic Distribution ⁽²⁾								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2010	5,046	86.4	3.4	6.8	0.7	2.2	0.5	0.0
2009	5,212	87.1	3.2	6.6	0.7	2.1	0.3	0.1
2008	4,751	87.7	3.4	6.3	0.7	1.6	0.3	0.0
2007	4,596	86.4	3.6	7.0	0.8	1.8	0.5	0.0
2006	4,487	89.0	3.3	5.1	0.4	1.5	0.4	0.1
2005	4,902	90.7	3.7	1.1	0.1	1.2	0.0	3.2
2004	4,966	91.2	2.6	3.3	0.1	1.2	0.0	1.6
2003	4,760	94.3	1.5	3.3	0.0	0.9	0.0	0.0
2002	4,814	97.0	0.9	1.5	0.0	0.5	0.0	0.0
2001	4,869	97.0	1.0	1.5	0.0	0.5	0.0	0.0
2000	3,125	97.8	1.0	1.0	0.0	0.2	0.0	0.0
1999	4,519	97.7	0.8	1.0	0.0	0.5	0.0	0.0
1998	4,744	99.1	0.3	0.4	0.0	0.1	0.0	0.0
1997	3,989	99.2	0.2	0.4	0.0	0.1	0.0	0.0
1996	0	*	*	*	*	*	*	*
1995	0	*	*	*	*	*	*	*
1994	0	*	*	*	*	*	*	*
1993	0	*	*	*	*	*	*	*
1992	0	*	*	*	*	*	*	*
1991	470	98.1	0.2	1.3	0.0	0.4	0.0	0.0
1990	355	94.9	2.8	1.7	0.0	0.6	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 15C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2010	5,046	0.1	4.1	12.1	63.9	18.3	1.5	0.1
2009	5,212	0.2	4.6	12.3	63.7	18.1	1.1	0.0
2008	4,751	0.2	5.2	14.6	60.9	17.6	1.5	0.0
2007	4,596	0.2	5.1	14.4	63.4	15.4	1.4	0.1
2006	4,487	0.2	5.0	13.6	62.3	17.3	1.5	0.1
2005	4,902	0.2	5.3	13.7	61.4	17.8	1.6	0.0
2004	4,966	0.1	5.7	12.9	60.7	18.8	1.7	0.0
2003	4,760	0.2	5.4	14.3	60.8	18.1	1.2	0.0
2002	4,814	0.1	5.8	14.3	59.1	19.2	1.4	0.0
2001	4,869	0.2	6.4	13.9	60.3	18.0	1.2	0.0
2000	3,125	0.2	5.7	15.6	59.7	17.5	1.2	0.0
1999	4,519	0.3	7.1	15.5	59.5	16.8	0.8	0.0
1998	4,744	0.4	8.2	15.1	58.7	16.6	1.0	0.0
1997	3,989	0.2	7.4	15.2	59.1	17.2	0.9	0.0
1996	0	*	*	*	*	*	*	*
1995	0	*	*	*	*	*	*	*
1994	0	*	*	*	*	*	*	*
1993	0	*	*	*	*	*	*	*
1992	0	*	*	*	*	*	*	*
1991	470	0.2	6.6	13.4	59.4	19.6	0.6	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 15C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1990	355	0.3	9.3	14.9	54.4	20.0	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 16C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2010	4,894	5.3	51.3	4,094	18.1	54.7	734	26.3	983	22.8
2009	5,030	4.8	51.7	4,044	17.4	54.8	762	29.4	1,157	23.0
2008	4,674	4.8	48.5	3,951	16.6	56.2	747	27.7	1,898	21.3
2007	4,493	4.7	48.5	3,915	17.1	56.3	740	29.1	1,397	21.9
2006	4,400	4.6	48.4	3,842	17.2	56.0	812	25.2	1,539	22.6
2005	4,133	5.2	46.9	3,293	15.2	57.0	1,999	18.4	2,278	21.7
2004	3,707	5.4	48.4	2,775	54.1	25.6	1,296	18.5	1,219	22.0
2003	3,562	7.2	45.8	2,620	65.3	14.7	760	26.3	888	26.0
2002	3,601	7.1	45.3	2,878	68.2	14.0	760	27.2	957	27.9
2001	3,537	7.0	45.7	2,899	67.8	14.0	789	30.7	980	29.1
2000	2,262	8.1	46.0	2,076	68.6	14.0	443	28.7	705	25.1
1999	3,286	7.2	44.1	2,984	68.5	13.0	658	24.5	981	21.4
1998	3,389	7.2	43.1	3,111	68.3	14.4	671	22.8	1,309	24.1
1997	3,116	8.0	41.3	2,669	71.7	11.1	642	20.1	1,898	19.6
1996	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 16C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
1995	0	*	*	0	*	*	0	*	0	*
1994	0	*	*	0	*	*	0	*	0	*
1993	0	*	*	0	*	*	0	*	0	*
1992	0	*	*	0	*	*	0	*	0	*
1991	244	11.1	32.0	1	*	*	192	14.1	108	12.0
1990	138	8.0	34.8	23	*	*	91	*	135	14.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2010	3,744	89.8	2.4	5,027	37.1
2009	3,650	90.5	1.4	5,202	33.9
2008	2,981	90.3	1.5	3,732	38.8
2007	3,035	88.8	2.6	3,714	36.9
2006	2,604	88.8	1.0	4,090	30.2
2005	3,718	92.2	0.7	3,721	37.2
2004	3,611	91.5	0.9	3,856	40.5
2003	3,325	90.9	0.8	3,718	39.7
2002	3,766	43.1	53.1	3,778	40.9
2001	4,251	0.2	99.7	3,707	37.8
2000	2,962	0.1	99.9	2,362	39.3
1999	4,231	2.3	97.4	3,449	38.7
1998	2,360	79.4	7.6	3,567	40.4
1997	632	82.8	4.0	3,259	37.9
1996	0	*	*	0	*
1995	0	*	*	0	*
1994	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
1993	0	*	*	0	*
1992	0	*	*	0	*
1991	208	72.1	2.9	433	0.7
1990	232	75.9	1.7	319	1.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 18C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Trends in Smoking Indicators

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2010	4,713	41.1	3,517	25.0	1,145	29.8	3,967	22.2	3,620	16.1
2009	4,586	43.2	3,349	25.4	1,086	27.3	3,744	31.8	3,446	28.1
2008	3,857	42.8	2,659	27.2	853	22.0	2,996	36.9	2,672	34.8
2007	3,671	43.4	2,267	29.2	741	17.0	2,965	40.7	2,336	36.9
2006	3,323	44.1	2,286	27.4	635	22.7	1,311	40.8	2,248	38.6
2005	3,121	48.9	3,045	29.9	986	25.7	5	*	297	34.7
2004	3,086	51.3	2,855	33.0	985	22.6	0	*	0	*
2003	2,992	48.7	2,596	30.8	821	19.1	0	*	0	*
2002	3,099	50.0	2,764	32.1	896	21.0	0	*	0	*
2001	3,177	49.2	3,080	34.7	1,026	19.8	0	*	0	*
2000	2,053	48.9	2,332	33.7	783	19.7	0	*	0	*
1999	3,112	50.4	3,138	35.2	1,102	19.3	0	*	0	*
1998	3,297	52.3	3,262	36.3	1,196	21.7	0	*	0	*
1997	2,881	51.6	2,445	32.4	925	22.8	0	*	0	*
1996	0	*	0	*	0	*	0	*	0	*
1995	0	*	0	*	0	*	0	*	0	*
1994	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 18C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Trends in Smoking Indicators

Summary of Trends in Smoking Indicators (2)								
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household		
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum	
	Number	%	Number	%	Number	%	Number	%
1993	0	*	0	*	0	*	0	*
1992	0	*	0	*	0	*	0	*
1991	164	56.1	93	*	0	*	0	*
1990	46	*	86	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2010	2,505	7.7	9.3	4,203	6.9	2,318	4.3	4,183	73.9
2009	3,426	7.3	10.0	4,194	7.1	3,172	4.0	4,170	72.4
2008	3,697	6.2	9.9	3,023	7.5	2,662	2.1	3,644	74.3
2007	3,697	6.7	10.4	3,175	8.2	2,730	2.9	3,536	66.7
2006	3,414	6.7	8.9	3,525	7.5	2,878	3.8	3,600	65.3
2005	1,887	6.0	10.7	1,473	6.3	1,264	3.2	2,064	69.1
2004	3,345	6.6	10.2	2,559	9.7	2,298	1.7	3,409	61.8
2003	4,191	7.5	11.2	3,235	9.7	2,900	2.7	4,277	60.8
2002	4,240	6.8	10.9	3,274	9.0	2,959	2.8	4,311	60.3
2001	4,331	8.2	10.6	3,260	10.8	2,877	2.9	4,422	54.0
2000	2,900	8.2	12.2	2,211	10.9	1,944	3.2	2,974	50.7
1999	4,143	8.1	11.5	3,207	8.6	2,883	3.3	4,275	49.9
1998	4,290	6.3	11.8	3,306	8.7	2,926	2.8	4,483	46.9
1997	3,316	7.2	11.4	2,809	10.2	2,386	2.1	3,541	47.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1996	0	*	*	0	*	0	*	0	*
1995	0	*	*	0	*	0	*	0	*
1994	0	*	*	0	*	0	*	0	*
1993	0	*	*	0	*	0	*	0	*
1992	0	*	*	0	*	0	*	0	*
1991	202	6.9	9.9	177	11.3	147	2.0	199	64.3
1990	235	5.1	13.2	202	7.4	185	2.7	226	71.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2010	4,226	5.4	51.8	3,547	17.6	55.6	646	24.5	874	21.5
	2009	4,386	4.8	52.9	3,540	16.9	56.0	674	28.9	1,030	22.5
	2008	4,106	4.8	49.2	3,487	16.1	56.7	654	26.6	1,731	20.9
	2007	3,891	4.8	48.7	3,401	16.4	57.5	646	27.2	1,285	21.0
	2006	3,921	4.6	48.9	3,418	16.7	57.3	705	23.7	1,411	21.8
	2005	3,774	5.1	47.6	3,018	14.8	58.1	1,771	17.3	2,111	20.7
	2004	3,390	5.3	49.4	2,549	54.1	26.0	1,172	17.7	1,156	21.6
	2003	3,372	7.3	46.0	2,494	65.0	14.9	707	25.7	861	25.8
	2002	3,479	7.1	45.4	2,767	67.5	14.3	737	27.4	930	28.0
	2001	3,429	7.1	45.6	2,803	67.8	14.0	771	30.5	956	28.9
Black, Not Hispanic	2010	163	2.5	54.0	128	26.6	43.8	22	*	25	*
	2009	158	3.2	53.2	130	24.6	44.6	21	*	32	*
	2008	154	1.3	51.3	126	21.4	51.6	38	*	42	*
	2007	151	2.6	52.3	131	28.2	41.2	29	*	25	*
	2006	144	1.4	53.5	132	18.9	50.8	32	*	38	*
	2005	127	5.5	40.2	92	*	*	69	*	73	*
	2004	85	*	*	66	*	*	44	*	26	*
	2003	47	*	*	31	*	*	13	*	10	*
	2002	33	*	*	30	*	*	6	*	7	*
	2001	34	*	*	32	*	*	6	*	10	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2010	335	3.0	50.7	280	19.3	51.4	37	*	54	*
	2009	332	4.2	45.5	247	16.6	50.2	45	*	48	*
	2008	293	4.1	45.1	234	20.1	55.6	34	*	72	*
	2007	318	2.8	51.6	272	17.3	58.1	40	*	49	*
	2006	229	6.1	44.5	197	19.8	48.2	53	*	48	*
	2005	42	*	*	34	*	*	26	*	18	*
	2004	130	8.5	36.9	91	*	*	18	*	10	*
	2003	114	4.4	48.2	78	*	*	24	*	10	*
	2002	67	*	*	61	*	*	12	*	18	*
	2001	56	*	*	49	*	*	5	*	12	*
American Indian/Alaska Native	2010	35	*	*	27	*	*	5	*	8	*
	2009	35	*	*	30	*	*	6	*	12	*
	2008	31	*	*	27	*	*	3	*	14	*
	2007	32	*	*	26	*	*	2	*	5	*
	2006	18	*	*	17	*	*	2	*	7	*
	2005	4	*	*	3	*	*	2	*	2	*
	2004	3	*	*	2	*	*	2	*	0	*
	2003	1	*	*	0	*	*	0	*	0	*
	2002	2	*	*	1	*	*	1	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2010	109	14.7	24.8	88	*	*	20	*	21	*
	2009	104	8.7	28.8	85	*	*	15	*	27	*
	2008	76	*	*	64	*	*	14	*	32	*
	2007	80	*	*	67	*	*	20	*	26	*
	2006	68	*	*	65	*	*	16	*	29	*
	2005	37	*	*	33	*	*	21	*	36	*
	2004	31	*	*	20	*	*	9	*	11	*
	2003	28	*	*	17	*	*	16	*	7	*
	2002	20	*	*	19	*	*	4	*	2	*
	2001	18	*	*	15	*	*	7	*	2	*
Multiple Races	2010	26	*	*	24	*	*	4	*	1	*
	2009	15	*	*	12	*	*	0	*	5	*
	2008	14	*	*	13	*	*	4	*	7	*
	2007	21	*	*	18	*	*	3	*	7	*
	2006	18	*	*	13	*	*	4	*	6	*
	2005	1	*	*	1	*	*	1	*	1	*
	2004	0	*	*	0	*	*	0	*	0	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
All Other/Unknown	2010	0	*	*	0	*	*	0	*	0	*
	2009	0	*	*	0	*	*	1	*	3	*
	2008	0	*	*	0	*	*	0	*	0	*
	2007	0	*	*	0	*	*	0	*	0	*
	2006	2	*	*	0	*	*	0	*	0	*
	2005	148	5.4	39.2	112	16.1	48.2	109	22.0	37	*
	2004	68	*	*	47	*	*	51	*	16	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*
Total All Race/Ethnic Groups	2010	4,894	5.3	51.3	4,094	18.1	54.7	734	26.3	983	22.8
	2009	5,030	4.8	51.7	4,044	17.4	54.8	762	29.4	1,157	23.0
	2008	4,674	4.8	48.5	3,951	16.6	56.2	747	27.7	1,898	21.3
	2007	4,493	4.7	48.5	3,915	17.1	56.3	740	29.1	1,397	21.9
	2006	4,400	4.6	48.4	3,842	17.2	56.0	812	25.2	1,539	22.6
	2005	4,133	5.2	46.9	3,293	15.2	57.0	1,999	18.4	2,278	21.7
	2004	3,707	5.4	48.4	2,775	54.1	25.6	1,296	18.5	1,219	22.0
	2003	3,562	7.2	45.8	2,620	65.3	14.7	760	26.3	888	26.0
	2002	3,601	7.1	45.3	2,878	68.2	14.0	760	27.2	957	27.9
	2001	3,537	7.0	45.7	2,899	67.8	14.0	789	30.7	980	29.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2010	3,239	90.6	2.2	4,344	37.3
	2009	3,221	91.0	1.3	4,529	34.2
	2008	2,648	90.6	1.4	3,272	39.1
	2007	2,665	89.0	2.3	3,200	37.4
	2006	2,316	89.0	1.1	3,648	31.3
	2005	3,398	92.3	0.7	3,370	38.5
	2004	3,296	91.5	0.9	3,530	41.7
	2003	3,144	91.1	0.9	3,516	40.3
	2002	3,638	43.0	53.1	3,651	41.1
	2001	4,119	0.2	99.7	3,591	38.0
Black, Not Hispanic	2010	123	79.7	5.7	171	28.1
	2009	117	85.5	3.4	164	29.9
	2008	96	*	*	129	31.0
	2007	98	*	*	130	24.6
	2006	77	*	*	130	22.3
	2005	125	91.2	2.4	127	27.6
	2004	98	*	*	90	*
	2003	49	*	*	52	*
	2002	39	*	*	36	*
	2001	46	*	*	38	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2010	257	86.4	3.9	340	43.2
	2009	212	89.6	2.4	345	36.8
	2008	162	89.5	1.9	244	41.0
	2007	188	87.8	4.8	273	41.0
	2006	145	86.9	1.4	210	22.4
	2005	32	*	*	37	*
	2004	119	94.1	1.7	129	29.5
	2003	105	86.7	0.0	118	29.7
	2002	65	*	*	68	*
	2001	65	*	*	59	*
American Indian/Alaska Native	2010	22	*	*	35	*
	2009	26	*	*	37	*
	2008	22	*	*	21	*
	2007	22	*	*	29	*
	2006	13	*	*	18	*
	2005	4	*	*	4	*
	2004	2	*	*	3	*
	2003	0	*	*	1	*
	2002	1	*	*	2	*
	2001	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2010	79	*	*	110	25.5
	2009	66	*	*	108	22.2
	2008	43	*	*	54	*
	2007	47	*	*	64	*
	2006	43	*	*	62	*
	2005	38	*	*	34	*
	2004	40	*	*	33	*
	2003	27	*	*	31	*
	2002	23	*	*	21	*
	2001	21	*	*	19	*
Multiple Races	2010	24	*	*	27	*
	2009	8	*	*	15	*
	2008	10	*	*	12	*
	2007	15	*	*	18	*
	2006	9	*	*	17	*
	2005	1	*	*	1	*
	2004	0	*	*	0	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
All Other/Unknown	2010	0	*	*	0	*
	2009	0	*	*	4	*
	2008	0	*	*	0	*
	2007	0	*	*	0	*
	2006	1	*	*	5	*
	2005	120	95.0	0.0	148	29.1
	2004	56	*	*	71	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*
Total All Race/Ethnic Groups	2010	3,744	89.8	2.4	5,027	37.1
	2009	3,650	90.5	1.4	5,202	33.9
	2008	2,981	90.3	1.5	3,732	38.8
	2007	3,035	88.8	2.6	3,714	36.9
	2006	2,604	88.8	1.0	4,090	30.2
	2005	3,718	92.2	0.7	3,721	37.2
	2004	3,611	91.5	0.9	3,856	40.5
	2003	3,325	90.9	0.8	3,718	39.7
	2002	3,766	43.1	53.1	3,778	40.9
	2001	4,251	0.2	99.7	3,707	37.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes ⁽³⁾		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2010	4,060	44.8	3,022	27.7	1,078	29.2	3,394	24.2	3,119	17.3
	2009	3,980	47.0	2,908	28.0	1,029	26.8	3,231	34.0	3,002	29.8
	2008	3,341	46.7	2,283	30.7	814	21.0	2,594	39.9	2,299	38.1
	2007	3,133	48.4	1,881	33.7	700	16.4	2,505	44.5	1,946	41.1
	2006	2,934	47.9	1,963	31.0	607	22.1	1,113	44.3	1,932	42.0
	2005	2,822	52.3	2,783	32.1	958	25.3	4	*	276	35.1
	2004	2,807	55.1	2,616	35.6	967	22.3	0	*	0	*
	2003	2,826	50.8	2,440	32.4	812	19.0	0	*	0	*
	2002	2,984	51.2	2,653	33.1	882	20.7	0	*	0	*
	2001	3,083	50.1	2,986	35.4	1,012	19.8	0	*	0	*
Black, Not Hispanic	2010	164	21.3	125	10.4	19	*	140	6.4	122	6.6
	2009	157	14.0	122	4.9	13	*	142	10.6	125	9.6
	2008	144	14.6	111	4.5	11	*	116	11.2	109	9.2
	2007	148	12.8	108	9.3	10	*	122	18.9	104	17.3
	2006	120	11.7	104	3.8	6	*	48	*	97	*
	2005	116	18.1	110	9.1	10	*	0	*	6	*
	2004	77	*	77	*	6	*	0	*	0	*
	2003	42	*	40	*	4	*	0	*	0	*
	2002	30	*	33	*	6	*	0	*	0	*
	2001	29	*	34	*	10	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes ⁽³⁾		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2010	325	17.5	243	6.6	32	*	296	11.1	251	9.2
	2009	313	18.5	214	9.3	33	*	271	21.4	211	21.8
	2008	266	17.3	195	5.1	18	*	220	16.8	196	14.8
	2007	285	10.9	216	4.2	17	*	251	16.3	219	12.8
	2006	188	18.6	162	6.2	17	*	120	20.8	162	19.1
	2005	37	*	31	*	3	*	0	*	4	*
	2004	116	13.8	100	3.0	9	*	0	*	0	*
	2003	104	9.6	99	*	4	*	0	*	0	*
	2002	65	*	62	*	8	*	0	*	0	*
	2001	51	*	45	*	2	*	0	*	0	*
American Indian/Alaska Native	2010	32	*	23	*	8	*	24	*	24	*
	2009	30	*	25	*	9	*	27	*	26	*
	2008	29	*	23	*	5	*	17	*	22	*
	2007	24	*	15	*	8	*	23	*	16	*
	2006	15	*	12	*	1	*	7	*	12	*
	2005	4	*	4	*	1	*	0	*	0	*
	2004	2	*	1	*	1	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	2	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
New Hampshire
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2010	105	3.8	83	*	3	*	89	*	82	*
	2009	90	*	69	*	1	*	64	*	69	*
	2008	66	*	37	*	2	*	40	*	37	*
	2007	62	*	34	*	1	*	47	*	38	*
	2006	56	*	40	*	1	*	19	*	40	*
	2005	22	*	26	*	1	*	1	*	11	*
	2004	22	*	22	*	0	*	0	*	0	*
	2003	20	*	17	*	1	*	0	*	0	*
	2002	18	*	16	*	0	*	0	*	0	*
	2001	14	*	15	*	2	*	0	*	0	*
Multiple Races	2010	27	*	21	*	5	*	24	*	22	*
	2009	13	*	9	*	1	*	8	*	10	*
	2008	11	*	10	*	3	*	9	*	9	*
	2007	19	*	13	*	5	*	17	*	13	*
	2006	8	*	5	*	3	*	4	*	5	*
	2005	1	*	1	*	0	*	0	*	0	*
	2004	0	*	0	*	0	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2010 Pregnancy Nutrition Surveillance (1)
New Hampshire
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
All Other/Unknown	2010	0	*	0	*	0	*	0	*	0	*
	2009	3	*	2	*	0	*	1	*	3	*
	2008	0	*	0	*	0	*	0	*	0	*
	2007	0	*	0	*	0	*	0	*	0	*
	2006	2	*	0	*	0	*	0	*	0	*
	2005	119	16.0	90	*	13	*	0	*	0	*
	2004	62	*	39	*	2	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*
Total All Race/Ethnic Groups	2010	4,713	41.1	3,517	25.0	1,145	29.8	3,967	22.2	3,620	16.1
	2009	4,586	43.2	3,349	25.4	1,086	27.3	3,744	31.8	3,446	28.1
	2008	3,857	42.8	2,659	27.2	853	22.0	2,996	36.9	2,672	34.8
	2007	3,671	43.4	2,267	29.2	741	17.0	2,965	40.7	2,336	36.9
	2006	3,323	44.1	2,286	27.4	635	22.7	1,311	40.8	2,248	38.6
	2005	3,121	48.9	3,045	29.9	986	25.7	5	*	297	34.7
	2004	3,086	51.3	2,855	33.0	985	22.6	0	*	0	*
	2003	2,992	48.7	2,596	30.8	821	19.1	0	*	0	*
	2002	3,099	50.0	2,764	32.1	896	21.0	0	*	0	*
	2001	3,177	49.2	3,080	34.7	1,026	19.8	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾

New Hampshire
Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)	Number	%	Number	%	Number	%
		Number	%	%						
White, Not Hispanic	2010	2,164	7.3	9.9	3,635	6.7	2,012	4.0	3,621	72.5
	2009	2,987	7.3	10.3	3,667	6.9	2,774	4.1	3,649	70.9
	2008	3,260	6.1	10.2	2,665	7.5	2,351	2.0	3,190	72.2
	2007	3,208	6.7	10.7	2,740	8.0	2,365	2.8	3,058	63.9
	2006	3,044	6.4	8.7	3,140	7.3	2,578	3.6	3,194	63.5
	2005	1,780	5.7	11.1	1,376	6.3	1,191	2.9	1,930	67.9
	2004	3,070	6.6	10.5	2,351	9.9	2,107	1.6	3,125	60.2
	2003	3,944	7.5	11.4	3,054	9.7	2,739	2.6	4,025	60.0
	2002	4,104	6.9	10.9	3,154	8.9	2,853	2.8	4,172	60.1
2001	4,204	8.2	10.5	3,155	10.8	2,786	2.9	4,287	53.6	
Black, Not Hispanic	2010	66	*	*	136	5.9	60	*	132	85.6
	2009	112	7.1	7.1	135	11.1	98	*	135	86.7
	2008	125	6.4	8.8	103	7.8	90	*	129	92.2
	2007	124	8.9	7.3	112	15.2	86	*	130	87.7
	2006	106	11.3	13.2	117	8.5	85	*	128	87.5
	2005	59	*	*	54	*	40	*	76	*
	2004	75	*	*	58	*	52	*	79	*
	2003	66	*	*	49	*	41	*	68	*
	2002	40	*	*	34	*	26	*	43	*
2001	46	*	*	37	*	30	*	48	*	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾

New Hampshire
Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)							
		Number	%	%	Number	%	Number	%	Number	%
Hispanic	2010	178	11.8	5.1	287	10.1	155	7.7	286	81.5
	2009	222	6.8	6.8	257	7.8	204	2.9	255	79.6
	2008	221	8.1	7.2	186	7.5	162	4.3	228	89.9
	2007	261	6.1	11.1	234	8.1	203	3.4	248	87.9
	2006	185	10.8	10.8	179	10.6	147	6.8	186	74.7
	2005	9	*	*	5	*	4	*	10	*
	2004	145	4.8	9.0	117	5.1	110	0.0	150	80.7
	2003	143	8.4	8.4	107	9.3	96	*	145	80.7
	2002	69	*	*	63	*	60	*	69	*
	2001	60	*	*	50	*	45	*	65	*
American Indian/Alaska Native	2010	15	*	*	27	*	15	*	27	*
	2009	21	*	*	31	*	17	*	30	*
	2008	22	*	*	16	*	12	*	27	*
	2007	28	*	*	23	*	20	*	22	*
	2006	16	*	*	16	*	14	*	16	*
	2005	2	*	*	2	*	2	*	2	*
	2004	2	*	*	2	*	2	*	2	*
	2003	0	*	*	0	*	0	*	0	*
	2002	2	*	*	2	*	1	*	2	*
	2001	0	*	*	0	*	0	*	0	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾

New Hampshire
Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)							
		Number	%	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2010	67	*	*	93	*	63	*	92	*
	2009	72	*	*	89	*	68	*	86	*
	2008	56	*	*	42	*	36	*	58	*
	2007	58	*	*	51	*	44	*	62	*
	2006	52	*	*	59	*	45	*	62	*
	2005	18	*	*	21	*	13	*	26	*
	2004	53	*	*	31	*	27	*	53	*
	2003	38	*	*	25	*	24	*	39	*
	2002	25	*	*	21	*	19	*	25	*
	2001	21	*	*	18	*	16	*	22	*
Multiple Races	2010	15	*	*	25	*	13	*	25	*
	2009	9	*	*	12	*	8	*	12	*
	2008	13	*	*	11	*	11	*	12	*
	2007	18	*	*	15	*	12	*	16	*
	2006	10	*	*	12	*	8	*	12	*
	2005	0	*	*	0	*	0	*	0	*
	2004	0	*	*	0	*	0	*	0	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾

New Hampshire
Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
All Other/Unknown	2010	0	*	*	0	*	0	*	0	*
	2009	3	*	*	3	*	3	*	3	*
	2008	0	*	*	0	*	0	*	0	*
	2007	0	*	*	0	*	0	*	0	*
	2006	1	*	*	2	*	1	*	2	*
	2005	19	*	*	15	*	14	*	20	*
	2004	0	*	*	0	*	0	*	0	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*
Total All Race/Ethnic Groups	2010	2,505	7.7	9.3	4,203	6.9	2,318	4.3	4,183	73.9
	2009	3,426	7.3	10.0	4,194	7.1	3,172	4.0	4,170	72.4
	2008	3,697	6.2	9.9	3,023	7.5	2,662	2.1	3,644	74.3
	2007	3,697	6.7	10.4	3,175	8.2	2,730	2.9	3,536	66.7
	2006	3,414	6.7	8.9	3,525	7.5	2,878	3.8	3,600	65.3
	2005	1,887	6.0	10.7	1,473	6.3	1,264	3.2	2,064	69.1
	2004	3,345	6.6	10.2	2,559	9.7	2,298	1.7	3,409	61.8
	2003	4,191	7.5	11.2	3,235	9.7	2,900	2.7	4,277	60.8
	2002	4,240	6.8	10.9	3,274	9.0	2,959	2.8	4,311	60.3
	2001	4,331	8.2	10.6	3,260	10.8	2,877	2.9	4,422	54.0

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

