

THIS IS AN OFFICAL NH DHHS HEALTH ALERT


Distributed by the NH Health Alert Network
Health.Alert@nh.gov
April 16, 2019; 1000 EDT (2:00 PM EDT)
NH-HAN 20190416



Hepatitis A Virus (HAV) Outbreak Accelerating Update #1

Key Points and Recommendations:

1. The hepatitis A virus (HAV) outbreak in New Hampshire is worsening. There have been 79 new acute hepatitis A virus (HAV) infections diagnosed since the outbreak began in November 2018, with increasing numbers every month (historically, NH has 6-7 cases *per year*, or <1 case *per month*).



2. Most people infected with hepatitis A in this outbreak have been hospitalized (63%), and one person has died. This individual had a co-infection with hepatitis C and died from fulminant liver failure after developing an acute HAV infection.
3. Vaccinate the following groups as soon as possible because they are currently at highest risk for acquiring HAV infection or developing serious complications from HAV infection:
 - People who use drugs (injection or non-injection)
 - People experiencing homelessness
 - Gay and bisexual men (i.e., men who have sex with men)
 - People who are, or were recently, incarcerated
 - People with chronic liver disease, including cirrhosis, hepatitis B, or hepatitis C
4. All healthcare providers at all levels of patient care (including outpatient offices, emergency departments, urgent care facilities, inpatient wards, jails/prisons, etc.) should screen patients for and vaccinate those with these risk factors. Checking bloodwork for immunity (either from prior infection or prior vaccination) is not recommended as this will delay vaccination.
5. The following individuals should also be vaccinated:

- All children beginning at age 1 year should be vaccinated as part of the routine childhood immunization recommendations.
 - Any person wishing to obtain immunity can/should be vaccinated.
6. Healthcare workers and first responders are not at increased risk for HAV infection, so do not prioritize them for vaccination as part of control of this outbreak. However, encourage anybody who wants to be protected to talk to their primary care providers or seek vaccination at a commercial pharmacy; vaccination is covered by insurance.
 7. Test all patients who inject drugs for HIV, hepatitis C, and hepatitis B infection.
 8. If you need assistance obtaining and administering vaccines, contact the New Hampshire Immunization Program within the NH Division of Public Health Services (DPHS) at (603) 271-4482. Community-based organizations that serve individuals at highest risk should reach out to their Regional Public Health Network representatives listed here to coordinate vaccination opportunities for clients: <https://www.dhhs.nh.gov/dphs/rphn/documents/rphncontactlist.pdf>.
 9. Report any suspect or confirmed case of hepatitis A promptly to the NH DPHS (within 24 hours) by calling 603-271-4496 (after hours 603-271-5300).
 10. Please also review the following information:
 - The attached CDC HAN
 - Prior NH DHHS HAN (February 2019): <https://www.dhhs.nh.gov/dphs/cdcs/alerts/documents/hepatitis-a-2019.pdf>
 - NH DHHS Healthcare provider FAQs (February 2019): <https://www.dhhs.nh.gov/dphs/cdcs/hepatitisa/documents/hepa-faqs-hcp.pdf>

Situation:

There have been 79 new cases of acute hepatitis A virus (HAV) infections since the outbreak began in November 2018, with increasing numbers every month. 33 of these hepatitis A diagnoses were in March alone. 63% of infected individuals have required hospitalization, and there has been one death in an individual with hepatitis C co-infection who developed fulminant liver failure. 55% are males, and the average age is 37.9 (Range: 20 - 75). Individuals have been diagnosed in most New Hampshire counties highlighting this as a state-wide problem. However, a majority of individuals have been from Hillsborough (36), Strafford (17), Rockingham (10), and Merrimack (10) Counties. A majority of infections have been in people reporting injectable and non-injectable drug use (67%) and people experiencing homelessness (35%). A number of individuals diagnosed with HAV have been incarcerated (11%).

To facilitate vaccination of high-risk individuals, our Regional Public Health Networks have been mobilized to target vaccination to individuals at highest risk of HAV infection. Any facility serving high-risk individuals (e.g., medication assisted therapy clinics, homeless shelters, transitional housing, soup kitchens, etc.) that does not have the ability to deliver hepatitis A vaccination should reach out to the Emergency Preparedness Coordinator in their Regional Public Health Network. Contact information can be found here: <https://www.dhhs.nh.gov/dphs/rphn/documents/rphncontactlist.pdf>.

Please visit our NH DPHS hepatitis A virus resource page for routine updates, educational materials and other information:

<https://www.dhhs.nh.gov/dphs/cdcs/hepatitisa/hepa-nh.htm>.

- ▶ For any questions regarding the contents of this message, please contact NH DHHS, DPHS, Bureau of Infectious Disease Control at 603-271-4496 (after hours 603-271-5300).
- ▶ To change your contact information in the NH Health Alert Network, contact Adnela Alic at 603-271-7499 or Adnela.Alic@dhhs.nh.gov.

Status: Actual
Message Type: Alert
Severity: Moderate
Sensitivity: Not Sensitive
Message Identifier: NHHAN 20190416
Delivery Time: 12 hours
Acknowledgement: No
Distribution Method: Email, Fax
Distributed to: Physicians, Physician Assistants, Practice Managers, Infection Control Practitioners, Infectious Disease Specialists, Community Health Centers, Hospital CEOs, Hospital Emergency Departments, Nurses, Long-Term Care Facilities, NHHA, Laboratory Response Network, Manchester Health Department, Nashua Health Department, Public Health Network, DHHS Outbreak Team, DPHS Investigation Team, DPHS Management Team, Northeast State Epidemiologists
From: Benjamin P. Chan, MD, MPH – State Epidemiologist
Originating Agency: NH Department of Health and Human Services, Division of Public Health Services

Attachments: CDC HAN 418

Follow us on Twitter @NHIDWatch or Facebook @NHBIDC

This is an official **CDC HEALTH UPDATE**

Distributed via the CDC Health Alert Network
March 25, 2019, 13:30 ET (1:30 PM ET)
CDCHAN-00418

Update: Widespread Outbreaks of Hepatitis A among People Who Use Drugs and People Experiencing Homelessness across the United States

Summary

Multiple states across the country have reported outbreaks of hepatitis A, primarily among people who use drugs and people experiencing homelessness. Since the hepatitis A outbreaks were first identified in 2016, more than 15,000 cases, 8,500 (57%) hospitalizations, and 140 deaths as a result of hepatitis A virus (HAV) infection have been reported. This Health Alert Network (HAN) update recommends that public health departments, healthcare facilities, and partners and programs providing services to affected populations vaccinate at-risk groups against hepatitis A, applying the updated recommendations of the Advisory Committee on Immunization Practices (ACIP).

This is an update to the Health Alert Network (HAN) advisory released on June 11, 2018 titled *Outbreak of Hepatitis A Virus (HAV) Infections among Persons Who Use Drugs and Persons Experiencing Homelessness* (<https://emergency.cdc.gov/han/han00412.asp>).

Background

Multiple states across the country have reported outbreaks of hepatitis A, primarily among people who use drugs and people experiencing homelessness. Since these outbreaks were first identified in 2016, more than 15,000 cases and 8,500 (57%) hospitalizations have been reported. Hospitalization rates have been higher than typically associated with HAV infection.^{1, 2} Severe complications have also been reported, sometimes leading to liver transplantation or death; at least 140 deaths have occurred nationwide.

HAV is highly transmissible from person-to-person. States experiencing large-scale outbreaks have reported widespread transmission soon after their jurisdictions first recognized hepatitis A cases among populations being affected by these outbreaks. For many states, this has resulted in an unprecedented number of hepatitis A cases among unvaccinated adults since hepatitis A vaccine became available in 1996, and has led to prolonged community outbreaks that have been challenging and costly to control.

CDC recommends that public health departments, healthcare providers, and other partners serving affected populations launch a rapid and effective public health response with the following strategies.

Recommendations

Offer Vaccination to the Following Groups to Prevent or Control an Outbreak

The best way to prevent HAV infection is through vaccination with the hepatitis A vaccine. The following groups are at highest risk for acquiring HAV infection or developing serious complications from HAV infection in these outbreaks and should be offered the hepatitis A vaccine:

- **People who use drugs (injection or non-injection)**
- **People experiencing homelessness**
- **Men who have sex with men (MSM)**
- **People who are, or were recently, incarcerated**
- **People with chronic liver disease, including cirrhosis, hepatitis B, or hepatitis C**

One dose of single-antigen hepatitis A vaccine has been shown to control outbreaks of hepatitis A and provides up to 95% seroprotection in healthy individuals for up to 11 years.^{3,4}

Pre-vaccination serologic testing is not required to administer hepatitis A vaccine. Vaccinations should not be postponed if vaccination history cannot be obtained or records are unavailable.

New ACIP Recommendations since the June 2018 HAN00412

<https://emergency.cdc.gov/han/han00412.asp>

1. As of November 2, 2018, ACIP recommends hepatitis A vaccine for post-exposure prophylaxis (PEP) for people 12 months of age and older. Providers may also administer immunoglobulin to adults older than 40 years of age, if indicated, and persons who are immunocompromised or have chronic liver disease.⁵
2. As of February 15, 2019, ACIP recommends hepatitis A vaccination for people experiencing homelessness.⁶

Health Departments

Outreach

1. Identify venues serving populations at-risk for HAV infection, including correctional facilities, syringe service programs, medication-assisted treatment (MAT) facilities, substance use disorder treatment facilities, homeless shelters, emergency departments, and sexually transmitted diseases (STD) clinics. Where ongoing relationships with these facilities and services providers do not exist, engage with partners serving these populations to promote education and vaccination efforts.
2. Employ novel approaches to improve vaccine delivery to hard-to-reach populations (e.g., Point of Dispensing sites (PODs), mobile outreach teams).
3. Include hepatitis A vaccination for ACIP-recommended risk groups in routine clinical services to increase vaccination coverage.
4. Engage multidisciplinary stakeholders (e.g., viral hepatitis or communicable disease experts, epidemiologists, immunization program staff, emergency preparedness staff, disease investigator specialists, health educators, behavioral scientists, harm reduction partners), which is critical for effective response efforts.

Case investigation, contact tracing, and outbreak response monitoring

1. Implement the new 2019 Acute Hepatitis A Case Definition from the Council of State and Territorial Epidemiologists (CSTE).⁷
2. Follow established procedures to interview cases and perform contact tracing for all new hepatitis A diagnoses.
3. Provide or encourage PEP of previously unvaccinated contacts as soon as possible, within 2 weeks after exposure.⁵
4. Track vaccine delivery and administration to at-risk populations to monitor the outbreak response efforts and improve vaccine coverage among at-risk populations.

Preventing outbreaks

1. States that are not currently experiencing hepatitis A outbreaks should remain vigilant for hepatitis A cases and proactively develop and implement prevention strategies. This includes outreach to and vaccination of ACIP-recommended risk groups, particularly people who use drugs (injection or non-injection), people experiencing homelessness, MSM, and people with chronic liver disease.^{6,8}
2. As soon as a hepatitis A case is identified in the at-risk populations, states should rapidly mobilize a response to mitigate the threat of HAV transmission.

Healthcare Providers

1. Screen patients for risk factors (e.g., drug use, homelessness, incarceration, MSM, and chronic liver disease).
2. Recommend and administer hepatitis A vaccine to at-risk patients, regardless of the original presenting complaint or the type of clinical facility. In particular, the emergency department may be an individual's only interaction with the healthcare system and is an important opportunity for prevention.
3. Record immunizations in the state immunization information system (registry).
4. Consider hepatitis A as a diagnosis in anyone with jaundice or clinically compatible symptoms.
5. Rapidly report all persons diagnosed with hepatitis A to the health department to ensure timely case investigation and follow-up of contacts.

For More Information

1. CSTE's 2019 Acute Hepatitis A Case Definition. <https://wwwn.cdc.gov/nndss/conditions/hepatitis-a-acute/case-definition/2019/>
2. MMWR. *Hepatitis A Virus Outbreaks Associated with Drug Use and Homelessness – California, Kentucky, Michigan, and Utah, 2017*. <https://www.cdc.gov/mmwr/volumes/67/wr/mm6743a3.htm>
3. CDC's Hepatitis A Outbreak website. <https://www.cdc.gov/hepatitis/outbreaks/2017March-HepatitisA.htm>
4. Outbreak specific considerations for hepatitis A vaccine administration. <https://www.cdc.gov/hepatitis/outbreaks/InterimOutbreakGuidance-HAV-VaccineAdmin.htm>
5. CDC's Hepatitis A Virus website. <https://www.cdc.gov/hepatitis/hav/index.htm>
6. Viral Hepatitis Surveillance – United States, 2016. <https://www.cdc.gov/hepatitis/statistics/2016surveillance/pdfs/2016HepSurveillanceRpt.pdf>
7. Hepatitis A General Information Fact Sheet. <https://www.cdc.gov/hepatitis/hav/pdfs/hepageneralfactsheet.pdf>
8. CDC's The Pink Book. Chapter 9: Hepatitis A. <https://www.cdc.gov/vaccines/pubs/pinkbook/downloads/hepa.pdf>

References

1. Ly K and Klevens RM. Trends in disease and complications of hepatitis A virus infection in the United States, 1999-2011: A new concern for adults. *J Infect Dis* 2015;212:176-182.
2. CDC. Viral hepatitis surveillance, United States, 2016. Atlanta, GA: CDC. <https://www.cdc.gov/hepatitis/statistics/2016surveillance/pdfs/2016HepSurveillanceRpt.pdf>
3. McMahon BJ, Beller M, Williams J, Schloss M, Tanttila H, Bulkow L. A program to control an outbreak of hepatitis A in Alaska by using an inactivated hepatitis A vaccine. *Arch Pediatr Adolesc Med* 1996;150(7):733-739.
4. Ott JJ, Wiersma ST. Single-dose administration of inactivated hepatitis A vaccination in the context of hepatitis A vaccine recommendations. *Int J Infect Dis* 2013;17(11):e939-944.
5. Nelson NP, Link-Gelles R, Hofmeister MG, et al. Update: Recommendations of the Advisory Committee on Immunization Practices for use of hepatitis A vaccine for post exposure prophylaxis and for preexposure prophylaxis for international travel. *MMWR Morb Mortal Wkly Rep* 2018;67(43):1216-1220. https://www.cdc.gov/mmwr/volumes/67/wr/mm6743a5.htm?s_cid=mm6743a5_w
6. Doshani M, Weng M, Moore K, Romero J, Nelson NP. Recommendations of the Advisory Committee on Immunization Practices for Use of Hepatitis A Vaccine for Persons Experiencing Homelessness. *MMWR Morb Mortal Wkly Rep* 2019;68:153-156. <https://www.cdc.gov/mmwr/volumes/68/wr/mm6806a6.htm>
7. National Notifiable Diseases Surveillance System. Hepatitis A, Acute 2019 Case Definition. <https://wwwn.cdc.gov/nndss/conditions/hepatitis-a-acute/case-definition/2019/>
8. Fiore AE, Wasley A, Bell BP; Advisory Committee on Immunization Practices (ACIP). Prevention of hepatitis A through active or passive immunization: Recommendations of the Advisory

Committee on Immunization Practices (ACIP). *MMWR Recomm Rep* 2006;55(No. RR-7).
<https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5507a1.htm>

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages:

Health Alert Requires immediate action or attention; highest level of importance
Health Advisory May not require immediate action; provides important information for a specific incident or situation
Health Update Unlikely to require immediate action; provides updated information regarding an incident or situation
HAN Info Service Does not require immediate action; provides general public health information

##This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, epidemiologists, HAN coordinators, and clinician organizations##