

**STATE OF NEW HAMPSHIRE
HEALTHCARE-ASSOCIATED INFECTIONS
2018 HOSPITAL REPORT**

September 23, 2019

*New Hampshire Department of Health and Human Services
Division of Public Health Services*

TABLE OF CONTENTS

LIST OF DATA TABLES	4
LIST OF FIGURES	5
INDIVIDUAL HOSPITAL REPORTS	6
ABBREVIATIONS USED IN THIS DOCUMENT	7
CONTRIBUTORS AND ACKNOWLEDGMENTS	8
EXECUTIVE SUMMARY	9
I. INTRODUCTION	12
A. Purpose	12
B. Audience.....	12
C. How to Use this Document	12
D. Background on Healthcare-Associated Infections.....	13
E. New Hampshire Healthcare-Associated Infections Program.....	13
F. State of New Hampshire Healthcare-Associated Infections Plan	14
G. Overview of Healthcare-Associated Infections Prevention Efforts	14
H. Healthcare-Associated Infections Technical Advisory Workgroup.....	15
II. SURVEILLANCE METHODS	18
A. 2018 Healthcare-Associated Infections Reporting Requirements for New Hampshire Hospitals	18
B. Selection of Reporting Requirements	18
C. Accuracy of Reported Healthcare-Associated Infections Surveillance Data	20
D. National Healthcare Safety Network	20
E. Comparisons with National Data	22
F. Central Line-Associated Bloodstream Infections Surveillance	22
G. Central Line Insertion Practices Monitoring	23
H. Catheter-Associated Urinary Tract Infections Surveillance	23
I. Surgical Site Infections Surveillance	24
J. Surgical Antimicrobial Prophylaxis Administration Monitoring	27
K. Influenza Vaccination Percentage Monitoring.....	27
III. STATEWIDE DATA	29
A. Statewide Standardized Infection Ratios.....	29
B. Overall Standardized Infection Ratios by Hospital.....	32

C. Central Line-Associated Bloodstream Infections	38
D. Central Line Insertion Practices	44
E. Catheter-Associated Urinary Tract Infections	50
F. Surgical Site Infections	55
G. Surgical Antimicrobial Prophylaxis Administration	70
H. Influenza Vaccination Percentages	70
IV. CONCLUSIONS	80
V. ACUTE CARE HOSPITAL REPORTS	81
APPENDIX 1: Technical Notes	140
APPENDIX 2: Influenza Vaccination Survey Questions, 2018-2019 Season	144
APPENDIX 3: Understanding the Relationship between Healthcare-Associated Infection Rates and Standardized Infection Ratio Comparison Metrics	146
APPENDIX 4: Preventing Healthcare-Associated Infections	148
APPENDIX 5: Map of New Hampshire Hospitals, 2018	153
REFERENCES	154

LIST OF DATA TABLES

Table 1. Overall statewide standardized infection ratios	31
Table 2. Overall healthcare-associated infections standardized infection ratios by hospital	33
Table 3. Overall healthcare-associated infections standardized infection ratios, comparison between 2017 and 2018	35
Table 4. Overall healthcare-associated infections standardized infection ratios by hospital, comparison between 2017 and 2018	37
Table 5. Central line-associated bloodstream infections rates,	39
Table 6. Central line-associated bloodstream infections rates in neonatal intensive care units.	40
Table 7. Central line-associated bloodstream infections standardized infection ratios	41
Table 8. Central line-associated bloodstream infections standardized infection ratios, comparison between 2017 and 2018	43
Table 9. Central line insertion practices adherence by occupation of inserter	45
Table 10. Central line insertion practices adherence percentages by hospital	46
Table 11. Central line insertion practices adherence percentages by occupation of inserter, comparison between 2017 and 2018	48
Table 12. Central line insertion practices adherence percentages by hospital, comparison between 2017 and 2018	49
Table 13. Catheter-associated urinary tract infection rates	51
Table 14. Catheter-associated urinary tract infections standardized infection ratios	52
Table 15. Catheter-associated urinary tract infections standardized infection ratios, comparison between 2017 and 2018	54
Table 16. Surgical site infections standardized infection ratios	57
Table 17. Surgical site infections standardized infection ratios, comparison between 2017 and 2018.....	59
Table 18. Coronary artery bypass graft procedure-associated surgical site infections standardized infection ratios	60
Table 19. Colon procedure-associated surgical site infections standardized infection ratios	61
Table 20. Abdominal hysterectomy procedure-associated surgical site infections standardized infection ratios	63
Table 21. Knee arthroplasty procedure-associated surgical site infections standardized infection ratios	65
Table 22. Post-discharge surveillance methods in New Hampshire hospitals, 2017-2018.....	69
Table 23. Influenza vaccination percentages for hospital healthcare personnel by hospital, 2017–18 influenza season.....	72
Table 24. Influenza vaccination percentages for hospital healthcare personnel by hospital, comparison between 2017-18 and 2018-19 influenza seasons	76
Table 25. Influenza vaccination policies for healthcare personnel by hospital, 2018-19 influenza season	78

LIST OF FIGURES

Figure 1. Types of healthcare-associated infections reported to New Hampshire Department of Health and Human Services	17
Figure 2. Overall statewide standardized infection ratios.....	32
Figure 3. Overall healthcare-associated infections standardized infection ratios	34
Figure 4. Statewide standardized infection ratios, comparison between 2017-2018	36
Figure 5. Central line-associated bloodstream infections standardized infection ratios.....	42
Figure 6. Overall central line-associated bloodstream infections standardized infection ratios by year, 2017-2018	44
Figure 7. Central line insertion practices adherence percentages by hospital	47
Figure 8. Catheter-associated urinary tract infections standardized infection ratios.....	53
Figure 9. Overall catheter-associated urinary tract infections standardized infection ratios by year, 2017-2018	55
Figure 10. Surgical site infections standardized infection ratios	58
Figure 11. Coronary artery bypass graft procedure-associated surgical site infections standardized infection ratios	60
Figure 12. Colon procedure-associated surgical site infections standardized infection ratios....	62
Figure 13. Abdominal hysterectomy procedure-associated surgical site infections standardized infection ratios	64
Figure 14. Knee arthroplasty procedure-associated surgical site infections standardized infection ratios	66
Figure 15. Overall surgical site infection standardized infection ratios by year, 2011-2017	67
Figure 16. Overall coronary artery bypass graft procedure standardized infection ratios by year, 2011-2017	67
Figure 17. Overall colon procedure standardized infection ratios by year, 2017-2018.....	68
Figure 18. Overall abdominal hysterectomy standardized infection ratios by year, 2017-2018.	68
Figure 19. Overall knee arthroplasty standardized infection ratios by year, 2017-2018.....	68
Figure 20. Statewide influenza vaccination percentages for hospital healthcare personnel by influenza season.....	71
Figure 21. Influenza vaccination percentages for hospital healthcare personnel by hospital, 2018-19 influenza season	73
Figure 22. Influenza vaccination percentages for hospital healthcare personnel by hospital, 2017-18 and 2018-19 influenza seasons	75
Figure 23. Influenza vaccination percentages for hospitals with and without vaccination policies, 2018-19 influenza season	77

INDIVIDUAL HOSPITAL REPORTS

Alice Peck Day Memorial Hospital.....	82
Androscoggin Valley Hospital.....	84
Catholic Medical Center.....	86
Cheshire Medical Center.....	88
Concord Hospital.....	90
Cottage Hospital.....	92
Dartmouth-Hitchcock Medical Center.....	94
Elliot Hospital.....	96
Exeter Hospital.....	98
Franklin Regional Hospital.....	100
Frisbie Memorial Hospital.....	102
Huggins Hospital.....	104
Lakes Region General Hospital.....	106
Littleton Regional Hospital.....	108
Monadnock Community Hospital.....	110
New London Hospital.....	112
Parkland Medical Center.....	114
Portsmouth Regional Hospital.....	116
Southern New Hampshire Medical Center.....	118
Speare Memorial Hospital.....	120
St. Joseph Hospital.....	122
The Memorial Hospital.....	124
Upper Connecticut Valley Hospital	126
Valley Regional Hospital.....	128
Weeks Medical Center.....	130
Wentworth-Douglass Hospital.....	132
Hampstead Hospital.....	134
Encompass Health Rehabilitation Hospital.....	135
New Hampshire Hospital	136
Northeast Rehabilitation Hospital at The Elliot.....	137
Northeast Rehabilitation Hospital at Pease	138
Northeast Rehabilitation Hospital, Salem.....	139
Northeast Rehabilitation Hospital, Southern New Hampshire Medical Center	140

Note: DHMC refers to Dartmouth-Hitchcock Medical Center (Mary Hitchcock Memorial Hospital). NE Rehab refers to Northeast Rehabilitation Hospital and is presented as four facilities with specified locations.

ABBREVIATIONS USED IN THIS DOCUMENT

ASA Score	American Society of Anesthesiologists (ASA) Classification of Physical Status
ASC	Ambulatory surgical center(s)
BSI	Bloodstream infection(s)
CABG	Coronary artery bypass graft procedure(s)
CAUTI	Catheter-associated urinary tract infection(s)
CBGB	NHSN operative code for coronary artery bypass graft procedure(s) with both a chest and donor site incision
CBGC	NHSN operative code for coronary artery bypass graft procedure(s) with chest incision site only
CCN	CMS Certification Number
CDC	U.S. Centers for Disease Control and Prevention
CLABSI	Central line-associated bloodstream infection(s)
CLIP	Central line insertion practices
CMS	Centers for Medicare and Medicaid Services
COLO	NHSN operative code for colon procedure(s)
Crotched Mountain	Crotched Mountain Rehabilitation Center
DHMC	Dartmouth-Hitchcock Medical Center (Mary Hitchcock Memorial Hospital)
DHHS	New Hampshire Department of Health and Human Services
HAI	Healthcare-associated infection(s)
HCP	Healthcare personnel
HICPAC	Healthcare Infection Control Practices Advisory Committee
HHS	U.S. Department of Health and Human Services
HYST	NHSN operative code for abdominal hysterectomy procedure(s)
ICU	Intensive care unit(s)
KPRO	NHSN operative code for knee arthroplasty procedure(s)
NE Rehab	Northeast Rehabilitation Hospital
NH	New Hampshire
NHHCQAC	New Hampshire Healthcare Quality Assurance Commission
NHSN	National Healthcare Safety Network
RSA	Revised Statutes Annotated
SCIP	Surgical Care Improvement Project
SIR	Standardized infection ratio(s)
SSI	Surgical site infection(s)
TAW	Healthcare-Associated Infections Technical Advisory Workgroup
VAP	Ventilator-associated pneumonia(s)

CONTRIBUTORS AND ACKNOWLEDGMENTS

The following individuals contributed to the review and analysis of data and other content provided in this report:

Trevor Bauer, MPH, Healthcare-Associated Infections Surveillance Coordinator

Yvette Perron, MPH, Healthcare Associated Infections Program Manager

Katrina E. Hansen, MPH, Chief, Infectious Disease Surveillance Section

Elizabeth R. Daly, MPH, Chief, Bureau of Infectious Disease Control

Elizabeth A. Talbot, MD, Deputy State Epidemiologist

Benjamin Chan, MD, MPH, State Epidemiologist

Renelle Gagnon, MPH, Infectious Disease Prevention Statistical Assistant

Tylor Young, GIS Analyst, Infectious Disease Surveillance Section

The HAI Program would also like to thank the Infection Prevention, Quality, and Information Technology staff at New Hampshire hospitals for collaborating to provide the data presented in this report. Finally, the HAI Program acknowledges the review, comments, input, and other program contributions provided by the members of the Healthcare-Associated Infections Technical Advisory Workgroup as listed on page 16.

For questions about this report, please contact:

New Hampshire Healthcare-Associated Infections Program

Infectious Disease Surveillance Section

Division of Public Health Services

NH Department of Health and Human Services

29 Hazen Drive, Concord, NH 03301-6504

Phone: (603) 271-4496

Email: haiprogram@dhhs.nh.gov

Website: <http://www.dhhs.nh.gov/dphs/cdcs/hai/index.htm>

EXECUTIVE SUMMARY

A healthcare-associated infection (HAI) is an infection that a patient acquires during the course of receiving treatment for another condition within a healthcare setting. An estimated 722,000 HAI and 75,000 associated deaths occurred in United States (U.S.) acute care hospitals in 2011.ⁱ During the 2006 legislative season, the New Hampshire (NH) Legislature passed a bill creating NH Revised Statutes Annotated (RSA) 151:32-35, which requires hospitals to identify, track, and report selected HAI to the NH Department of Health and Human Services (DHHS). All 26 of NH's acute care hospitals began reporting data to DHHS on two infections and three process measures in January 2009, and eight specialty hospitals reported influenza vaccination coverage. This report represents the tenth summary of HAI-related data reported by hospitals in NH.

Healthcare-Associated Infections in New Hampshire Hospitals

There were more infections than predicted in NH based on national data, however these data were not statistically significant. A total of 196 HAI were reported, representing 133 surgical site infections (SSI), 20 central line-associated bloodstream infections (CLABSI), and 43 catheter-associated urinary tract infections (CAUTI). The observed number of HAI in NH hospitals was 4% higher than predicted based on national data. There were 27% fewer CLABSI, 16% more CAUTI, and 7% more SSI than predicted. Seventeen hospitals had sufficiently robust data¹ to present hospital-specific data for overall HAI. Two hospitals observed a lower number of infections as predicted, one hospital observed a higher number of infections, and fourteen hospitals observed similar infections based on national data. The total number of infections increased in 2018 when compared to 2017; however, this difference was not statistically significant.

Central Line-Associated Bloodstream Infections

Nineteen hospitals² with intensive care units (ICU) reported CLABSI data from their ICU. Data were robust enough for 16 hospitals to present data for 20 individual ICU in this report. All 20 ICU experienced similar rates of CLABSI when compared with national data. The total number of CLABSI reported decreased in 2018 compared to 2017; however, this difference was not statistically significant.

Central Line Insertion Practices

The hospitals² with ICU reported information on central line insertion practices (CLIP) for central lines inserted in their ICU. Statewide adherence to the four infection prevention practices during central line insertions was 98.3%. Compared with attending physicians, intern/residents less frequently adhered to the four infection prevention practices during central line insertions (98.2% versus 95.6%). Data were sufficiently robust for 11 hospitals to present hospital-specific data in this report. All eleven hospitals reported CLIP adherence percentages that were similar to the State percentage. In 2018, the statewide adherence percentage for CLIP decreased from 2017; however, this decrease was not statistically significant. Of the 11 hospitals for which

¹ Data are not shown for facilities with less than one predicted infection, fewer than 50 central lines or catheter days, and fewer than 20 central line insertions performed.

² Of the 34 hospitals licensed in 2018, only 21 hospitals had ICU in which to monitor CAUTI, CLABSI, and CLIP.

availability of data allowed a comparison between the two years, 10 hospitals' CLIP adherence was similar and one hospital was lower in 2018 compared to 2017.

Catheter-Associated Urinary Tract Infections

Data were sufficiently robust enough for 19 hospitals to present CAUTI data for 23 individual ICU in this report. All twenty-two ICU experienced similar rates of CAUTI and one medical ICU was lower in comparison with national rates. The total number of CAUTI reported in 2018 was lower when compared to 2017; however, this decrease was not statistically significant.

Surgical Site Infections

Twenty-four³ acute care hospitals reported SSI data for four surgical procedures.

- Coronary Artery Bypass Graft (CABG) procedures: Four acute care hospitals performed CABG, and data were sufficiently robust for three hospitals to present hospital-specific data in this report. All three hospitals reported a similar number of CABG procedure-associated SSI compared with national data. Overall, there were lower number of CABG SSI than predicted based on national data.
- Colon (COLO) procedures: Twenty-two acute care hospitals performed the procedure, and data were sufficiently robust for 13 hospitals to present hospital-specific data in this report. Twelve hospitals reported a similar number of colon procedure-associated SSI, and one hospital reported a lower number of colon procedures when compared to national data. Overall, there were a similar number of colon procedure-associated SSI as predicted based on national data.
- Abdominal Hysterectomy (HYST) procedures: Twenty-two acute care hospitals performed the procedure, and data were sufficiently robust for four hospitals to present hospital-specific data in this report. All four hospitals reported a similar number of abdominal hysterectomy procedure-associated SSI compared with national data. Overall, there were a similar number of abdominal hysterectomy procedure-associated SSI as predicted based on national data.
- Knee Arthroplasty (KPRO) procedures: Twenty-four acute care hospitals performed the procedure, and data were sufficiently robust for 9 hospitals to present hospital-specific data in this report. Eight hospitals reported a similar number of knee arthroplasty procedure-associated SSI, and one hospital had a higher number when compared with national data. Overall, there were similar knee arthroplasty-related SSI than predicted based on national data.

Surgical Antimicrobial Prophylaxis Administration

Surgical antimicrobial prophylaxis data is reported to the Centers for Medicare and Medicaid Services (CMS) through the Surgical Care Improvement Project (SCIP). In previous years, DHHS accessed these data and presented it in this report. At the time of this report's publication, 2014-2018 data was unavailable and consequently is not included in this report.

³ Of the 33 hospitals licensed in 2018, only 26 hospitals performed procedures in which to monitor SSI.

Influenza Vaccination Coverage in Hospital Healthcare Personnel

All 33 acute care, psychiatric, and rehabilitation hospitals reported healthcare personnel (HCP) influenza vaccination percentages. Vaccination coverage by hospital ranged from 65.9% to 99.6%, and the hospital State percentage was 94.0%. Ten hospitals had vaccination percentages similar to the overall State vaccination percentage, thirteen hospitals reported vaccination percentages that were significantly higher than the overall State vaccination percentage, and ten hospitals reported vaccination percentages that were significantly lower than the overall State vaccination percentage. The statewide hospital HCP vaccination percentage increased from the 2017-18 influenza season (93.3%) to the 2018-19 influenza season (94.0%); this was statistically significant. Specifically, five hospitals increased HCP influenza vaccination coverage in 2018-19 compared to the 2017-18 influenza season, 24 hospitals had similar vaccination coverage, and four hospitals decreased vaccination coverage.

Conclusion

This tenth report of hospital HAI data displays continuous progress toward the goal of eliminating HAI in NH. This report provides a picture of selected HAI data, which can be used by healthcare facilities in the state to identify areas for improvement and prevention as well as healthcare consumers to make informed healthcare decisions.

I. INTRODUCTION

A. Purpose

This report represents the tenth summary of healthcare-associated infection (HAI)-related data reported by hospitals in New Hampshire (NH) during calendar year 2018. This report can be used by healthcare facilities in the state to identify areas for improvement as well by healthcare consumers to make informed healthcare decisions.

B. Audience

The intended audience may include, but is not limited to: healthcare personnel (HCP), infection control and prevention staff, facility leadership and management, clinicians, and healthcare consumers.

C. How to Use this Document

This document includes aggregate data reported by all 33 acute care, critical access, and specialty hospitals in NH. This report also includes individual hospital reports on page 81. The document consists of six sections:

- I) Introduction
- II) Surveillance methods
- III) Statewide data
 - a. Overall NH data
 - b. Central line-associated bloodstream infections (CLABSI)
 - c. Central line insertion practices (CLIP)
 - d. Catheter-associated urinary tract infections (CAUTI)
 - e. Surgical site infections (SSI) following coronary artery bypass graft (CABG), colon (COLO), abdominal hysterectomy (HYST), and knee arthroplasty (KPRO) procedures
 - f. Post-discharge surveillance methods
 - g. Surgical antimicrobial prophylaxis administration
 - h. Percentage of HCP receiving influenza vaccination
- IV) Conclusions
- V) Individual hospital reports
- VI) Appendices
 - a. Technical notes
 - b. Influenza vaccination survey questions, 2017-18 season
 - c. Understanding the relationship between HAI rates and standardized infection ratio (SIR) comparison metrics
 - d. Preventing HAI
 - e. Map of NH hospitals

Please contact the NH Department of Health and Human Services (DHHS) Healthcare-Associated Infections Program (603-271-4496) with any questions about the content or how to use this document.

D. Background on Healthcare-Associated Infections

An HAI is an infection that a patient acquires during the course of receiving treatment for another condition within a healthcare setting. An estimated 722,000 HAI and 75,000 associated deaths occurred in United States (U.S.) acute care hospitals in 2011.ⁱⁱ This may represent a decreasing trend because previous studies depict higher numbers of HAI; 1.7 million infections and 99,000 deaths each year.ⁱⁱⁱ By these estimates, HAI are among the top 10 leading causes of death in the U.S., and 5–10% of all hospital admissions are complicated by HAI.^{iv} The economic burden of HAI is substantial and increasing. The total cost of HAI has been estimated at \$33 billion per year in U.S. hospitals.^v The most common HAI are pneumonia, gastrointestinal illness, primary bloodstream infections (BSI), and SSI.ⁱⁱ

E. New Hampshire Healthcare-Associated Infections Program

The NH DHHS has been developing and improving a HAI surveillance program since 2007. During the 2006 legislative season, the NH Legislature passed a bill creating NH Revised Statutes Annotated (RSA) 151:32-35, which requires hospitals to identify, track, and report HAI to DHHS. RSA 151:33 specifically requires reporting of CLABSI, SSI, ventilator-associated pneumonia (VAP), CLIP, surgical antimicrobial prophylaxis, and influenza vaccination coverage. The intent of the bill is to provide HAI data by hospital in a publicly accessible forum. Because the bill did not include funding to carry out these activities, mandatory reporting was not fully implemented until January 2009.

DHHS, with consideration of the law, required that eligible hospitals initially report the following measures:

- CLABSI in adult intensive care units (ICU) (via NHSN). Only those hospitals with ICU enroll and report data to NHSN.
- CLIP in all ICU (via NHSN). Only those hospitals with ICU enroll and report data to NHSN.
- SSI following CABG, colon, and knee arthroplasty procedures (via NHSN). Only those hospitals that perform the selected procedures enroll and report data to NHSN.
- Surgical antimicrobial prophylaxis (via Centers for Medicare and Medicaid Services [CMS]). Only those hospitals that administer antimicrobial prophylaxis report these data.
- Influenza vaccination in patients and HCP (via DHHS web-based survey). All hospitals (including rehabilitation and psychiatric) report influenza vaccination in HCP.

All 26 acute care hospitals successfully enrolled in NHSN and began reporting the required data in January 2009.

During the 2010 legislative season, the NH Legislature passed House Bill 1548 (2010) amending RSA 151:32-35 to require all licensed ambulatory surgery centers (ASCs) to report HAI to DHHS.

HAI data reported by ASC is published in a separate report and posted to the HAI Program publications website: <http://www.dhhs.nh.gov/dphs/cdcs/hai/publications.htm>.

The administrative rules related to HAI reporting were revised in 2011 to include additional reporting measures for eligible hospitals. Starting January 2012, hospitals were also required to report:

- CLABSI in all ICU (via NHSN)
- CLIP in all ICU (via NHSN)
- Catheter-associated urinary tract infections (CAUTI) in all pediatric and adult ICU (via NHSN)
- SSI following abdominal hysterectomy (HYST) procedures (via NHSN)

F. State of New Hampshire Healthcare-Associated Infections Plan

In response to increasing concerns about the public health impact of HAI, the U.S. Department of Health and Human Services (HHS) developed its “Action Plan to Prevent Healthcare-Associated Infections” (HHS Action Plan) in 2009. The HHS Action Plan includes recommendations for surveillance, research, communication, and metrics for measuring progress toward national goals. In a concurrent development, the 2009 Omnibus Appropriations Act required states receiving Preventive Health and Health Services Block Grant funds to certify that they would submit a plan to reduce HAI to the Secretary of HHS not later than January 1, 2010. In order to assist states in responding within the short timeline required by that language and to facilitate coordination with national HAI prevention efforts, the Centers for Disease Control and Prevention (CDC) provided a template to assist state planning efforts in the prevention of HAI. The template targeted four areas: 1) Development or Enhancement of HAI Program Infrastructure; 2) Surveillance, Detection, Reporting, and Response; 3) Prevention; and 4) Evaluation, Oversight, and Communication. In 2009, DHHS drafted its State HAI Plan and submitted it to HHS. Updates to the plan is posted to the HAI Program website: <http://www.dhhs.nh.gov/dphs/cdcs/hai/index.htm>.

G. Overview of Healthcare-Associated Infections Prevention Efforts

DHHS participates in statewide prevention activities through the NH Health Care Quality Assurance Commission (NHHQCAC), on which the Division of Public Health Services Director serves. DHHS is active in various projects coordinated by the NHHQCAC and the CMS Quality Innovation Network-Quality Improvement Organization (QIN-QIO). Major statewide initiatives through these organizations have included hand hygiene campaigns, patient safety checklists, and programs to prevent BSI, antimicrobial resistance, and *Clostridium difficile*. Additionally, the Foundation for Healthy Communities received a large grant through the Partnership for Patients program to conduct additional large, statewide prevention initiatives. For additional information on these various efforts, the following websites may be helpful:

New Hampshire Health Care Quality Assurance Commission:

<http://www.healthynh.com/fhc-initiatives/nh-health-care-quality-assurance-commission.html>

CMS QIN-QIO for Connecticut, Maine, Massachusetts, NH, Rhode Island, and Vermont:

www.HealthCareForNewEngland.org

Foundation for Healthy Communities Partnership for Patients:

<http://www.healthynh.com/partnership-for-patients.html>

In addition to supporting and engaging in prevention activities with patient safety groups, the HAI Program provides educational opportunities to healthcare facilities across the state in order to share best practices for infection prevention and ultimately reduce HAI.

H. Healthcare-Associated Infections Technical Advisory Workgroup

In the spring of 2009, DHHS formed an HAI Technical Advisory Workgroup (TAW). The purpose of the TAW is to provide scientific and infection prevention expertise to the HAI Program. The TAW meets quarterly, and as a forum for stakeholder participation in decision-making around the HAI Program. The TAW is currently a 25-member group that includes representation from stakeholders across NH and includes representatives from various sizes and types of hospitals and ASC, infection control associations, a consumer advocate, the NH Hospital Association, the New Hampshire Healthcare Quality Assurance Commission, the New Hampshire Ambulatory Surgery Association, and the Northeast Health Care Quality Foundation (see page 16 for a list of TAW members during the 2018 reporting year).


HEALTHCARE-ASSOCIATED INFECTIONS TECHNICAL ADVISORY WORKGROUP

Members	
Benjamin Chan, MD, MPH	Gloria Thorington
Beth Daly, MPH	Anne Diefendorf, MS, RD, LD
Katrina Hansen, MPH	Terri E. Kangas-Feller, BS, RN-BC, CIC
Yvette Perron, MPH	Tanya Lord MPH, PhD
Trevor Bauer, MPH	Elissa Malcolm, MS
Carly Zimmermann, MPH, MLS(ASCP)cm	Claudia Alvarado, RN
Elizabeth Talbot, MD	Donna Quinn, RN, BSN, MBA
Michael Fleming	Cathy Foti, RN
Darlene Burrows, RN, BSN, CIC	Dr. Michael Calderwood
Lynda Caine, RN, MPH, CIC	Dr. Antonia Altomare
Ashley Conley, MS, CPH	Nancy Gooch, RN

*Served on TAW for part of 2018

DHHS: New Hampshire Department of Health and Human Services

Figure 1. Types of healthcare-associated infections reported to NH Department of Health and Human Services



II. SURVEILLANCE METHODS

A. 2018 Healthcare-Associated Infections Reporting Requirements for New Hampshire Hospitals

Reporting requirements are governed by RSA 151:33 with authority given to DHHS to develop administrative rules to provide specific reporting instructions and methodology. Administrative rules, “He-P 309 Healthcare Associated Infections,” were drafted in 2010 with stakeholder input and approved January 14, 2011 by the Joint Legislative Committee on Administrative Rules. Reporting requirements for 2009-2018 included the following required measures for hospitals:

- CLABSI in adult ICU
- CLIP in adult ICU
- SSI following CABG, colon, and knee arthroplasty procedures
- Surgical antimicrobial prophylaxis
- Influenza vaccination in patients and HCP

The rules were updated in 2012 to include the following required measures for hospitals:

- CLABSI in all ICU
- CLIP in all ICU
- CAUTI in all adult and ICU
- SSI following CABG, colon, abdominal hysterectomy, and knee arthroplasty procedures
- Surgical antimicrobial prophylaxis
- Influenza vaccination in patients and HCP

While all licensed hospitals including acute care and specialty hospitals are required to report the selected measures under RSA 151:33, specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CAUTI, CLABSI and CLIP, because they do not have ICU, nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries. The five rehabilitation and two psychiatric hospitals in NH are only required to report influenza vaccination coverage for patients and HCP.

B. Selection of Reporting Requirements

RSA 151:33 broadly requires reporting of all SSI and CLABSI; however, it is not feasible to perform surveillance for all of these infections using NHSN. In order to generate infection measures for hospitals and compare them with national data, infection reporting was limited to the capabilities of NHSN and measures were selected in accordance with national recommendations for HAI surveillance in the context of public reporting.

In 2005, the CDC released a report titled “Guidance on Public Reporting of Healthcare-Associated Infections: Recommendations of the Healthcare Infection Control Practices Advisory Committee (HICPAC).”^{vi} The group recommended selecting outcome measures for reporting based on the

frequency, severity, and preventability of the outcomes and the likelihood that they can be detected and reported accurately. Specifically, the group recommended monitoring the following outcome measures:

- CLABSI in ICU
- SSI following selected operations
- CAUTI and VAP were not recommended because of lower morbidity and mortality resulting in less prevention effectiveness relative to the burden of data collection and reporting (in the case of CAUTI), and difficulty in detecting infections accurately resulting in invalid and misleading comparisons of infection rates for consumers (in the case of VAP)

Additionally, the group recommended monitoring the following process measures:

- CLIP
- Surgical antimicrobial prophylaxis
- Influenza vaccination of patients and HCP

In 2008, the Healthcare-Associated Infections Working Group⁴ of the Joint Public Policy Committee released “Essentials of Public Reporting of Healthcare-Associated Infections: A Tool Kit.”^{vii} The working group agreed with the CDC/HICPAC document, “Guidance on Public Reporting of Healthcare-Associated Infections” (referenced above) and recommended exclusion of outcome measures related to VAP and CAUTI because the existing surveillance criteria are difficult to apply consistently, making case counts unreliable. The toolkit recommends monitoring the following outcome measures:

- CLABSI in ICU
- Surgical procedures that are performed with adequate frequency to permit meaningful comparisons among institutions. Specific reasonable options listed were: 1) CABG; 2) colon resection; 3) total hip arthroplasty; 4) total knee arthroplasty; 5) laminectomy; and 6) total abdominal hysterectomy

The only process measure the group recommended monitoring was HCP influenza vaccination coverage.

Within the context of RSA 151:33, DHHS reviewed the national guidelines and capabilities of NHSN in selecting infection and process measures. It is expected that these reporting requirements may change in the future as we learn from public reporting, as HAI epidemiology evolves, and as new surveillance methods and reporting technologies become available.

⁴ The Healthcare-Associated Infection Working Group of the Joint Public Policy Committee is a multi-organizational group represented by the Association for Professionals in Infection Control and Epidemiology, CDC, Council of State and Territorial Epidemiologists, and Society for Healthcare Epidemiology of America.

C. Accuracy of Reported Healthcare-Associated Infections Surveillance Data

DHHS conducted a validation study of 2014-2015 data to assess the degree of under- and over-reporting and to provide additional training to address any common or systematic errors in reporting processes. DHHS contracted with an independent, external agency to perform the validation study and HAI Program staff participated in activities including NHSN data review, medical record review, data analysis, corrections, and follow-up for deficiencies. Overall, validation of 2014-2015 data showed that there was approximately 15% under-reporting of CLABSI, CAUTI and SSI combined across all NH hospitals. This under-reporting was mostly due to misunderstandings of the NHSN definitions for HAI. In addition to under-reporting, the validation studies also found 10% of CLABSI, CAUTI and SSI were over reported or not classified accurately (i.e., reporting an infection that was not truly a CLABSI, CAUTI or SSI). The 2014-2015 data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over reporting of HAI. However, the HAI Program is currently in the process of validating data on a rolling basis.

Several processes are used to ensure that these 2018 data are as accurate as possible. First, DHHS selected NHSN for mandatory reporting, which requires the use of standardized infection definitions and reporting methods. Secondly, DHHS provided preliminary data reports to each hospital with the request to confirm accuracy. This reconciliation process was iterative until all hospitals made corrections and agreed to the reported data. Lastly, 2009-2010 and 2014-2015 data validation was performed, reducing systematic errors that may have occurred during the reporting process; this has likely resulted in a lasting improvement to data quality, even in years when validation does not take place.

Despite the above measures, there are several limitations to the reporting methods that may limit comparison of data across hospitals. Definitions for classifying an infection as healthcare-associated are standardized through the use of NHSN; however, methods to identify the infection in each hospital are not. For example, hospitals may use different methods to identify CLABSI (e.g., reviewing laboratory records, reviewing ICU records) or may have different approaches to diagnosing and managing suspect CLABSI in the ICU. For SSI, identifying patients who develop infections after discharge from the hospital can be difficult, and each hospital may use a different method of post-discharge surveillance (e.g., letters to surgeons, conducting chart reviews for surgical patients, calling surgeon offices). These different approaches vary in sensitivity. See page 24 for more details about how hospitals identify SSI.

D. National Healthcare Safety Network

NHSN is a voluntary, secure, internet-based surveillance system for healthcare facilities to monitor patient safety and infection prevention measures. Enrollment is open to all types of healthcare facilities in the U.S. DHHS selected NHSN because it is widely used across the entire U.S., it offers already developed and accepted surveillance definitions and methods, it provides national comparison data, and there is no cost to use or join the system.

NHSN collects and analyzes healthcare-associated infections (HAI) data reported from healthcare-facilities across the U.S. to track HAI incidence, identify opportunities to eliminate HAIs, and measure the progress of HAI prevention efforts. Progress is measured using a summary statistic called the standardized infection ratio (SIR). This comparative metric is calculated after risk adjusting the data reported into NHSN. Risk adjusting is a statistical process used to account for differences in patient characteristics that may influence health care outcomes. The SIR compares the number of infections in a facility or state to the number of infections that were “predicted”, or would be expected, to have occurred based on previous years of reported data (i.e., baseline data). The number of predicted infections is an estimate based on aggregated data reported to CDC’s NHSN during a specific baseline period. The current risk adjustment methods and baseline periods vary by HAI type and/or healthcare facility type.

January 2017, NHSN updated both the source of aggregate data and the risk adjustment methodology used to create the original baseline. This update is known as “2015 rebaseline”. HAI prevention progress will be measured in comparison to infection data reported to NHSN using updated risk-adjustment models.

Previously calculated SIRs have different baseline years for each infection type and facility type. Starting with 2017 data, SIRs will only be calculated under the new risk models. The 2016 Annual HAI report was the last year data was analyzed using the original baseline in NH. The 2017 Annual HAI report includes 2016 and 2017 data analyzed using the 2015 baseline.

Starting 2018, NH DHHS annual HAI reports will be measured using the new 2015 baseline from 2015. The data analyzed in this report utilized the new national baseline based on 2015 data, which includes new risk adjustments methodology. Therefore, the data in this report cannot be compared to previous New Hampshire State HAI reports. Of note, facility SIRs have increased and shifted closer to 1 with the new baseline (click here for more information about the updated NHSN baseline: <https://www.cdc.gov/nhsn/2015rebaseline/index.html>). The higher SIRs observed in the report may partially be a reflection of using the new 2015 baseline.



Additional NH HAI Reports are available at:
<https://www.dhhs.nh.gov/dphs/cdcs/hai/publications.htm>

More information about NHSN is available at: <http://www.cdc.gov/nhsn/index.html>.

E. Comparisons with National Data

All SSI comparisons with national data use 2013 NHSN data published in the “National Healthcare Safety Network (NHSN) report: Data summary for 2013, Device Associated Module, issued December 2015.”^{viii} All device-associated infection (CLABSI and CAUTI) comparisons with national data use 2013 data.

These reports are available at: <https://www.cdc.gov/nhsn/datastat/index.html>.

F. Central Line-Associated Bloodstream Infections Surveillance

A CLABSI is a laboratory-confirmed BSI that develops after insertion of a central line and is not secondary to an infection at another body site. A central line is an intravascular catheter that terminates at or close to the heart or in one of the great vessels and is used for infusion, withdrawal of blood, or hemodynamic monitoring. Hospitals are required to monitor and report CLABSI in adult ICU. This monitoring includes reporting the number of infections identified as well as the total number of central line days in the unit. These metrics are monitored following NHSN protocols and definitions, and reported in NHSN.

Central line days are the number of patients with one or more central lines of any type, which are counted at the same time each day and aggregated over the reporting period. For example, a patient with a central line in place for five days would be counted as five central line days.

Detailed descriptions of the NHSN CLABSI surveillance protocols are available at: http://www.cdc.gov/nhsn/PDFs/pscManual/4PSC_CLABScurrent.pdf.

Limitations for CLABSI surveillance:

- NHSN only allows for monitoring CLABSI in inpatient units. In NH in 2017, CLABSI were monitored in all ICU (including pediatric and neonatal ICU) and not in other inpatient locations.
- Validation of 2014-2015 data showed that there was approximately 19% under-reporting of CLABSI across all NH hospitals. This under-reporting was mostly due to misunderstandings about the NHSN definition for CLABSI. In addition to under-reporting, the validation studies also found 3% of over-reporting (i.e., reporting an infection that was not truly a CLABSI). The 2018 CLABSI data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under-and-overreporting of infections.

G. Central Line Insertion Practices Monitoring

CLIP monitoring assesses key infection prevention practices that occur during the insertion of a central line. A central line is any intravascular catheter used for infusion, blood withdrawal, or hemodynamic monitoring that terminates at or close to the heart or in one of the great vessels. In order to comply with all infection prevention practices during the insertion, the inserter must 1) perform hand hygiene prior to insertion; 2) use all five barriers (gloves, gown, cap, mask, and drape); 3) use an appropriate skin preparation agent; and 4) ensure skin is dry prior to insertion.

Hospitals monitor and report CLIP data through NHSN using all NHSN protocols and definitions. In 2018, hospitals were required to monitor all central line insertions that were placed in ICU (which includes pediatric, neonatal, and step down units). The NHSN CLIP protocols are available at: http://www.cdc.gov/nhsn/PDFs/pscManual/5psc_CLIPcurrent.pdf.

Occupational groups are compared with the overall State compliance percentage since there are no national data for comparison. Groups with a confidence interval that overlaps the State's confidence interval are considered to be similar to the State adherence percentage. Any occupation or hospital with a confidence interval that is higher than the State's confidence interval is considered to have a significantly higher percentage than the State adherence percentage. Groups with a confidence interval that is lower than the State's confidence interval are considered to have a significantly lower percentage than the State adherence percentage.

Limitations for central line insertion practices monitoring:

- In NH, CLIP was monitored in all ICU (including pediatric and neonatal ICU) and not in other settings where central lines may be inserted (e.g., operating room, procedure rooms, emergency room, dialysis centers).
- The person recording the insertion practices may differ in each hospital. This person may be an observer or the person doing the insertion, which may impact quality of data on adherence reported.

H. Catheter-Associated Urinary Tract Infections Surveillance

A CAUTI is a urinary tract infection that develops after insertion of an indwelling urinary catheter and is not secondary to an infection at another body site. An indwelling urinary catheter is a drainage tube that is inserted into the urinary bladder through the urethra and left in place, and is connected to a drainage bag. They are sometimes called Foley catheters and are used for intermittent or continuous irrigation or urine drainage. Hospitals are required to monitor and report CAUTI in all ICU (excluding neonatal ICU and step down units). This monitoring includes reporting the number of infections identified as well as the total number of catheter days in the unit. These metrics are monitored following NHSN protocols and definitions and reported in NHSN.

Catheter days represent the number of patients with one or more indwelling urinary catheters of any type, counted at the same time each day and aggregated over the reporting period. For example, a patient with a catheter in place for five days would be counted as five catheter days;

one patient with a catheter for one day and another with a catheter for four days are also counted as five catheter days.

Detailed descriptions of the NHSN CAUTI surveillance protocols are available at:

<http://www.cdc.gov/nhsn/PDFs/pscManual/7pscCAUTICurrent.pdf>.

Limitations for CAUTI surveillance:

- NHSN only allows for monitoring CAUTI in inpatient units. In NH in 2018, CAUTI were monitored in all ICU (excluding neonatal ICU) and not in other inpatient locations.
- The 2018 CAUTI data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.

I. Surgical Site Infections Surveillance

An SSI is an infection that develops at the site of a surgical procedure. There are different ways to classify an SSI, such as whether it is superficial, in deep tissue, or in the organ/space. Monitoring for an SSI may continue for as little as 30 days or as long as 90 days based on depth and procedure type (e.g., knee arthroplasty, CABG). In 2018, hospitals were required to monitor and report SSI for four procedures:

- Coronary Artery Bypass Graft (chest incision and donor site)
 - NHSN Operative Procedures CBGC (coronary artery bypass graft procedures with chest incision site only) and CBGB (coronary artery bypass graft procedures with both a chest and donor site incision)
- Colon Surgery (incision, resection, or anastomosis of the large intestine; includes large-to-small and small-to-large bowel anastomosis; does not include rectal operations)
 - NHSN Operative Procedure COLO
- Abdominal Hysterectomy (includes that by laparoscope)
 - NHSN Operative Procedure HYST
- Knee Arthroplasty
 - NHSN Operative Procedure KPRO

Specific ICD-10 codes can be found at:

<https://www.cdc.gov/nhsn/xls/2017-icd-10-pcs-code-mapping-opc.xlsx>

SSI monitoring includes total counts as well as patient-level information for all patients undergoing the same procedure. This allows for appropriate risk adjustment, because risk for development of an SSI can be influenced by patient- and procedure-specific factors. Patient and procedure risk factors that are considered when assessing SSI SIR by hospital vary by type of procedure but include factors such as:

- Operation lasting more than the duration of cut point hours⁵
- Contaminated (Class III) or Dirty/Infected (Class IV) surgical wound class
- American Society of Anesthesiologists (ASA) Classification of Physical Status score of 3, 4, or 5 (see below)
- Age of the patient
- Gender of the patient
- Hospital bed size
- Hospital's medical school affiliation
- Whether the surgery was the result of trauma

The wound class is a way of determining how clean or dirty the operative body site was at the time of the operation. Operation body sites are divided into four classes:

Class I/Clean: An uninfected operation body site is encountered and the respiratory, digestive, genital, or uninfected urinary tracts are not entered.

Class II/Clean-Contaminated: Operation body sites in which the respiratory, digestive, genital, or urinary tracts are entered under controlled conditions and without unusual contamination.

Class III/Contaminated: Operation body sites that have recently undergone trauma, operations with major breaks in sterile technique (e.g., open cardiac massage), or gross spillage from the gastrointestinal tract.

Class IV/Dirty or Infected: Includes old traumatic wounds with retained dead tissue and those that involve existing infection or perforated intestines.

The ASA score is a scale used by the anesthesiologist to classify the patient's physical condition prior to surgery. It is one of the factors that help determine a patient's risk of possibly developing SSI.

The ASA scale is:

1. Normally healthy patient
2. Patient with mild systemic disease
3. Patient with severe systemic disease
4. Patient with an incapacitating systemic disease that is a constant threat to life
5. A patient who is not predicted to survive with or without the operation

All SSI metrics are monitored following NHSN protocols and definitions and reported in NHSN.

The NHSN SSI protocols are available at:

<http://www.cdc.gov/nhsn/PDFs/pscManual/9pscSSIcurrent.pdf>.

⁵ Cut points are assigned based upon the time that the majority (75%) of a specific procedure takes to perform. The duration cut point is measured in minutes and is the time between the skin incision and skin closure.

In general, most SSI identified during the initial hospital encounter or those that require readmission are thought to be well-represented in HAI surveillance data. However, the infections that develop after the patient is discharged home that do not require readmission are thought to be less well-represented, as inclusion in surveillance requires the healthcare facility to proactively seek out these infections, a process known as post-discharge surveillance.

The proportion of infections detected through post-discharge surveillance in comparison to the state average may provide an indicator of how well the facility is able to identify these infections, which ultimately can impact the facility's SSI SIR (better surveillance may result in a higher SIR). SSI data detected through post-discharge surveillance were analyzed for 2013-2014 and infection control staff were interviewed regarding methods of SSI surveillance in 2011. The percent of SSI detected post-discharge was calculated for each hospital and compared to a moving state average (hospital vs. all other hospitals). Statistical significance was calculated using the NHSN Statistics Calculator.

Limitations for SSI surveillance:

- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections. Post-discharge surveillance methods were analyzed to better understand these differences between facilities and are presented in this report on page 69.
- SSI reporting in NHSN requires not only reporting of infections but also detailed information on each patient undergoing the procedure being monitored. This allows for risk adjustment. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- Some procedures require monitoring for SSI for up to 90 days after the procedure depending on the depth of infection (in NH, this includes CABG and knee arthroplasty). Due to the reporting deadlines required for producing a data report such as this, it is possible that deep or organ/space SSI associated with surgeries performed at the end of 2018 may not be included in this report. As such, this report may not account for all SSI that developed as a result of procedures performed in 2018.
- The SSI data presented in this report includes all types of infections, including superficial SSI, which can occur as a result of care in the hospital but also as a result of the patient's care of the wound site once discharged.
- Validation of 2014-2015 data showed that there was approximately 15% under-reporting of SSI across all NH hospitals. This under-reporting was mostly due to misunderstandings about the NHSN definition for SSI. In addition to under-reporting, the validation studies also found 3% over-reporting (i.e., reporting an infection that was not truly a SSI). The 2018 SSI data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.

J. Surgical Antimicrobial Prophylaxis Administration Monitoring

All NH hospitals report surgical antimicrobial prophylaxis data and other measures to CMS through the Surgical Care Improvement Project (SCIP). For this reason, DHHS does not collect surgical antimicrobial prophylaxis data directly from hospitals. In addition to other measures required by CMS, measures relative to NH RSA 151:33 include the following:

- SCIP 1: Number and percentage of patients who received prophylactic antibiotic within one hour prior to surgery
- SCIP 2: Number and percentage of patients who received the appropriate prophylactic antibiotic
- SCIP 3: Number and percentage of patients whose prophylactic antibiotic was discontinued within 24 hours after surgery

These process measures show a hospital's adherence rate to best practices designed to reduce surgical complications. Hospitals follow the CMS specification manual appropriate to the date of discharge found at:

<http://qualitynet.org/dcs/ContentServer?cid=1141662756099&pagename=QnetPublic%2FPage%2FQnetTier2&c=Page>.

In previous years, DHHS accessed hospital data on surgical antimicrobial prophylaxis administration from the New Hampshire Quality Care website at:

<http://www.healthynh.com/fhc-initiatives/nh-health-care-quality-assurance-commission.html>

As of July 1, 2016, SCIP data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.

K. Influenza Vaccination Percentage Monitoring

HCP can become infected with the influenza virus through contact with infected patients and can transmit influenza to patients and other HCP. Despite documented benefits of HCP influenza vaccination on patient outcomes and HCP absenteeism nationally, vaccination coverage among HCP remains low. In a CDC survey, influenza vaccination coverage in HCP nationally was 88.7% during the 2017-18 influenza season.^{ix} Because HCP provide care to patients at high risk for complications of influenza, they should be offered influenza vaccine each year. Currently there are no regulations requiring vaccination in NH, and HCP are free to decline vaccination for any reason. However, some hospitals do have policies requiring mandatory HCP vaccination. Vaccination coverage in hospital HCP have been monitored in NH for several years.

All hospitals are required to report HCP and patient vaccination data directly to DHHS. This reporting occurs either solely via a web-based survey provided to facilities, or via NHSN in combination with an abbreviated web-based survey, newly for the 2018-19 influenza season and according to facility discretion. See Appendix 2 for the 2018-19 survey questions regarding influenza vaccination. Data for the 2018-19 influenza season were reported by all hospitals.

Submission of these data meets the requirements of both the HAI law (RSA 151:32-35) and the healthcare immunization law (RSA 151:9-b).

HCP influenza vaccination percentages were calculated by dividing the total number of HCP that worked or volunteered in each facility for at least one working day between October 1, 2018 and March 31, 2019 by the total number of HCP immunized against influenza for the 2018-19 influenza season.

Limitations for influenza vaccination monitoring:

- The data collection tools ask for the total number of HCP vaccinated. This may not reflect the number of HCP to whom the vaccine was offered. Hospitals may vary in the refusal percentage for vaccination among HCP and the reasons for such refusal. Additionally, some HCP may not be eligible to receive the vaccine. DHHS attempted to assess why unvaccinated HCP did not receive the vaccine; however, not all hospitals were able to report this information.
- Because the web-based survey did not include options for facilities to report unknown vaccination status, patients and HCP with unknown vaccination status were analyzed as though they were not vaccinated. This results in a conservative estimate of vaccination status (e.g., lower than in reality).
- Vaccination status is not uniformly available by location where the vaccination was received (e.g., at the reporting facility or elsewhere).
- Data collection techniques at hospitals may vary from season to season, potentially affecting comparison of data. DHHS continues to work towards improving the validity and utility of this measure in order to eliminate issues that pose problems for such comparison.
- Reporting patient vaccination percentages is limited by availability of vaccine and by hospitals' ability to track why patients did not receive the vaccine. For example, the survey asks for admissions through March 31, 2019, by which time some hospitals may have used their vaccine supply and are unable to order more. This scenario would result in a lower vaccination percentage because the survey counts all admissions through March, even though there was no opportunity to vaccinate these patients due to supply. DHHS has elected not to report patient vaccination percentages until a better way to collect the information is identified so that results are reliable, accurate, and informative.
- For the 2014-15 influenza season, CMS began requiring all facilities sharing the same CMS Certification Number (CCN) to report this measure in aggregate via NHSN. Because some ASC and hospitals may share the same CCN, it is possible that HCP influenza vaccination data contains more duplicate data than in prior influenza seasons.

III. STATEWIDE DATA

HAI data are presented throughout this report as both SIR and rates as appropriate. Presenting data as an SIR allows for aggregating data across risk groups, procedures, and hospitals to gain a better understanding of the incidence of HAI while still adjusting for underlying patient or hospital factors that may affect the occurrence of infections. The SIR allows comparison between how many infections actually occurred and how many were predicted to occur based on national data. Specific annual rate information is also provided where possible, which represents the number of infections that occurred. Rate data are limited in that they must be stratified by certain factors, such as hospital and type of ICU; they cannot be aggregated over these categories for the purpose of analysis. See technical notes for additional information on rates and the SIR.

Because an SIR is a comparison of the number of actual observed infections to the number predicted based on national data, an SIR of 1.0 means that exactly the same number of infections was observed as was predicted. An SIR of less than one means that fewer infections were observed than were predicted (for example, SIR = 0.70 would be interpreted as 30% fewer infections observed than predicted). An SIR of more than one means that more infections were observed than were predicted (for example, SIR = 1.30 would be interpreted as 30% more infections observed than predicted). A confidence interval is calculated to determine whether the difference between observed and predicted infections is statistically significant. If the difference is not statistically significant, the observed and predicted numbers of infections are considered similar. See technical notes for additional information on confidence intervals.

This report provides comparisons with national and State data where appropriate. Comparisons are color coded consistently throughout. For infections, yellow represents infection rates or SIR that are similar to national data, red represents infection rates or SIR that are significantly higher than national data, and green represents infection rates or SIR that are significantly lower than national data.

 SIR: fewer than predicted  SIR: similar to predicted  SIR: more than predicted

For process measures, yellow represents percentages that are similar to the State percentage, red represents percentages that are significantly lower than the State percentage, and green represents percentages that are significantly higher than the State percentage.

 higher than State  similar to State  lower than State

Statistical significance is affected by sample size. If a value is almost or just barely significant, just a few additional observations can push significance one way or the other (i.e., not significant or significant).

A. Statewide Standardized Infection Ratios

In 2018, 196 HAI were reported by all 26 acute care hospitals in NH. These infections represent CLABSI and CAUTI in ICU and SSI following colon, knee arthroplasty, abdominal hysterectomy, and CABG procedures. A total of 188.89 infections were predicted based on national data; the overall observed number of HAI was 4% higher than predicted. More specifically, there were 27%


fewer CLABSI, 16% more CAUTI and 7% more SSI. Looking individually at the specific procedures tracked for SSI by NH hospitals, there were 9% fewer infections following CABG procedures, 1% more infections following colon procedures, 6% more infections following abdominal hysterectomy procedures, and 36% more infections following knee arthroplasty procedures. However, the differences for colon, and abdominal hysterectomy procedures are not statistically significant, and the number of infections observed are considered similar to national data. The difference for coronary artery bypass was statistically significant when compared to national data, and the number of infections observed are considered lower than national data. These data are shown in Table 1 and Figure 2.

Table 1. Statewide standardized infection ratios, Jan 1–Dec 31, 2018

	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted Number of Infections
Overall HAI SIR	196	188.89	1.04	0.90 , 1.19	Similar
	The overall observed number of HAI in New Hampshire hospitals was 4% more than predicted based on national data. This difference is not statistically significant, which means the overall number of HAI in the state is SIMILAR than the number seen nationally.				
CLABSI SIR	20	27.38	0.73	0.46 , 1.11	Similar
	The overall observed number of CLABSI in New Hampshire hospitals was 27% fewer than predicted based on national data. This difference is not statistically significant, which means the overall number of CLABSI in the state is SIMILAR than the number seen nationally.				
CAUTI SIR	43	36.98	1.16	0.85 , 1.55	Similar
	The overall observed number of CAUTI in New Hampshire hospitals was 16% higher than predicted based on national data. This difference is not statistically significant, which means the overall number of CAUTI in the state is SIMILAR to the number seen nationally.				
Overall SSI SIR	133	124.53	1.07	0.90 , 1.26	Similar
	The overall observed number of SSI in New Hampshire hospitals was 7% more than predicted based on national data. This difference is not statistically significant, which means the overall number of SSI in the state is SIMILAR than the number seen nationally.				
CABG SIR	11	12.12	0.91	0.48 , 1.58	Similar
	The overall observed number of CABG infections in New Hampshire hospitals was 9% fewer than predicted based on national data. This difference is not statistically significant, which means the overall number of CABG infections in the state is SIMAR than the number seen nationally.				
COLO SIR	74	73.01	1.01	0.80 , 1.27	Similar
	The overall observed number of COLO infections in New Hampshire hospitals was 1% higher than predicted based on national data. This difference is not statistically significant, which means the overall number of COLO infections in the state is SIMILAR to the number seen nationally.				
HYST SIR	15	14.17	1.06	0.62 , 1.71	Similar
	The overall observed number of HYST infections in New Hampshire hospitals was 6% more than predicted based on national data. This difference is not statistically significant, which means the overall number of HYST infections in the state is SIMILAR to the number seen nationally.				
KPRO SIR	33	24.32	1.36	0.95 , 1.88	Similar
	The overall observed number of KPRO infections in New Hampshire hospitals was 36% more than predicted based on national data. This difference is not statistically significant, which means the overall number of KPRO infections in the state is SIMILAR to the number seen nationally.				

HAI: Healthcare-associated infection, CLABSI: Central line-associated blood stream infections, CAUTI: Catheter-associated urinary tract infections, SSI: Surgical site infections, CABG: Surgical site infections associated with coronary artery bypass graft procedures, COLO: Surgical site infections associated with colon procedures, HYST: Surgical site infections associated with abdominal hysterectomy procedures, KPRO: Surgical site infections associated with knee arthroplasty procedures

Figure 2. Overall statewide standardized infection ratios, Jan 1–Dec 31, 2018



HAI: Healthcare-associated infection
 CLABSI: Central line-associated blood stream infections
 CAUTI: Catheter-associated urinary tract infections
 SSI: Surgical site infections
 CABG: Surgical site infections associated with coronary artery bypass graft procedures
 COLO: Surgical site infections associated with colon procedures
 HYST: Surgical site infections associated with abdominal hysterectomy procedures
 KPRO: Surgical site infections associated with knee arthroplasty procedures

B. Overall Standardized Infection Ratios by Hospital

Table 2 and Figure 3 below show the total number of HAI reported by each hospital. These infections represent CLABSI and CAUTI in ICU and SSI following colon, abdominal hysterectomy, knee arthroplasty, and coronary artery bypass graft procedures. Seventeen hospitals had sufficiently robust data to present. Fourteen hospitals observed a similar number of infections as were predicted based on national data, two observed a lower number of infections, and one observed a higher number of infections.

Table 2. Overall healthcare-associated infections standardized infection ratios, Jan 1–Dec 31, 2018


Hospital Name	Observed Number of Infections	Predicted Number of Infections	SIR	Confidence Interval	Comparison to Expected Number of Infections
Alice Peck Day Memorial Hospital	2	1.15	1.74	0.29 , 5.75	Similar
Androscoggin Valley Hospital	†	†	†	†	†
Catholic Medical Center	17	21.78	0.78	0.47 , 1.22	Similar
Cheshire Medical Center	8	4.62	1.73	0.80 , 3.29	Similar
Concord Hospital	11	14.58	0.75	0.40 , 1.31	Similar
Cottage Hospital	†	†	†	†	†
DHMC	92	68.04	1.35	1.10 , 1.65	Higher
Elliot Hospital	8	15.78	0.51	0.24 , 0.96	Lower
Exeter Hospital	7	5.85	1.20	0.52 , 2.37	Similar
Franklin Regional Hospital	†	†	†	†	-
Frisbie Memorial Hospital	5	4.30	1.16	0.43 , 2.58	Similar
Huggins Hospital	1	1.01	0.99	0.05 , 4.88	Similar
Lakes Region General Hospital	2	5.39	0.37	0.06 , 1.23	Similar
Littleton Regional Hospital	3	2.10	1.43	0.36 , 3.89	Similar
Monadnock Community Hospital	†	†	†	†	†
New London Hospital	†	†	†	†	†
Parkland Medical Center	7	4.19	1.67	0.73 , 3.31	Similar
Portsmouth Regional Hospital	17	13.02	1.31	0.79 , 2.05	Similar
Southern NH Medical Center	4	7.93	0.50	0.16 , 1.22	Similar
Speare Memorial Hospital	2	1.16	1.72	0.29 , 5.70	Similar
St. Joseph Hospital	0	4.67	0.00	- , 0.78	Lower
The Memorial Hospital	†	†	†	†	†
Upper Connecticut Valley Hospital	†	†	†	†	-
Valley Regional Hospital	†	†	†	†	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass Hospital	6	9.00	0.67	0.27 , 1.39	Similar
State Total	196	188.89	1.04	0.90 , 1.19	Similar

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not report any data contributing to a SIR during this time period.

* Observed number of infections includes all infections that are required to be reported (central line-associated bloodstream infections, catheter-associated urinary tract infections, and surgical site infections following coronary artery bypass, colon, abdominal hysterectomy, and knee arthroplasty procedures).

Figure 3. Overall healthcare-associated infections standardized infection ratios, Jan 1–Dec 31, 2018



Note: Data are not shown for hospitals with less than one predicted infection. Observed number of infections includes all infections that are required to be reported (central line-associated bloodstream infections, catheter-associated urinary tract infections and surgical site infections following coronary artery bypass, colon, abdominal hysterectomy, and knee arthroplasty procedures).


Overall Statewide Standardized Infection Ratios: Comparison to 2017 Data

Table 3 shows that the statewide SIR in 2018 increased in comparison to 2017; however, this difference was not statistically significant. In 2018, a total of 196 HAI were reported, representing 133 SSI, 20 CLABSI, and 43 CAUTI compared to 183 HAI (116 SSI, 19 CLABSI, and 48 CAUTI) in 2017.

Table 3. Overall healthcare-associated infections standardized infection ratios, comparison between 2017 and 2018

Hospital	Standardized Infection Ratio (SIR) 2018	95% Confidence Interval 2018	Standardized Infection Ratio (SIR) 2017	95% Confidence Interval 2017	2018 Compared to 2017
Overall HAI SIR	1.04	0.90 , 1.19	0.99	0.86 , 1.15	Similar
CLABSI SIR	0.73	0.46 , 1.11	0.75	0.47 , 1.15	Similar
CAUTI SIR	1.16	0.85 , 1.55	1.37	1.02 , 1.80	Similar
Overall SSI SIR	1.07	0.90 , 1.26	0.94	0.78 , 1.12	Similar
CABG SIR	0.91	0.48 , 1.58	0.34	0.11 , 0.82	Similar
COLO SIR	1.01	0.80 , 1.27	1.04	0.82 , 1.30	Similar
HYST SIR	1.06	0.62 , 1.71	0.94	0.56 , 1.50	Similar
KPRO SIR	1.36	0.95 , 1.88	0.92	0.60 , 1.36	Similar

Figure 4. Statewide standardized infection ratios, comparison between 2017 and 2018



HAI: Healthcare-associated infection, CLABSI: Central line-associated blood stream infections, CAUTI: Catheter-associated urinary tract infections, SSI: Surgical site infections, CABG: SSI associated with coronary artery bypass graft procedures, COLO: SSI associated with colon procedures, HYST: SSI associated with abdominal hysterectomy procedures, KPRO: SSI associated with knee arthroplasty procedures

Table 4. Overall healthcare-associated infections standardized infection ratios by hospital, comparison between 2017 and 2018

Hospital	Standardized Infection Ratio (SIR) 2018	95% Confidence Interval 2018	Standardized Infection Ratio (SIR) 2017	95% Confidence Interval 2017	2017 Compared to 2018
Alice Peck Day Memorial	1.74	0.29 , 5.75	†	†	†
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	0.78	0.47 , 1.22	0.66	0.35 , 1.12	Similar
Cheshire Medical Center	1.73	0.80 , 3.29	1.18	0.32 , 3.02	Similar
Concord Hospital	0.75	0.40 , 1.31	0.64	0.30 , 1.17	Similar
Cottage Hospital	†	†	†	†	†
DHMC	1.35	1.10 , 1.65	1.04	0.81 , 1.31	Similar
Elliot Hospital	0.51	0.24 , 0.96	0.74	0.38 , 1.29	Similar
Exeter Hospital	1.20	0.52 , 2.37	0.82	0.27 , 1.92	Similar
Franklin Regional	†	†	†	†	†
Frisbie Memorial	1.16	0.43 , 2.58	1.22	0.39 , 2.85	Similar
Huggins Hospital	0.99	0.05 , 4.88	†	†	†
Lakes Region General	0.37	0.06 , 1.23	1.29	0.47 , 2.81	Similar
Littleton Regional	1.43	0.36 , 3.89	0.57	0.01 , 3.20	Similar
Monadnock Community	†	†	†	†	†
New London Hospital	†	†	†	†	†
Parkland Medical Center	1.67	0.73 , 3.31	2.01	0.86 , 3.95	Similar
Portsmouth Regional	1.31	0.79 , 2.05	1.04	0.54 , 1.82	Similar
Southern NH Medical	0.50	0.16 , 1.22	1.01	0.41 , 2.09	Similar
Speare Memorial Hospital	1.72	0.29 , 5.70	†	†	†
St. Joseph Hospital	0.00	- , 0.78	0.96	0.31 , 2.23	Similar
The Memorial Hospital	†	†	-	-	N/A
Upper Connecticut Valley	†	†	†	†	†
Valley Regional Hospital	†	†	†	†	†
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	0.67	0.27 , 1.39	1.01	0.48 , 1.86	Similar
State Total	1.04	0.90 , 1.19	0.99	0.86 , 1.15	Similar

† Data are not shown for hospitals with less than one predicted infection.

N/A or not applicable: Comparison between two years of data at a given facility is not applicable if no data were reported by that facility and/or if data must be censored for one or more of the years presented.

C. Central Line-Associated Bloodstream Infections

Table 5 shows the number of CLABSI identified in ICU at each hospital in NH. Among ICU with sufficiently robust data to present, 20 ICU observed a similar CLABSI rate to national rates. As shown in Table 6, two hospital observed similar CLABSI rates for all birthweight categories in comparison to the national data for Birthweight categories A, B, C, D, and E⁶. See methods for additional information on data collection.

⁶ Birthweight Category A \leq 750 grams, Birthweight Category B =751-1000 grams, Birthweight Category C =1001-1500 grams, Birthweight Category D =1501 2500 grams, and Birthweight Category E $>$ 2500 grams

Table 5. Central line-associated bloodstream infections rates, Jan 1–Dec 31, 2018

Hospital Name	Type of Unit	Observed Infections	Number of Central line Days	Hospital Rate	National Rate	P-Value	Hospital Rate Compared to National Rate
Androscoggin Valley Hospital	Medical ICU	†	†	†	†	†	†
Catholic Medical Center	Med/Surg ICU	1	2,969	0.3	0.8	0.387	Similar
Cheshire Medical Center	Medical ICU	1	488	2.0	1.1	0.508	Similar
Concord Hospital	Med/Surg ICU	1	1,978	0.5	0.8	0.714	Similar
Cottage Hospital	Med/Surg ICU	†	†	†	†	†	†
Dartmouth-Hitchcock Medical Center	Cardiac ICU	4	2,138	1.9	1.0	0.244	Similar
	Medical ICU	2	2,479	0.8	1.2	0.621	Similar
	Medical ICU	1	689	1.5	1.2	0.770	Similar
	Med/Surg Ped ICU	1	426	2.3	1.2	0.506	Similar
	Surg ICU	3	2,719	1.1	1.1	0.970	Similar
Elliot Hospital	Med/Surg ICU	0	1,381	0.0	0.8	0.322	Similar
Exeter Hospital	Med/Surg ICU	1	804	1.2	0.8	0.624	Similar
Frisbie Memorial Hospital	Med/Surg ICU	0	539	0.0	0.8	0.643	Similar
Huggins Hospital	Med/Surg ICU	0	52	0.0	0.8	0.958	Similar
Lakes Region General Hospital	Med/Surg ICU	0	338	0.0	0.8	0.758	Similar
Littleton Regional Hospital	Med/Surg ICU	0	112	0.0	0.8	0.912	Similar
Parkland Medical Center	Medical ICU	1	1,130	0.9	1.1	0.951	Similar
Portsmouth Regional Hospital	Med/Surg ICU	2	3,157	0.6	0.8	0.793	Similar
Southern NH Medical Center	Med/Surg ICU	0	521	0.0	0.8	0.652	Similar
Speare Memorial Hospital	Med/Surg ICU	0	67	0.0	0.8	0.947	Similar
St. Joseph Hospital	Med/Surg ICU	0	585	0.0	0.8	0.620	Similar
Weeks Medical Center	Med/Surg ICU	†	†	†	†	†	†
Wentworth-Douglass Hospital	Med/Surg ICU	2	1,457	1.4	0.8	0.454	Similar

Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections.

† Data are not shown for hospitals with fewer than 50 central line days. Med/Surg = medical surgical ICU = intensive care unit Ped = pediatric Surg=surgical

Table 6. Central line-associated bloodstream infections rates in neonatal intensive care units by birthweight category, Jan 1–Dec 31, 2018

Organization ID	Birth Weight Category	Observed Infections	Number of Device Days	Hospital Rate	National Rate	P-Value	Hospital Rate Compared to National Rate
Dartmouth-Hitchcock Medical Center	BW Category A < 751g	0	206	0	2.1	0.648	Similar
	BW Category B = 751-1000g	0	275	0	1.3	0.692	Similar
	BW Category C = 1001-1500g	0	306	0	0.8	0.787	Similar
	BW Category D = 1501-2500g	†	†	†	†	†	†
	BW Category E > 2500g	0	219	0	0.7	0.850	Similar
Elliot Hospital	BW Category A < 751g	0	79	0	2.1	0.847	Similar
	BW Category B = 751-1000g	0	288	0	1.3	0.680	Similar
	BW Category C = 1001-1500g	0	315	0	0.8	0.782	Similar
	BW Category D = 1501-2500g	0	569	0	0.6	0.725	Similar
	BW Category E > 2500g	0	345	0	0.7	0.775	Similar
Southern NH Medical Center	BW Category A < 751g	†	†	†	†	†	†
	BW Category B = 751-1000g	†	†	†	†	†	†
	BW Category C = 1001-1500g	†	†	†	†	†	†
	BW Category D = 1501-2500g	†	†	†	†	†	†
	BW Category E > 2500g	†	†	†	†	†	†

Note: DHMC, Elliot, and Southern NH Medical have neonatal intensive care units. All other hospitals do not and as such, had no data to report.

† Data are not shown for hospitals with fewer than 50 central line days for each birthweight category.

CLABSI Standardized Infection Ratios

Overall, the observed number of CLABSI was 27% fewer than predicted based on national data. The analysis presented in Table 7 and Figure 5 shows that six hospitals observed a similar number of infections as predicted based on national data.

Table 7. Central line-associated bloodstream infections standardized infection ratios, Jan 1–Dec 31, 2018


Hospital Name	Observed Infections	Predicted Infections	Standardized Infection Ratio	95% Confidence Interval	Comparison to Predicted Number of Infections
Alice Peck Day Memorial	-	-	-	-	-
Androscoggin Valley Hospital	†	†	†	†	†
Catholic Medical Center	1	3.35	0.30	0.02 , 1.47	Similar
Cheshire Medical Center	†	†	†	†	†
Concord Hospital	1	1.99	0.50	0.03 , 2.48	Similar
Cottage Hospital	†	†	†	†	†
DHMC	11	11.23	0.98	0.52 , 1.70	Similar
Elliot Hospital	0	2.93	0.00	- , 1.02	Similar
Exeter Hospital	†	†	†	†	†
Franklin Regional Hospital	-	-	-	-	-
Frisbie Memorial	†	†	†	†	†
Huggins Hospital	†	†	†	†	†
Lakes Region General Hospital	†	†	†	†	†
Littleton Regional Hospital	†	†	†	†	†
Monadnock Community	-	-	-	-	-
New London Hospital	-	-	-	-	-
Parkland Medical Center	†	†	†	†	†
Portsmouth Regional Hospital	2	2.74	0.73	0.12 , 2.41	Similar
Southern NH Medical Center	†	†	†	†	†
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	†	†	†	†	†
The Memorial Hospital	-	-	-	-	-
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	-	-	-	-	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass Hospital	2	1.10	1.82	0.31 , 6.02	Similar
State Total	20	27.38	0.73	0.46 , 1.11	Similar

Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections.

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not report any data contributing to a SIR during this time period.

Figure 5. Central line-associated bloodstream infections standardized infection ratios, Jan 1-Dec 31, 2018



Note: Data are not shown for hospitals with less than one predicted infection. Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections.

Central Line-Associated Bloodstream Infections: Comparison to 2017 Data

Overall, in 2018 the statewide CLABSI SIR was similar to 2017. The analysis presented in Table 8 shows that all six hospitals for which data are shown observed a similar number of infections in 2018 when compared to 2017. Figure 6 shows the CLABSI SIR between 2017 and 2018.

Table 8. Central line-associated bloodstream infections standardized infection ratios, comparison between 2017 and 2018


Hospital	Standardized Infection Ratio (SIR) 2018	95% Confidence Interval 2018	Standardized Infection Ratio (SIR) 2017	95% Confidence Interval 2017	2018 Compared to 2017
Alice Peck Day Memorial	-	-	-	-	-
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	0.30	0.02 , 1.47	0.70	0.12 , 2.31	Similar
Cheshire Medical Center	†	†	†	†	†
Concord Hospital	0.50	0.03 , 2.48	0.00	- , 1.92	Similar
Cottage Hospital	†	†	†	†	†
DHMC	0.98	0.52 , 1.70	0.84	0.43 , 1.50	Similar
Elliot Hospital	0.00	- , 1.02	0.37	0.02 , 1.83	Similar
Exeter Hospital	†	†	†	†	†
Franklin Regional	-	-	-	-	-
Frisbie Memorial	†	†	†	†	†
Huggins Hospital	†	†	†	†	†
Lakes Region General	†	†	†	†	†
Littleton Regional	†	†	†	†	†
Monadnock Community	-	-	-	-	-
New London Hospital	-	-	-	-	-
Parkland Medical Center	†	†	†	†	†
Portsmouth Regional	0.73	0.12 , 2.41	0.94	0.16 , 3.09	Similar
Southern NH Medical	†	†	†	†	†
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	†	†	†	†	†
The Memorial Hospital	-	-	-	-	-
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	-	-	-	-	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	1.82	0.31 , 6.02	0.00	- , 2.50	Similar
State Total	0.73	0.46 , 1.11	0.75	0.47 , 1.15	Similar

Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections.

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not report any data contributing to a SIR during this time period.

Figure 6. Overall central line-associated bloodstream infections standardized infection ratios by year, 2017-2018



D. Central Line Insertion Practices

CLIP monitoring assesses infection prevention practices that occur during the insertion of a central line. See methods section for information on monitoring CLIP.

Tables 9 through 11 and Figure 7 show the number of insertions during which all four infection prevention practices were appropriately followed, which is referred to as bundle adherence. A confidence interval is provided to assess any statistically significant differences in bundle adherence between groups.

The analysis presented in Table 9 suggests that, as an occupational group, interns and residents adhere to all four practices during central line insertions statistically less frequently than other occupations. As an occupational group, all other occupations adhere at a similar percentage when compared to the State adherence percentage. The analysis presented in Table 10 and Figure 7 show that, of the 11 hospitals with sufficiently robust data to present hospital-specific data, all 11 hospitals had similar adherence compared with the State adherence percentage.

Table 9. Central line insertion practices adherence percentages by occupation of inserter, Jan 1–Dec 31, 2018

Occupation of Inserter	Insertions that Adhered to Bundle*	Total Number of Insertions	% Adherence *	95% Confidence Interval	Occupation % Compared to State %
Advanced Practitce	403	414	97.3	95.4 , 98.6	Similar
Attending Physician	542	552	98.2	96.8 , 99.1	Similar
Fellow	181	182	99.5	97.3 , -	Similar
Intern/Resident	306	320	95.6	92.9 , 97.5	Lower
Other	†	†	†	†	†
Other Medical Staff	48	49	98.0	90.4 , 99.9	Similar
Physician Assistant	147	148	99.3	96.7 , -	Similar
Registered Nurse	862	868	99.3	98.6 , 99.7	Similar
State Total	2504	2548	98.3	97.7 , 98.7	

Note: Other Medical Staff represents other (non-attending) physicians.

* Bundle adherence refers to performing all four infection prevention practices during central line insertion.

† Data are not shown when fewer than 20 insertions were performed.

Table 10. Central line insertion practices adherence percentages by hospital, Jan 1–Dec 31, 2018

Hospital	Insertions that Adhered to Bundle*	Total Number of Insertions	% Adherence*	95% Confidence Interval	Hospital % Compared to State %
Alice Peck Day Memorial	-	-	-	-	-
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	54	54	100.0	94.6 , -	Similar
Cheshire Medical Center	55	55	100.0	94.7 , -	Similar
Concord Hospital	117	117	100.0	97.5 , -	Similar
Cottage Hospital	†	†	†	†	†
DHMC	1,085	1,112	97.6	96.5 , 98.4	Similar
Elliot Hospital	597	605	98.7	97.5 , 99.4	Similar
Exeter Hospital	85	88	96.6	91.0 , 99.1	Similar
Franklin Regional	-	-	-	-	-
Frisbie Memorial	†	†	†	†	†
Huggins Hospital	†	†	†	†	†
Lakes Region General	†	†	†	†	†
Littleton Regional	†	†	†	†	†
Monadnock Community	-	-	-	-	-
New London hospital	-	-	-	-	-
Parkland Medical Center	51	53	96.2	88.1 , 99.4	Similar
Portsmouth Regional	111	111	100.0	97.3 , -	Similar
Southern NH Medical	67	67	100.0	95.6 , -	Similar
Speare Memorial	†	†	†	†	†
St. Joseph Hospital	26	26	100.0	89.1 , -	Similar
The Memorial Hospital	-	-	-	-	-
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	-	-	-	-	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	142	144	98.6	95.5 , 99.8	Similar
State Total	2,504	2,548	98.3	97.7 , 98.7	


Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor insertion practices.

† Data are not shown when fewer than 20 insertions were performed.

* Bundle adherence refers to performing all four infection prevention practices during central line insertion.

- Facility did not report any data contributing to an adherence percentage during this time period.

Figure 7. Central line insertion practices adherence percentages by hospital, Jan 1–Dec 31, 2018



Note: Data are not shown when fewer than 20 insertions were performed. Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor insertion practices.

* Bundle adherence refers to performing all four infection prevention practices during central line insertion.

Central Line Insertion Practices: Comparison to 2017 Data

Overall, in 2018 the statewide adherence percentage for CLIP was similar to that in 2017. The analysis presented in Table 11 shows that CLIP adherence in 2018 was similar to 2017 for all occupations. Table 12 shows that, 11 hospitals with sufficiently robust data to present hospital-specific percentages, had similar CLIP adherence and 1 hospital had lower CLIP adherence in 2018 compared to 2017.

Table 11. Central line insertion practices adherence percentages by occupation of inserter, comparison between 2018 and 2017

Occupation of Inserter	% Adherence* 2018	95% Confidence Interval 2018	% Adherence* 2017	95% Confidence Interval 2017	2018 Compared to 2017
Advanced Practice Nurse	97.3	95.4 , 98.6	98.6	97.3 , 99.4	Similar
Attending Physician	98.2	96.8 , 99.1	97.7	96.2 , 98.7	Similar
Fellow	99.5	97.3 , -	98.5	96.0 , 99.7	Similar
Intern/Resident	95.6	92.9 , 97.5	98.8	97.1 , 99.6	Similar
Medical Student	†	†	†	†	†
Other	†	†	†	†	†
Other Medical Staff	98.0	90.4 , 99.9	99.2	96.0 , -	Similar
Physician Assistant	99.3	96.7 , -	98.7	95.7 , 99.8	Similar
Registered Nurse	99.3	98.6 , 99.7	99.2	98.4 , 99.6	Similar
State Total	98.3	97.7 , 98.7	98.5	98.1 , 99.0	Similar

Note: Other Medical Staff represents other (non-attending) physicians.

† Data are not shown when fewer than 20 insertions were performed.

* Bundle adherence refers to performing all four infection prevention practices during central line insertion.

N/A or not applicable: Comparison between two years of data at a given facility is not applicable if no data were reported by that facility and/or if data must be censored for one or more of the years presented.

- Facility did not report any data contributing to an adherence percentage during this time period.

Table 12. Central line insertion practices adherence percentages by hospital, comparison between 2018 and 2017

Hospital	% Adherence* 2018	95% Confidence Interval 2018	% Adherence* 2017	95% Confidence Interval 2017	2018 Compared to 2017
Alice Peck Day Memorial	-	-	-	-	-
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	100.0	94.6 , -	100.0	94.3 , -	Similar
Cheshire Medical Center	100.0	94.7 , -	95.0	77.7 , 99.8	Similar
Concord Hospital	100.0	97.5 , -	99.6	97.9 , -	Similar
Cottage Hospital	†	†	†	†	†
DHMC	97.6	96.5 , 98.4	99.1	98.5 , 99.5	Lower
Elliot Hospital	98.7	97.5 , 99.4	97.3	95.3 , 98.6	Similar
Exeter Hospital	96.6	91.0 , 99.1	100.0	97.3 , -	Similar
Franklin Regional	-	-	-	-	-
Frisbie Memorial	†	†	†	†	†
Huggins Hospital	†	†	100.0	87.8 , -	N/A
Lakes Region General	†	†	†	†	†
Littleton Regional	†	†	†	†	†
Monadnock Community	-	-	-	-	-
New London Hospital	-	-	-	-	-
Parkland Medical Center	96.2	88.1 , 99.4	92.4	87.1 , 95.9	Similar
Portsmouth Regional	100.0	97.3 , -	99.1	97.1 , 99.9	Similar
Southern NH Medical	100.0	95.6 , -	100.0	96.5 , -	Similar
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	100.0	89.1 , -	93.5	83.3 , 98.3	Similar
The Memorial Hospital	-	-	-	-	†
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	-	-	-	-	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	98.6	95.5 , 99.8	100.0	97.7 , -	Similar
State Total	98.3	97.7 , 98.7	98.5	98.1 , 99.0	Similar

Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor insertion practices.

† Data are not shown when fewer than 20 insertions were performed.

* Bundle adherence refers to performing all four infection prevention practices during central line insertion.

N/A or not applicable: Comparison between two years of data at a given facility is not applicable if no data were reported by that facility and/or if data must be censored for one or more of the years presented.

- Facility did not report any data contributing to an adherence percentage during this time period.

E. Catheter-Associated Urinary Tract Infections

Tables 13 through 15 and Figure 8 show the number of infections that were identified in adult ICU at NH hospitals. The analysis presented in Table 13 shows the number of CAUTI identified in individual ICU by hospitals in NH. Among ICU with sufficiently robust data to present, all individual ICU observed a similar CAUTI rate based on national data. See methods for additional information on data collection.

Table 13. Catheter-associated urinary tract infection rates, Jan 1–Dec 31, 2018

	Unit Type	Infections	Catheter days	Hospital Rate	National Rate	P-value	Hospital Rate Compared to National Rate
Androscoggin Valley Hospital	Medical ICU	0	163	0	2.0	0.717	Similar
Catholic Medical Center	Med/Surg ICU	3	3,271	0.9	1.7	0.284	Similar
Cheshire Medical Center	Medical ICU	0	989	0	2.0	0.133	Similar
Concord Hospital	Med/Surg ICU	4	3,223	1.2	1.7	0.572	Similar
Cottage Hospital	Med/Surg ICU	0	81	0	1.3	0.902	Similar
Dartmouth-Hitchcock Medical Center	Cardiac ICU	5	2,193	2.3	2.3	0.936	Similar
	Medical ICU	3	3,046	1.0	3.5	0.009	Lower
	Medical ICU	2	1,087	1.8	3.5	0.382	Similar
	Med/Surg Pediatric ICU	0	188	0	2.5	0.629	Similar
	Surg ICU	14	4,040	3.5	3.4	0.933	Similar
Elliot Hospital	Med/Surg ICU	1	1,818	0.6	1.7	0.234	Similar
Exeter Hospital	Med/Surg ICU	1	857	1.2	1.3	0.961	Similar
Frisbie Memorial Hospital	Med/Surg ICU	2	884	2.3	1.3	0.414	Similar
Huggins Hospital	Med/Surg ICU	0	176	0	1.3	0.799	Similar
Lakes Region General Hospital	Med/Surg ICU	0	1,120	0	1.3	0.241	Similar
Littleton Regional Hospital	Med/Surg ICU	0	348	0	1.3	0.642	Similar
Parkland Medical Center	Medical ICU	1	1,257	0.8	2.0	0.351	Similar
Portsmouth Regional Hospital	Med/Surg ICU	6	3,608	1.7	1.3	0.493	Similar
Southern NH Medical Center	Med/Surg ICU	1	1,378	0.7	1.7	0.421	Similar
Speare Memorial Hospital	Med/Surg ICU	0	213	0	1.3	0.763	Similar
St. Joseph Hospital	Med/Surg ICU	0	1,156	0	1.3	0.230	Similar
Weeks Medical Center	Med/Surg ICU	0	75	0	1.3	0.909	Similar
Wentworth-Douglass Hospital	Med/Surg ICU	0	1,526	0	1.3	0.144	Similar

Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections.

† Data are not shown for hospitals with fewer than 50 catheter days.

Med/Surg = medical surgical ICU = intensive care unit Ped=pediatric Surg=surgical ICU

Catheter-Associated Urinary Tract Infections Standardized Infection Ratios

The observed number of CAUTI was 16% higher than predicted based on national data, this was not statistically significant. The analysis presented in Table 14 shows that six hospitals observed a similar number of infections as predicted based on national data.

Table 14. Catheter-associated urinary tract infections standardized infection ratios, Jan 1–Dec 31, 2018


Hospital	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted Number of Infections
Alice Peck Day Memorial	-	-	-	-	-
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	3	4.26	0.70	0.18 , 1.92	Similar
Cheshire Medical Center	†	†	†	†	†
Concord Hospital	4	3.29	1.22	0.39 , 2.93	Similar
Cottage Hospital	†	†	†	†	†
DHMC	24	16.68	1.44	0.94 , 2.11	Similar
Elliot Hospital	1	1.63	0.61	0.03 , 3.03	Similar
Exeter Hospital	†	†	†	†	†
Franklin Regional	†	†	†	†	†
Frisbie Memorial	†	†	†	†	†
Huggins Hospital	†	†	†	†	†
Lakes Region General	†	†	†	†	†
Littleton Regional Hospital	†	†	†	†	†
Monadnock Hospital	-	-	-	-	-
New London Hospital	-	-	-	-	-
Parkland Medical Center	†	†	†	†	†
Portsmouth Regional	6	3.23	1.86	0.75 , 3.86	Similar
Southern NH Medical Center	1	1.01	0.99	0.05 , 4.90	Similar
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	†	†	†	†	†
The Memorial Hospital	-	-	-	-	-
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	-	-	-	-	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	0	1.12	0.00	- , 2.69	Similar
State total	43	36.98	1.16	0.85 , 1.55	Similar

Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections.

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not report any data contributing to a SIR during this time period.

Figure 8. Catheter-associated urinary tract infections standardized infection ratios, Jan 1–Dec 31, 2018



Note: Data are not shown for hospitals with less than one predicted infection. Alice Peck Day Memorial, Androscoggin Valley Hospital, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections.

Catheter-Associated Urinary Tract Infections: Comparison to 2017 Data

Overall, in 2018 the statewide CAUTI SIR was similar compared to 2017. The analysis in Table 15 shows that 6 hospitals with robust data observed similar number of infections in 2018 when compared to 2017. Figure 9 compares the 2018 statewide CAUTI SIR to 2017.

Table 15. Catheter-associated urinary tract infections standardized infection ratios, comparison between 2018 and 2017


Hospital	Standardized Infection Ratio (SIR) 2018	95% Confidence Interval 2018	Standardized Infection Ratio (SIR) 2017	95% Confidence Interval 2017	2018 Compared to 2017
Alice Peck Day Memorial	-	-	-	-	-
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	0.70	0.18 , 1.92	0.33	0.02 , 1.64	Similar
Cheshire Medical Center	†	†	†	†	†
Concord Hospital	1.22	0.39 , 2.93	1.53	0.49 , 3.69	Similar
Cottage Hospital	†	†	†	†	†
DHMC	1.44	0.94 , 2.11	1.47	0.99 , 2.11	Similar
Elliot Hospital	0.61	0.03 , 3.03	1.31	0.22 , 4.33	Similar
Exeter Hospital	†	†	†	†	†
Franklin Regional	†	†	-	-	-
Frisbie Memorial	†	†	†	†	†
Huggins Hospital	†	†	†	†	†
Lakes Region General	†	†	†	†	†
Littleton Regional	†	†	†	†	†
Monadnock Community	-	-	-	-	-
New London Hospital	-	-	-	-	-
Parkland Medical Center	†	†	†	†	†
Portsmouth Regional	1.86	0.75 , 3.86	1.65	0.60 , 3.65	Similar
Southern NH Medical	0.99	0.05 , 4.90	†	†	N/A
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	†	†	†	†	†
The Memorial Hospital	-	-	†	†	†
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	-	-	-	-	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	0.00	- , 2.69	2.98	0.94 , 7.18	Similar
State Total	1.16	0.85 , 1.55	1.37	1.02 , 1.80	Similar

Note: Alice Peck Day Memorial, Franklin Regional Hospital, Monadnock Community Hospital, New London Hospital, The Memorial Hospital, Upper Connecticut Valley Hospital and Valley Regional Hospital did not have an intensive care unit in which to monitor infections in 2018 and/or 2017.

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not report any data contributing to a SIR during this time period.

Figure 9. Overall catheter-associated urinary tract infections standardized infection ratios by year, 2017-2018



F. Surgical Site Infections

Tables 16-21 and Figures 10-14 below show the number of SSI following the four monitored procedures reported by each acute care hospital in NH. Overall, the observed number of SSI was 7% higher than predicted based on national data. The analysis presented in Table 16 shows that fifteen hospitals with robust data observed similar number of SSI as predicted. One hospital experienced a lower number of SSI, and one hospital experienced higher number of SSI as predicted when compared to national data. For CABG procedures (Table 18), all hospitals with sufficiently robust data to present observed a similar number of infections as predicted. For colon procedures (Table 19), twelve hospitals observed a similar number of infections, and one hospital observed a lower number of infections as predicted. For abdominal hysterectomy procedures (Table 20), all four hospitals with sufficiently robust data to present observed a similar number of infections as predicted based on national data. For knee arthroplasty procedures (Table 21), eight hospitals observed a similar number of infections, and one hospital observed a higher number of infections as predicted based on national data.

This report does not display SSI rates due to a change in analysis recommendations. SSI data are presented throughout this report as SIR. This allows more robust adjustment for underlying patient or hospital factors. The SSI SIR is calculated using logistic regression modeling, which provides better risk adjustment and more appropriate comparisons. See Appendix 1 for technical notes and more detail regarding the SIR.

Post-discharge Surveillance for Surgical Site Infections

Hospitals do not use a standard method to identify infections once a patient has been discharged (known as “post-discharge surveillance”). This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections. Table 22 shows the percentage of SSI identified through post-discharge surveillance at each acute care hospital in NH. Of the 26 hospitals with sufficiently robust data, 19 hospitals identified a similar number of SSI through post-discharge surveillance, three hospitals identified more SSI through post-discharge surveillance when compared to the State rolling average. Out of 298 SSI reported 2017 and 2018, 39.9% (119) were detected during admission, 6.4% (19) were detected during readmission at another facility, 37.6% (112) were detected during readmission to the original facility where the procedure took place, and 16.1% (48) were detected post-discharge. Most of the infections detected post-discharge were classified as intra-abdominal abscess 31.88% (95); 32.55% (97) were superficial infections and 16.11% (48) were deep infections. Of the 48 infections detected post-discharge, 50% (24) were colon procedure, 22.9% (11) were knee arthroplasty procedures, 25% (12) were abdominal hysterectomy procedures and 2.1% (1) were coronary artery bypass graft procedures. NH hospitals infection prevention staff rely primarily on follow-up letters to surgeons, culture reports, and outpatient clinic notes as forms of post-discharge surveillance. Other methods include patient letters and communication with other healthcare facilities.


Table 16. Surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018

Hospital	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted Number of Infections
Alice Peck Day Memorial	2	1.15	1.74	0.29 , 5.74	Similar
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	13	14.17	0.92	0.51 , 1.53	Similar
Cheshire Medical Center	7	3.52	1.99	0.87 , 3.93	Similar
Concord Hospital	6	9.30	0.65	0.26 , 1.34	Similar
Cottage Hospital	†	†	†	†	†
DHMC	57	40.13	1.42	1.08 , 1.82	Higher
Elliot Hospital	7	11.22	0.62	0.27 , 1.23	Similar
Exeter Hospital	5	4.62	1.08	0.39 , 2.39	Similar
Franklin Regional Hospital	-	-	-	-	-
Frisbie Memorial Hospital	3	3.29	0.91	0.23 , 2.47	Similar
Huggins Hospital	†	†	†	†	†
Lakes Region General	2	4.34	0.46	0.07 , 1.52	Similar
Littleton Regional Hospital	3	1.85	1.62	0.41 , 4.42	Similar
Monadnock Community	†	†	†	†	†
New London Hospital	†	†	†	†	†
Parkland Medical Center	5	2.73	1.83	0.67 , 4.06	Similar
Portsmouth Regional	9	7.05	1.28	0.62 , 2.34	Similar
Southern NH Medical	3	6.49	0.46	0.11 , 1.25	Similar
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	0	3.39	0.00	- , 0.88	Lower
The Memorial Hospital	†	†	†	†	†
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	†	†	†	†	†
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	4	6.78	0.59	0.18 , 1.42	Similar
State Total	133	124.53	1.07	0.89 , 1.26	Similar

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not perform any of the four procedures being monitored during 2018

Figure 10. Surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018



Note: Data are not shown for hospitals with less than one predicted infection.

Table 17. Surgical site infections standardized infection ratios, comparison between 2017 and 2018

Hospital	Standardized Infection Ratio (SIR) 2018	95% Confidence Interval 2018	Standardized Infection Ratio (SIR) 2017	95% Confidence Interval 2017	2018 Compared to 2017
Alice Peck Day Memorial	1.74	0.29 , 5.74	†	†	N/A
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	0.92	0.51 , 1.53	0.72	0.36 , 1.28	Similar
Cheshire Medical Center	1.99	0.87 , 3.93	1.14	0.29 , 3.11	Similar
Concord Hospital	0.65	0.26 , 1.34	0.52	0.21 , 1.08	Similar
Cottage Hospital	†	†	†	†	†
DHMC	1.42	1.08 , 1.82	0.91	0.64 , 1.24	Similar
Elliot Hospital	0.62	0.27 , 1.23	0.77	0.38 , 1.41	Similar
Exeter Hospital	1.08	0.39 , 2.39	0.85	0.27 , 2.05	Similar
Franklin Regional	-	-	-	-	-
Frisbie Memorial	0.91	0.23 , 2.47	0.91	0.23 , 2.47	Similar
Huggins Hospital	†	†	†	†	†
Lakes Region General	0.46	0.07 , 1.52	1.34	0.49 , 2.96	Similar
Littleton Regional	1.62	0.41 , 4.42	0.61	0.03 , 3.03	Similar
Monadnock Community	†	†	†	†	†
New London Hospital	†	†	†	†	†
Parkland Medical Center	1.83	0.67 , 4.06	2.52	1.10 , 4.99	Similar
Portsmouth Regional	1.28	0.62 , 2.34	0.79	0.29 , 1.75	Similar
Southern NH Medical	0.46	0.11 , 1.25	0.70	0.22 , 1.70	Similar
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	0.00	- , 0.88	1.19	0.44 , 2.64	Similar
The Memorial Hospital	†	†	†	†	†
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	†	†	-	-	N/A
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	0.59	0.18 , 1.42	0.82	0.33 , 1.70	Similar
State Total	1.07	0.89 , 1.26	0.94	0.78 , 1.12	Similar

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not perform any of the four procedures being monitored during 2017 or 2018

N/A or not applicable: Comparison between two years of data at a given facility is not applicable if no data were reported by that facility and/or if data must be censored for one or more of the years presented.

Table 18. Coronary artery bypass graft procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018

Hospital	Observed Infections	Predicted Infections	Standardized Infection Ratio	95% Confidence Interval	Comparison to Expected Number of Infections
Catholic Medical Center	4	6.79	0.59	0.19 , 1.42	Similar
Concord Hospital	†	†	†	†	†
Dartmouth-Hitchcock Medical Center	2	3.15	0.64	0.11 , 2.10	Similar
Portsmouth Regional Hospital	5	2.19	2.29	0.84 , 5.07	Similar
State Total	11	12.12	0.91	0.48 , 1.58	Similar

Figure 11. Coronary artery bypass graft procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018





Table 19. Colon procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018

Hospital	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted Number of Infections
Alice Peck Day Memorial	-	-	-	-	-
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	7	6.22	1.13	0.49, 2.22	Similar
Cheshire Medical Center	5	2.09	2.39	0.87, 5.29	Similar
Concord Hospital	1	4.50	0.22	0.01, 1.09	Similar
Cottage Hospital	†	†	†	†	†
DHMC	36	25.67	1.40	1.00, 1.92	Similar
Elliot Hospital	2	8.19	0.24	0.04, 0.80	Lower
Exeter Hospital	4	3.50	1.14	0.36, 2.75	Similar
Franklin Regional Hospital	-	-	-	-	-
Frisbie Memorial Hospital	1	2.21	0.45	0.02, 2.22	Similar
Huggins Hospital	†	†	†	†	†
Lakes Region General	2	2.60	0.77	0.12, 2.54	Similar
Littleton Regional	†	†	†	†	†
Monadnock Community	†	†	†	†	†
New London Hospital	†	†	†	†	†
Parkland Medical Center	3	1.34	2.23	0.56, 6.07	Similar
Portsmouth Regional	4	3.15	1.27	0.40, 3.06	Similar
Southern NH Medical	3	4.02	0.75	0.19, 2.02	Similar
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	0	2.31	0.00	- , 1.29	Similar
The Memorial Hospital	†	†	†	†	†
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	-	-	-	-	-
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	1	4.20	0.24	0.01, 1.17	Similar
State Total	74	73.01	1.01	0.80 , 1.27	Similar

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not perform this procedure during 2018

Figure 12. Colon procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018



Note: Data are not shown for hospitals with less than one predicted infection.


Table 20. Abdominal hysterectomy procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018

Hospital	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted Number of Infections
Alice Peck Day Memorial	†	†	†	†	†
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	†	†	†	†	†
Cheshire Medical Center	†	†	†	†	†
Concord Hospital	†	†	†	†	†
Cottage Hospital	-	-	-	-	-
DHMC	11	6.28	1.75	0.92 , 3.04	Similar
Elliot Hospital	0	1.56	0.00	- , 1.92	Similar
Exeter Hospital	†	†	†	†	†
Franklin Regional Hospital	-	-	-	-	-
Frisbie Memorial Hospital	†	†	†	†	†
Huggins Hospital	†	†	†	†	†
Lakes Region General	†	†	†	†	†
Littleton Regional Hospital	†	†	†	†	†
Monadnock Community	†	†	†	†	†
New London Hospital	†	†	†	†	†
Parkland Medical Center	†	†	†	†	†
Portsmouth Regional	†	†	†	†	†
Southern NH Medical	0	1.11	0.00	- , 2.71	Similar
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	†	†	†	†	†
The Memorial Hospital	†	†	†	†	†
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	†	†	†	†	†
Weeks Medical Center	-	-	-	-	-
Wentworth-Douglass	1	1.04	0.96	0.05 , 4.73	Similar
State Total	15	14.17	1.06	0.62 , 1.70	Similar

† Data are not shown for hospitals with less than one predicted infection.

- Facility did not perform this procedure during 2018

Figure 13. Abdominal hysterectomy procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018



Note: Data are not shown for hospitals with less than one predicted infection


Table 21. Knee arthroplasty procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018

Hospital	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted Number of Infections
Alice Peck Day Memorial	2	1.06	1.88	0.32 , 6.22	Similar
Androscoggin Valley	†	†	†	†	†
Catholic Medical Center	†	†	†	†	†
Cheshire Medical Center	†	†	†	†	†
Concord Hospital	5	3.45	1.45	0.53 , 3.22	Similar
Cottage Hospital	†	†	†	†	†
DHMC	8	5.03	1.59	0.74 , 3.02	Similar
Elliot Hospital	5	1.47	3.40	1.25 , 7.53	Higher
Exeter Hospital	1	1.02	0.98	0.05 , 4.83	Similar
Franklin Regional Hospital	-	-	-	-	-
Frisbie Memorial Hospital	†	†	†	†	†
Huggins Hospital	†	†	†	†	†
Lakes Region General	0	1.44	0.00	- , 2.09	Similar
Littleton Regional Hospital	†	†	†	†	†
Monadnock Community	†	†	†	†	†
New London Hospital	†	†	†	†	†
Parkland Medical Center	†	†	†	†	†
Portsmouth Regional	0	1.56	0.00	- , 1.93	Similar
Southern NH Medical Center	0	1.36	0.00	- , 2.20	Similar
Speare Memorial Hospital	†	†	†	†	†
St. Joseph Hospital	†	†	†	†	†
The Memorial Hospital	†	†	†	†	†
Upper Connecticut Valley	-	-	-	-	-
Valley Regional Hospital	†	†	†	†	†
Weeks Medical Center	†	†	†	†	†
Wentworth-Douglass	2	1.54	1.30	0.22 , 4.29	Similar
State Total	33	24.32	1.36	0.95 , 1.88	Similar

† Data are not shown for hospitals with less than one predicted infection

- Facility did not perform this procedure during 2018

Figure 14. Knee arthroplasty procedure-associated surgical site infections standardized infection ratios, Jan 1–Dec 31, 2018



Note: Data are not shown for hospitals with less than one predicted infection

Overall Surgical Site Infections: Comparison to 2017 Data

Overall, in 2018 the statewide SSI SIR was similar to 2017. The analysis presented in Table 17 (above) indicates that all hospitals for which data are shown have similar numbers of infections observed in 2017 and 2018.

Figures 16-19 show the SIR for each procedure that was reportable from 2017-2018. There was a decrease in the SIR for colon (Figure 17); there was an increase in SIR for coronary bypass graft procedure (Figure 16), abdominal hysterectomy (Figure 18) procedures, and KPRO procedures (Figure 19) from 2017 to 2018; however, these differences are not statistically significant. NH had a similar SSI SIR than predicted when compared to national data, and no significant change across reporting years.

Figure 15. Overall surgical site infection standardized infection ratios by year, 2017-2018




Figure 16. Overall coronary artery bypass graft procedure standardized infection ratios by year, 2017-2018




Figure 17. Overall colon procedure standardized infection ratios by year, 2017-2018




Figure 18. Overall abdominal hysterectomy standardized infection ratios by year, 2017-2018




Figure 19. Overall knee arthroplasty standardized infection ratios by year, 2017-2018




Table 22. Post-discharge surveillance methods and percentage of SSI detected post-discharge in New Hampshire hospitals, 2017-2018

Hospital	Post-Discharge Surveillance Methods	% SSIs Identified Post-Discharge	Compared to State
Alice Peck Day Memorial	Surgeon Letters	50.00	Similar
Androscoggin Valley	Surgeon Letters, Culture Reports, Outpatient Clinic	0.00	Similar
Catholic Medical Center	Surgeon Letters, Culture Reports	7.14	Similar
Cheshire Medical Center	Culture Reports, Outpatient Clinic	21.43	Similar
Concord Hospital	Culture Reports	0.00	Similar
Cottage Hospital	Surgeon Letters, Culture Reports, Outpatient Clinic, other*	0.00	Similar
DHMC	Surgeon Letters, Culture Reports, Outpatient Clinic	13.08	Similar
Elliot Hospital	Surgeon Letters, Culture Reports	18.18	Similar
Exeter Hospital	Surgeon Letters	0.00	Similar
Franklin Regional	Surgeon Letters, Culture Reports	†	†
Frisbie Memorial	Surgeon Letters, Outpatient Clinic	55.56	Higher
Huggins Hospital	Surgeon Letters, Culture Reports, Patient/Family	33.33	Similar
Lakes Region General	Surgeon Letters, Culture Reports	25.00	Similar
Littleton Regional	Surgeon Letters, Culture Reports, Outpatient Clinic	20.00	Similar
Monadnock Community	Surgeon Letters, Culture Reports	100.00	Similar
New London Hospital	Surgeon Letters, Culture Reports, Outpatient Clinic	66.67	Similar
Parkland Medical Center	Surgeon Letters, Culture Reports	8.33	Similar
Portsmouth Regional	Surgeon Letters	5.56	Similar
Southern NH Medical	Surgeon Letters, Culture Reports	0.00	Similar
Speare Memorial Hospital	Surgeon Letters	25.00	Similar
St. Joseph Hospital	Surgeon Letters, Culture Reports, Outpatient Clinic	100.00	Higher
The Memorial Hospital	Surgeon Letters, Culture Reports, Other**	100.00	Higher
Upper Connecticut Valley	Surgeon Letters, Culture Reports, Patient/Family	†	†
Valley Regional Hospital	Surgeon Letters, Culture Reports, Outpatient Clinic	†	†
Weeks Medical Center	Surgeon Letters, Culture Reports, Outpatient Clinic	†	†
Wentworth-Douglass	Surgeon Letters, Culture Reports	9.09	Similar

†No SSIs reported or predicted number of infections is less than one during this time period. Note: Post-discharge surveillance methods may have changed since originally reported. These data are for 2017 and 2018 and are not directly comparable to the rest of the data in this report. These data are shown to assess the effectiveness of the post-discharge surveillance system implemented at each facility. Two years of data are used since the number of reported infections at many facilities is small.

G. Surgical Antimicrobial Prophylaxis Administration


SCIP 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report. In 2013, NH hospitals performed surgical antimicrobial prophylaxis correctly more often or similar to the national adherence percentage. For SCIP measure 1, 98.6% of patients received prophylactic antibiotic within one hour prior to surgery compared with 98.5% nationally. For SCIP measure 2, 99.3% of patients received the appropriate prophylactic antibiotic compared with 98.9% nationally. For SCIP measure 3, 98.0% of patients had his or her prophylactic antibiotic discontinued within 24 hours after surgery compared with 97.6% nationally. See methods section for additional information on how this information is collected.

Data for 2013 and earlier years is available in prior HAI Program reports and can be accessed here: <http://www.dhhs.nh.gov/dphs/cdcs/hai/publications.htm>.

H. Influenza Vaccination Percentages

Figure 20 shows 2018-19 influenza season, NH State observed an increase in influenza vaccination from the year prior. The State observed an overall increase in HCP vaccination percentages in NH hospitals between 2008-09 to 2018-19; 2008-09 to 2013-14 represents a statistically significant increase from the year prior. Table 23 and Figure 21 show the total number of HCP and the number of HCP vaccinated against seasonal influenza at each hospital during the 2018–19 influenza season. Vaccination percentages by hospital ranged from 65.9% to 100%, and the overall State vaccination percentage was 94.0%. The analysis presented in Table 23 shows that ten hospitals had vaccination percentages similar to the overall State vaccination percentage, thirteen hospitals reported vaccination percentages that were significantly higher than the overall State vaccination percentage, and ten hospitals reported vaccination percentages that were significantly lower than the overall State vaccination percentage.

Figure 20. Statewide influenza vaccination percentages for hospital HCP by influenza season




Note: Influenza season represents data for HCP between October 1st and March 31st the following calendar year, with the exception of 2008-09, which data were collected for October 1st through April 30th.

Table 23. Influenza vaccination percentages for hospital HCP by hospital, 2018–19 influenza season, Oct 1, 2018–Mar 31, 2019

Hospital	Total HCP	HCP Vaccinated	% HCP Vaccinated	95% Confidence Interval	Hospital % Compared to State %
Alice Peck Day Memorial Hospital	1,130	1,065	94.3	92.8 , 95.5	Similar
Androscoggin Valley Hospital	510	485	95.1	93.0 , 96.7	Similar
Catholic Medical Center	3,743	3,697	98.8	98.4 , 99.1	Higher
Cheshire Medical Center	1,599	1,467	91.7	90.3 , 93.0	Lower
Concord Hospital	4,997	4,707	94.2	93.5 , 94.8	Similar
Cottage Hospital	470	323	68.7	64.4 , 72.8	Lower
Dartmouth-Hitchcock Medical Center	9,348	8,938	95.6	95.1 , 96.0	Higher
Elliot Hospital	6,584	6,524	99.1	98.8 , 99.3	Higher
Exeter Hospital	2,232	2,069	92.7	91.6 , 93.7	Lower
Franklin Regional Hospital	167	155	92.8	88.1 , 96.1	Similar
Frisbie Memorial Hospital	1,678	1,597	95.2	94.1 , 96.1	Similar
Huggins Hospital	767	713	93.0	91.0 , 94.6	Similar
Lakes Region General Hospital	1,328	1,272	95.8	94.6 , 96.8	Higher
Littleton Regional Hospital	760	750	98.7	97.7 , 99.3	Higher
Monadnock Community Hospital	750	697	92.9	90.9 , 94.6	Similar
New London Hospital	694	675	97.3	95.8 , 98.3	Higher
Parkland Medical Center	1,249	1,153	92.3	90.7 , 93.7	Lower
Portsmouth Regional Hospital	2,638	2,380	90.2	89.0 , 91.3	Lower
Southern NH Medical Center	2,512	2,450	97.5	96.9 , 98.1	Higher
Speare Memorial Hospital	784	710	90.6	88.4 , 92.5	Lower
St. Joseph Hospital	2,474	2,410	97.4	96.7 , 98.0	Higher
The Memorial Hospital	627	538	85.8	82.9 , 88.4	Lower
Upper Connecticut Valley Hospital	224	204	91.1	86.8 , 94.3	Similar
Valley Regional Hospital	442	440	99.6	98.5 , 99.9	Higher
Weeks Medical Center	451	441	97.8	96.1 , 98.9	Higher
Wentworth-Douglass Hospital	3,782	3,146	83.2	82.0 , 84.4	Lower
Encompass Health Rehabilitation	258	239	92.6	88.9 , 95.4	Similar
Hampstead Hospital	208	137	65.9	59.2 , 72.1	Lower
New Hampshire Hospital	1,294	1,021	78.9	76.6 , 81.1	Lower
Northeast Rehabilitation at Pease	244	243	99.6	98.0 , -	Higher
Northeast Rehabilitation at Salem	711	707	99.4	98.6 , 99.8	Higher
Northeast Rehabilitation at SNHMC	101	100	99.0	95.2 , -	Higher
Northeast Rehabilitation at The Elliot	64	65	98.46	92.7 , 99.9	Similar
State Total	54820	51518	94.0	93.8 , 94.2	

Figure 21. Influenza vaccination percentages for hospital HCP by hospital, 2018-19 influenza season, Oct 1, 2017– Mar 31, 2019



Influenza Vaccination Percentages: Comparison to 2017-18 Data

The overall statewide hospital HCP vaccination percentage increased significantly from 2008–09 to 2009–10, which may have been explained by overall increased interest in influenza vaccination as a result of the 2009 H1N1 pandemic. However, the influenza vaccination percentage continued to increase between the 2009-10 and 2018-19 seasons, suggesting other influences such as the public reporting of influenza vaccination coverage and mandatory vaccination policies. The analysis presented in Table 24 shows that overall, five hospitals increased HCP influenza vaccination, 24 hospitals had similar vaccination percentages, and 4 hospitals decreased influenza vaccination percentages in 2018-19 compared to 2017-18.

Figure 22. Influenza vaccination percentages for hospital healthcare personnel by hospital, 2017-18 and 2018-19 influenza seasons




Table 24. Influenza vaccination percentages for hospital healthcare personnel by hospital, comparison between 2017-18 and 2018-19 influenza seasons

Hospital	% HCP Vaccinated 2018-19	95% Confidence Interval 2018-19	% HCP Vaccinated 2017-18	95% Confidence Interval 2017-18	2018-19 Compared to 2017-18
Alice Peck Day Memorial Hospital	94.3	92.8 , 95.5	94.8	93.2 , 96.1	Similar
Androscoggin Valley Hospital	95.1	93.0 , 96.7	94.7	92.6 , 96.4	Similar
Catholic Medical Center	98.8	98.4 , 99.1	98.9	98.6 , 99.2	Similar
Cheshire Medical Center	91.7	90.3 , 93.0	94.3	93.1 , 95.3	Lower
Concord Hospital	94.2	93.5 , 94.8	77.4	76.0 , 78.7	Higher
Cottage Hospital	68.7	64.4 , 72.8	91.3	88.5 , 93.6	Lower
Dartmouth-Hitchcock Medical Center	95.6	95.1 , 96.0	96.3	95.9 , 96.7	Similar
Elliot Hospital	99.1	98.8 , 99.3	99.1	98.8 , 99.3	Similar
Exeter Hospital	92.7	91.6 , 93.7	92.9	91.7 , 93.9	Similar
Franklin Regional Hospital	92.8	88.1 , 96.1	90.4	86.0 , 93.8	Similar
Frisbie Memorial Hospital	95.2	94.1 , 96.1	95.0	93.8 , 96.0	Similar
Encompass Health Rehabilitation	92.6	88.9 , 95.4	93.7	90.2 , 96.2	Similar
Huggins Hospital	93.0	91.0 , 94.6	89.1	86.6 , 91.2	Similar
Lakes Region General Hospital	95.8	94.6 , 96.8	97.2	96.2 , 98.0	Similar
Littleton Regional Hospital	98.7	97.7 , 99.3	96.2	94.6 , 97.4	Higher
Monadnock Community Hospital	92.9	90.9 , 94.6	96.2	94.6 , 97.4	Similar
New London Hospital	97.3	95.8 , 98.3	97.7	96.4 , 98.6	Similar
Parkland Medical Center	92.3	90.7 , 93.7	80.9	78.7 , 83.0	Higher
Portsmouth Regional Hospital	90.2	89.0 , 91.3	90.0	88.6 , 91.3	Similar
Southern NH Medical Center	97.5	96.9 , 98.1	98.0	97.3 , 98.5	Similar
Speare Memorial Hospital	90.6	88.4 , 92.5	89.5	87.0 , 91.6	Similar
St. Joseph Hospital	97.4	96.7 , 98.0	97.7	97.0 , 98.2	Similar
The Memorial Hospital	85.8	82.9 , 88.4	84.3	81.4 , 86.9	Similar
Upper Connecticut Valley Hospital	91.1	86.8 , 94.3	89.3	84.2 , 93.2	Similar
Valley Regional Hospital	99.6	98.5 , 99.9	100.0	99.2 , -	Similar
Weeks Medical Center	97.8	96.1 , 98.9	97.7	95.9 , 98.8	Similar
Wentworth-Douglass Hospital	83.2	82.0 , 84.4	96.3	95.6 , 96.9	Lower
Hampstead Hospital	65.9	59.2 , 72.1	73.0	66.8 , 78.6	Similar
New Hampshire Hospital	78.9	76.6 , 81.1	54.5	51.8 , 57.2	Higher
Northeast Rehabilitation at Pease (Portsmouth)	99.6	98.0 , -	99.1	96.9 , 99.8	Similar
Northeast Rehabilitation at Salem	99.4	98.6 , 99.8	99.5	98.8 , 99.8	Similar
Northeast Rehabilitation at SNHMC (Nashua)	99.0	95.2 , -	97.8	94.2 , 99.4	Similar
Northeast Rehabilitation at The Elliot (Manchester)	75.4	63.86 , 84.69	98.8	94.2 , 99.9	Lower
State Total	94.0	93.8 , 94.2	93.3	93.1 , 93.5	Higher

Influenza Vaccination Policies for Healthcare Personnel

During the 2018-19 influenza season, 31 (93.9%) of 33 hospitals had a HCP vaccination policy in place and two (6.1%) did not have one in place and one was considering one. Among the 31 hospitals with a policy, 14 (42.4%) allowed only medical and religious exemptions; and one (3.0%) allowed medical and personal/philosophical exemptions. One (3.0%) hospital allowed only medical exemptions. The remaining 15 (45.5%) allowed an exemption for medical, religious, and personal/philosophical reasons. Thirty (96.8%) hospitals with a policy required unvaccinated HCP with an approved exemption to wear a mask, and 19 (57.6%) compelled unvaccinated HCP without an acceptable reason for exemption to progressive discipline, potentially including termination. Hospitals with vaccination policies had significantly higher percentages of influenza vaccination as a whole (94.3%) than hospitals without mandatory policies (77.1%). Hospitals that utilized progressive discipline potentially including termination as a consequence for unvaccinated HCP without an acceptable exemption had a significantly higher vaccination percentage (96.3%) than hospitals that did not include potential termination as a consequence (88.2%).

Figure 23. Influenza vaccination percentages for hospitals with and without vaccination policies, 2018-19 influenza season




Table 25. Influenza vaccination policies and consequences for healthcare personnel by hospital, 2018-19 influenza season

Hospital	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP with Accepted Exemption	Consequences for Unvaccinated HCP without Accepted Exemption
Alice Peck Day Memorial Hospital	Medical, Religious	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination
Androscoggin Valley Hospital	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Receive verbal and/or written education
Catholic Medical Center	Medical, Religious	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education
Cheshire Medical Center	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination
Concord Hospital	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask, Progressive discipline, potentially including termination
Cottage Hospital	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask
Dartmouth Hitchcock Medical Center	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination
Elliot Hospital	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination
Encompass Health Hospital of Concord NH	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask
Exeter Hospital	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask
Franklin Regional Hospital	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Receive verbal and/or written education
Frisbie Memorial Hospital	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education
Huggins Hospital	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask
Lakes Region General Hospital	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Receive verbal and/or written education
Littleton Regional Healthcare	Medical, Religious	Wear a mask	Wear a mask, Progressive discipline, potentially including termination

*Exemptions include Medical, Religious, Personal/philosophical, and Other.

Table 25. (Continued) Influenza vaccination policies and consequences for healthcare personnel by hospital, 2018-19 influenza season

Hospital	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP with Accepted Exemption	Consequences for Unvaccinated HCP without Accepted Exemption
Monadnock Community Hospital	Medical, Religious	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education
NE Rehab, Pease	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination
NE Rehab, Salem	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination
NE Rehab, SNHMC	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination
NE Rehab, The Elliot	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination
New London Hospital Association	Medical, Religious	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination
Parkland Medical Center	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask
Portsmouth Regional Hospital	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask
Southern New Hampshire Medical Center	Medical, Religious	Other	Progressive discipline, potentially including termination
Speare Memorial Hospital	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Progressive discipline, potentially including termination
St Joseph Hospital	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination
The Memorial Hospital	Medical, Personal/philosophical	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education
Upper Connecticut Valley Hospital	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Receive verbal and/or written education
Valley Regional Hospital	Medical, Religious	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination
Weeks Medical Center	Medical	Wear a mask	Wear a mask
Wentworth-Douglass Hospital	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask

*Exemptions include Medical, Religious, Personal/philosophical, and Other.

Note: One hospital (3%) did not have mandatory vaccination policy during the 2018-19 influenza season, but was considering one at the time of the survey. One hospital (3%) did not have mandatory vaccination policies during the 2018-19 influenza season and were not considering one at the time of the survey. Response of "Other" under requirements for unvaccinated HCP with accepted exemptions indicated "nothing at this time".

IV. CONCLUSIONS

This tenth report on hospital HAI surveillance data displays continued progress toward the goal of eliminating HAI in NH. This report provides a picture of selected HAI data that can be used by healthcare facilities to identify areas for improvement and prevention, as well as healthcare consumers to make informed healthcare decisions.

Key findings described in this report include the following:

- All 33 individually licensed hospitals in NH complied with the HAI mandatory reporting law in 2018.
- NH hospitals reported less HAI associated with coronary artery bypass graft and central line associated blood stream infections based on national data; this difference was not statistically significant. Hospitals reported more HAI associated with catheter, colon procedures, knee replacement procedures, and abdominal hysterectomy procedures than predicted based on national data; this difference was not statistically significant.
- The majority of hospitals have similar number of infections than predicted based on national data.
- A few NH hospitals have more infections following certain procedures, which may warrant changes to current infection prevention practices in order to reduce infections.
- Statewide adherence to all four infection prevention practices during central line insertions was 98.3%, which is similar to 2017 (98.5%). Hospitals have made improvements since HAI reporting began, and should continue to work toward the goal of 100% adherence.
- Surgical antimicrobial prophylaxis data for 2014-2018 was not available and consequently not included in this report. However, in 2013, NH hospitals performed surgical antimicrobial prophylaxis correctly more often or similar to the national average.
- Vaccination coverage by hospitals during the 2018-19 influenza season ranged from 65.9% to 99.6%. The overall State percentage was 94.0%, which represents a slight increase from the 2017-18 influenza season when the statewide vaccination percentage was 93.3%. This was not statistically significant and that is to be expected as vaccination coverage approaches 100%.
- Thirty-one (93.9%) NH hospitals had an HCP vaccination policy in place during the 2018-19 season. This was similar to the 2017-18 season. Overall, hospitals with vaccination policies had significantly higher percentages of influenza vaccination as a whole (94.3%) than hospitals without mandatory policies (77.1%).

While this report only includes information on a subset of HAI, the information provided can be used as an important indicator of healthcare quality and infection prevention efforts in NH hospitals. Although data in this report have not been independently validated to assess reporting accuracy, this process is ongoing; a validation study is underway and will be the subject of a future report. Healthcare consumers can discuss the information provided in this

report with their healthcare provider and should review Appendix 4 for information on what individual patients can do to prevent HAI.

V. ACUTE CARE HOSPITAL REPORTS

Because data must be broken down into categories for risk adjustment and because rates must be suppressed if data are too sparse, data that can be presented for NH facilities may be limited. Due to restrictions on presenting data, there are several hospitals for which facility-specific infections data for specific measures cannot be presented. See technical notes for additional information on data restriction and presentation.




ALICE PECK DAY MEMORIAL

Lebanon, NH
 Not-for-profit, Critical Access
 # of Admissions: 1,315
 # of Beds: 32
 # of ICU Beds: 0
 # of Patient-days: 2,308

2018 HAI DATA REPORT

Standardized Infection Ratios



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	2	1.15	1.74	0.29 , 5.75	Similar
CLABSI	No ICU to monitor infections				
CAUTI	No ICU to monitor infections				
SSI	2	1.15	1.74	0.29 , 5.74	Similar
CABG	Facility does not perform this procedure				
COLO	Facility does not perform this procedure				
HYST	†	†	†	†	†
KPRO	2	1.06	1.88	0.32 , 6.22	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				

BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams


HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

ALICE PECK DAY MEMORIAL 2018 DATA REPORT

PROCESS MEASURES


Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP		98.3	
SCIP-1	Note: 2018 SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	94.3	94.0	Similar

Central Line Insertion Practices




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



ANDROSCOGGIN VALLEY

Berlin, NH

Not-for-profit, Critical Access

of Admissions: 1,590


of Beds: 25

of ICU Beds: 5

of Patient-days: 5,117

2018 HAI DATA REPORT

Standardized Infection Ratios



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	†	†	†	†	†
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical ICU (CAH)	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical ICU (CAH)	0	163	0.0	2.0	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

ANDROSCOGGIN VALLEY 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: 2018 SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	95.1	94.0	Similar




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Receive verbal and/or written education



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



CATHOLIC MEDICAL CENTER

Manchester, NH
 Not-for-profit, Acute Care
 # of Admissions: 26,694
 # of Beds: 264
 # of ICU Beds: 38

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	17	21.78	0.78	0.47 , 1.22	Similar
CLABSI	1	3.35	0.30	0.02 , 1.47	Similar
CAUTI	3	4.26	0.70	0.18 , 1.92	Similar
SSI	13	14.17	0.92	0.51 , 1.53	Similar
CABG	4	6.79	0.59	0.19 , 1.42	Similar
COLO	7	6.22	1.13	0.49 , 2.22	Similar
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	1	2,969	0.3	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	3	3,271	0.9	1.7	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

CATHOLIC MEDICAL CENTER 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	100.0	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	98.8	94.0	Higher




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



CHESHIRE MEDICAL CENTER

Keene, NH
 Not-for-profit, Acute Care
 # of Admissions: 6,292
 # of Beds: 100
 # of ICU Beds: 10
 # of Patient-days: 20,894

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	8	4.62	1.73	0.80 , 3.29	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	7	3.52	1.99	0.87 , 3.93	Similar
CABG	Facility does not perform this procedure				
COLO	5	2.09	2.39	0.87 , 5.29	Similar
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical ICU	1	488	2.0	1.1	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical ICU	0	989	0.0	2.0	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

CHESHIRE MEDICAL CENTER 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	100.0	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	91.7	94.0	Lower



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



CONCORD HOSPITAL
 Concord, NH
 Not-for-profit, Acute Care
 # of Admissions: 21,353
 # of Beds: 238
 # of ICU Beds: 18
 # of Patient-days: 71,302

2018 HAI DATA REPORT

Standardized Infection Ratios



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	11	14.58	0.75	0.40 , 1.31	Similar
CLABSI	1	1.99	0.50	0.03 , 2.48	Similar
CAUTI	4	3.29	1.22	0.39 , 2.93	Similar
SSI	6	9.30	0.65	0.26 , 1.34	Similar
CABG	†	†	†	†	†
COLO	1	4.50	0.22	0.01 , 1.09	Similar
HYST	†	†	†	†	†
KPRO	5	3.45	1.45	0.53 , 3.22	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	1	1,978	0.5	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	4	3,223	1.2	1.7	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

CONCORD HOSPITAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	100.0	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	94.2	94.0	Similar



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
NONE			



SCIP: Surgical care improvement project

CLIP: Central line insertion practices


DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



COTTAGE HOSPITAL
 Woodsville, NH
 Not-for-profit, Critical Access
 # of Admissions: 1,145
 # of Beds: 25
 # of ICU Beds: 3
 # of Patient-days: 4,289

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	†	†	†	†	†
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	Facility did not perform this procedure in 2017				
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	81	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

COTTAGE HOSPITAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	68.7	94.0	Lower



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical, Other	Wear a mask	Wear a mask



SCIP: Surgical care improvement project

CLIP: Central line insertion practices


DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



DHMC
 Lebanon, NH
 Not-for-profit, Acute Care
 # of Admissions: 26,751
 # of Beds: 444
 # of ICU Beds: 89
 # of Patient-days: 124,933

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	92	68.04	1.35	1.10 , 1.65	Higher
CLABSI	11	11.23	0.98	0.52, 1.70	Similar
CAUTI	24	16.68	1.44	0.94 , 2.11	Similar
SSI	57	40.13	1.42	1.08 , 1.82	Higher
CABG	2	3.15	0.64	0.11 , 2.10	Similar
COLO	36	25.67	1.40	0.99 , 1.92	Similar
HYST	11	6.28	1.75	0.92 , 3.04	Similar
KPRO	8	5.03	1.59	0.74 , 3.02	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Cardiac ICU	4	2,138	1.9	1.0	Similar
Medical ICU 1	2	2,479	0.8	1.2	Similar
Medical ICU 2	1	689	1.5	1.2	Similar
Med/Surg PED ICU	1	426	2.3	1.2	Similar
Surg ICU	3	2,719	1.1	1.1	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL ICU

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	0	206	0.0	2.1	Similar
BW Category B	0	275	0.0	1.3	Similar
BW Category C	0	306	0.0	0.8	Similar
BW Category D	†	†	†	†	†
BW Category E	0	219	0.0	0.7	Similar


CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Cardiac ICU	5	2,193	2.3	2.3	Similar
Medical ICU 1	3	3,046	1.0	3.5	Lower
Medical ICU 2	2	1,087	1.8	3.5	Similar
Med/Surg PED ICU	0	188	0.0	2.5	Similar
Surg ICU	14	4,040	3.5	3.4	Similar

DHMC 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	97.6	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	95.6	94.0	Higher



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:

BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams


HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



ELLIOT HOSPITAL
 Manchester, NH
 Not-for-profit, Acute Care
 # of Admissions: 6,163
 # of Beds: 270
 # of ICU Beds: 42
 # of Patient-days: 65,281

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	8	15.78	0.51	0.24 , 0.96	Lower
CLABSI	0	2.93	0.00	- , 1.02	Similar
CAUTI	1	1.63	0.61	0.03 , 3.03	Similar
SSI	7	11.22	0.62	0.27 , 1.23	Similar
CABG	Facility does not perform this procedure				
COLO	2	8.19	0.24	0.04 , 0.80	Lower
HYST	0	1.56	0.00	- , 1.92	Similar
KPRO	5	1.47	3.40	1.25 , 7.53	Higher

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	1,381	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	0	79	0.0	2.1	Similar
BW Category B	0	288	0.0	1.3	Similar
BW Category C	0	315	0.0	0.8	Similar
BW Category D	0	569	0.0	0.6	Similar
BW Category E	0	345	0.0	0.7	Similar

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	1	1,818	0.6	1.7	Similar

BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams


HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

ELLIOT HOSPITAL 2018 DATA REPORT

PROCESS MEASURES



Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	98.7	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	99.1	94.0	Higher

Central Line Insertion Practices



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination



SCIP: Surgical care improvement project

CLIP: Central line insertion practices


DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



EXETER HOSPITAL
 Exeter, NH
 Not-for-profit, Acute Care
 # of Admissions: 5,393
 # of Beds: 100
 # of ICU Beds: 10
 # of Patient-days: 21,414

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	7	5.85	1.20	0.52 , 2.37	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	5	4.62	1.08	0.39 , 2.39	Similar
CABG	Facility does not perform this procedure				
COLO	4	3.50	1.14	0.36 , 2.75	Similar
HYST	†	†	†	†	†
KPRO	1	1.02	0.98	0.05 , 4.83	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	1	804	1.2	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	1	857	1.2	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

EXETER HOSPITAL 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	96.6	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	92.7	94.0	Lower




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask



SCIP: Surgical care improvement project



CLIP: Central line insertion practices


DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



FRANKLIN REGIONAL
 Franklin, NH
 Not-for-profit, Critical Access
 # of Admissions: 926
 # of Beds: 25
 # of ICU Beds: 0
 # of Patient-days: 5,954

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	Facility did not report any data contributing to a SIR during this time period. <ul style="list-style-type: none"> No ICU to monitor central lines and urinary catheters No procedures of these types performed in 2017 				
CLABSI					
CAUTI					
SSI					
CABG					
COLO					
HYST					
KPRO					

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

FRANKLIN REGIONAL 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP		98.3	
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	92.8	94.0	Similar




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Receive verbal and/or written education



SCIP: Surgical care improvement project



CLIP: Central line insertion practices


DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



FRISBIE MEMORIAL HOSPITAL
 Rochester, NH
 Not-for-profit, Acute Care
 # of Admissions: 4,461
 # of Beds: 88
 # of ICU Beds: 6

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	5	4.30	1.16	0.43 , 2.58	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	3	3.29	0.91	0.23 , 2.47	Similar
CABG	Facility does not perform this procedure				
COLO	1	2.21	0.45	0.02 , 2.22	Similar
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	539	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	2	884	2.3	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

FRISBIE MEMORIAL HOSPITAL 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	95.2	94.0	Similar




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education



SCIP: Surgical care improvement project



CLIP: Central line insertion practices


DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



HUGGINS HOSPITAL
 Wolfeboro, NH
 Not-for-profit, Critical Access
 # of Admissions: 698
 # of Beds: 25
 # of ICU Beds: 4
 # of Patient-days: 4,402

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	1	1.01	0.99	0.05 , 4.88	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	52	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	176	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

HUGGINS HOSPITAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	93.0	94.0	Similar



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



LAKES REGION GENERAL

Laconia, NH
 Not-for-profit, Acute Care
 # of Admissions: 5,228
 # of Beds: 88
 # of ICU Beds: 10
 # of Patient-days: 23,087

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	2	5.39	0.37	0.06 , 1.23	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	2	4.34	0.46	0.07 , 1.52	Similar
CABG	Facility does not perform this procedure				
COLO	2	2.60	0.77	0.12 , 2.54	Similar
HYST	†	†	†	†	†
KPRO	0	1.44	0.00	- , 2.09	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	338	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	1,120	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

LAKES REGION GENERAL 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	95.8	94.0	Higher




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Receive verbal and/or written education



SCIP: Surgical care improvement project



CLIP: Central line insertion practices


DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



LITTLETON REGIONAL
 Littleton, NH
 Not-for-profit, Critical Access
 # of Admissions: 1,955
 # of Beds: 29
 # of ICU Beds: 4
 # of Patient-days: 5,917

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	3	2.10	1.43	0.36 , 3.89	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	3	1.85	1.62	0.41 , 4.42	Similar
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	112	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	348	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

LITTLETON REGIONAL 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	98.7	94.0	Higher




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



MONADNOCK COMMUNITY

Peterborough, NH
 Not-for-profit, Critical Access
 # of Admissions: 1,442
 # of Beds: 25
 # of ICU Beds: 0
 # of Patient-days: 4,451

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	†	†	†	†	†
CLABSI	No ICU to monitor central line infections				
CAUTI	No ICU to monitor catheter infections				
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	Facility did not perform central line insertions				

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	Facility did not perform catheter insertions in 2017				


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

MONADNOCK COMMUNITY 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP		98.3	
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	92.9	94.0	Similar




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.




NEW LONDON HOSPITAL

New London, NH
 Not For-profit, Critical Access
 # of Admissions: 1,335
 # of Beds: 25
 # of ICU Beds: 0
 # of Patient-days: 6,021

2018 HAI DATA REPORT

STANDARDIZED INFECTION RATIOS (SIR)



Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	†	†	†	†	†
CLABSI	No ICU to monitor central line infections				
CAUTI	No ICU to monitor catheter infections				
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

NEW LONDON HOSPITAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP		98.3	
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	97.3	94.0	Higher



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



PARKLAND MEDICAL CENTER

Derry, NH

For-profit, Acute Care


of Admissions: 4,401

of Beds: 86

of ICU Beds: 8

of Patient-days: 18,998

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	7	4.19	1.67	0.73 , 3.31	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	5	2.73	1.83	0.67 , 4.06	Similar
CABG	Facility does not perform this procedure				
COLO	3	1.34	2.23	0.56 , 6.07	Similar
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical ICU	1	1,130	0.9	1.1	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical ICU	1	1,257	0.8	2.0	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

PARKLAND MEDICAL CENTER 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	96.2	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	92.3	94.0	Lower




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.




PORTSMOUTH REGIONAL

Portsmouth, NH
 For-profit, Acute Care
 # of Admissions: 10,056
 # of Beds: 224
 # of ICU Beds: 14
 # of Patient-days: 52,175

2018 HAI DATA REPORT

STANDARDIZED INFECTION RATIOS (SIR)



Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	17	13.02	1.31	0.76 , 2.09	Similar
CLABSI	2	2.74	0.73	0.12 , 2.41	Similar
CAUTI	6	3.23	1.86	0.75 , 3.86	Similar
SSI	9	7.05	1.28	0.62 , 2.34	Similar
CABG	5	2.19	2.29	0.84 , 5.07	Similar
COLO	4	3.15	1.27	0.40 , 3.06	Similar
HYST	†	†	†	†	†
KPRO	0	1.56	0.00	- , 1.93	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	2	3,157	0.6	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	6	3,608	1.7	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

PORTSMOUTH REGIONAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	100	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	90.2	94.0	Lower



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014–2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



SOUTHERN NH MEDICAL

Nashua, NH
 Not-for-profit, Acute Care
 # of Admissions: 14,047
 # of Beds: 162
 # of ICU Beds: 17
 # of Patient-days: 38,672

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	4	7.93	0.50	0.16 , 1.22	Similar
CLABSI	†	†	†	†	†
CAUTI	1	1.01	0.99	0.05 , 4.90	Similar
SSI	3	6.49	0.46	0.11 , 1.25	Similar
CABG	Facility does not perform this procedure				
COLO	3	4.02	0.75	0.19 , 2.02	Similar
HYST	0	1.11	0.00	- , 2.71	Similar
KPRO	0	1.36	0.00	- , 2.20	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	521	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	†	†	†	†	†
BW Category B	†	†	†	†	†
BW Category C	†	†	†	†	†
BW Category D	†	†	†	†	†
BW Category E	†	†	†	†	†

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	1	1,378	0.7	1.7	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

SOUTHERN NH MEDICAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	100.0	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	97.5	94.0	Higher



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Other	Progressive discipline, potentially including termination



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.
- Response of “Other” under requirements for unvaccinated HCP with accepted exemptions indicated “no mask required; herd immunity”



SPEARE MEMORIAL HOSPITAL

Plymouth, NH
 Not-for-profit, Critical Access
 # of Admissions: 2,319
 # of Beds: 25
 # of ICU Beds: 4
 # of Patient-days: 6,277

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	2	1.16	1.72	0.29 , 5.70	Similar
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	Facility does not perform this procedure				
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	67	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	213	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

SPEARE MEMORIAL HOSPITAL 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	90.6	94.0	Lower




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/Philosophical	Wear a mask, Receive verbal and/or written education	Wear a mask, Progressive discipline, potentially including termination



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014–2017 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.




ST JOSEPH HOSPITAL

Nashua, NH
 Not-for-profit, Acute Care
 # of Admissions: 4,840
 # of Beds: 208
 # of ICU Beds: 11
 # of Patient-days: 26,866

2018 HAI DATA REPORT

STANDARDIZED INFECTION RATIOS (SIR)



Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	0	4.67	0.00	-, 0.78	Lower
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	0	3.39	0.00	-, 0.88	Lower
CABG	Facility does not perform this procedure				
COLO	0	2.31	0.00	-, 1.29	Similar
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	585	0.0	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	1,156	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

ST JOSEPH HOSPITAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	100.0	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	97.4	94.0	Higher



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/Philosophical	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



THE MEMORIAL HOSPITAL

North Conway, NH
 Not-for-profit, Critical Access
 # of Admissions: 1,626
 # of Beds: 25
 # of ICU Beds: 0
 # of Patient-days: 5,113

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	†	†	†	†	†
CLABSI	No ICU to monitor central line infections				
CAUTI	No ICU to monitor catheter infections				
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	†	†	†	†	†
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

THE MEMORIAL HOSPITAL 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP		98.3	
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	85.8	94.0	Lower



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Personal/philosophical	Wear a mask	Wear a mask, Progressive discipline, potentially including termination, Receive verbal and/or written education



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



UPPER CONNECTICUT VALLEY

Colebrook, NH
 Not-for-profit, Critical Access
 # of Admissions: 231
 # of Beds: 12
 # of ICU Beds: 0
 # of Patient-days: 875

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	Facility did not report any data contributing to a SIR during this time period. <ul style="list-style-type: none"> No ICU to monitor central lines or urinary catheters No procedures of these types performed in 2018 				
CLABSI					
CAUTI					
SSI					
CABG					
COLO					
HYST					
KPRO					

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

UPPER CONNECTICUT VALLEY 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP		98.3	
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	91.1	94.0	Similar



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
NO			



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



VALLEY REGIONAL HOSPITAL

Claremont, NH
 Not-for-profit, Critical Access
 # of Admissions: 644
 # of Beds: 21
 # of ICU Beds: 0
 # of Patient-days: 3,589

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	†	†	†	†	†
CLABSI	No ICU to monitor central line infections				
CAUTI	No ICU to monitor catheter infections				
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	Facility does not perform this procedure				
HYST	Facility did not perform this procedure				
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
No ICU	No ICU to monitor infections				


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

VALLEY REGIONAL HOSPITAL 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP		98.3	
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	99.6	94.0	Higher




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious,	Wear a mask, Receive verbal and/or written education	Progressive discipline, potentially including termination



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.




WEEKS MEDICAL CENTER

Lancaster, NH
 Not-for-profit, Critical Access
 # of Admissions: 1,347
 # of Beds: 25
 # of ICU Beds: 3
 # of Patient-days: 3,786

2018 HAI DATA REPORT

Standardized Infection Ratios



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	†	†	†	†	†
CLABSI	†	†	†	†	†
CAUTI	†	†	†	†	†
SSI	†	†	†	†	†
CABG	Facility does not perform this procedure				
COLO	†	†	†	†	†
HYST	Facility did not perform this procedure				
KPRO	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	†	†	†	†	†

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	75	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

WEEKS MEDICAL CENTER 2018 DATA REPORT


PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	†	98.3	†
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	97.8	94.0	Higher




INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical	Wear a mask	Wear a mask



SCIP: Surgical care improvement project



CLIP: Central line insertion practices

DATA NOTES:


- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.



WENTWORTH-DOUGLASS

Dover, NH
 Not-for-profit, Acute Care
 # of Admissions: 8,234
 # of Beds: 189
 # of ICU Beds: 11
 # of Patient-days: 39,617

2018 HAI DATA REPORT



STANDARDIZED INFECTION RATIOS (SIR)

Measure	Observed Infections	Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval	Comparison to Predicted
Overall HAI	6	9.00	0.67	0.27 , 1.39	Similar
CLABSI	2	1.10	1.82	0.31 , 6.02	Similar
CAUTI	0	1.12	0.00	- , 2.69	Similar
SSI	4	6.78	0.59	0.18 , 1.42	Similar
CABG	Facility does not perform this procedure				
COLO	1	4.20	0.24	0.01 , 1.17	Similar
HYST	1	1.04	0.96	0.05 , 4.73	Similar
KPRO	2	1.54	1.30	0.22 , 4.29	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES

Type of Unit	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	2	1,457	1.4	0.8	Similar

CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION RATES BY BIRTHWEIGHT IN NEONATAL INTENSIVE CARE UNITS

Birthweight Category	Number of Infections	Number of Central Line Days	Rate per 1,000 Central Line Days	National Rate	Comparison to National Rate
BW Category A	No Neonatal ICU to monitor infections				
BW Category B					
BW Category C					
BW Category D					
BW Category E					

CATHETER-ASSOCIATED URINARY TRACT INFECTION RATES

Type of Unit	Number of Infections	Number of Catheter Days	Rate per 1,000 Catheter Days	National Rate	Comparison to National Rate
Medical/Surgical ICU	0	1,526	0.0	1.3	Similar


BW Category A: Equal or less than 750 grams BW Category B: Equal and between 751 and 1,000 grams BW Category C: Equal and between 1,001 and 1,500 grams BW Category D: Equal and between 1,501 and 2,500 grams BW Category E: More than 2,500 grams

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty

WENTWORTH-DOUGLASS 2018 DATA REPORT



PROCESS MEASURES

Measure	Percent Adherence	State Coverage	Comparison to State Coverage
CLIP	98.6	98.3	Similar
SCIP-1	Note: SCIP data were not available at time of publication and as a result are excluded from this report.		
SCIP-2			
SCIP-3			
Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	83.2	94.0	Lower



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious, Personal/philosophical	Wear a mask	Wear a mask



SCIP: Surgical care improvement project

CLIP: Central line insertion practices

DATA NOTES:

- The 2018 central line-associated blood stream infections (CLABSI), surgical site infections (SSI) and catheter-associated urinary tract infections (CAUTI) data presented in this report have not been validated and must be interpreted with the understanding that in general there are both under- and over-reporting of infections.
- In New Hampshire in 2018, CLABSI were monitored in all intensive care units (including pediatric and neonatal units) and not in other inpatient locations. CAUTI were monitored in all intensive care units (including pediatric units and excluding neonatal ICU) and not in other inpatient locations.
- Hospitals do not use a standard method of post-discharge surveillance to identify infections once a patient has been discharged. This poses a challenge for data interpretation, because a higher SSI rate (for example) at a given hospital may represent either poor infection prevention practices or, conversely, a better system for identifying infections.
- SSI reporting requires not only reporting of infections but also detailed information on every patient who underwent the procedure being monitored. As such, DHHS has elected to monitor a subset of procedures based on national recommendations since it would not be feasible for hospitals to report information on every patient receiving a surgical procedure due to the burden of reporting.
- As of July 1, 2015, SCIP 2014– 2018 data reported to CMS was unavailable and the HAI Program was unable to conduct further analysis of this measure as routinely included in this report.




HAMPSTEAD HOSPITAL
 Hampstead, NH
 Private
 # of Admissions: 1,924 (2016)
 # of Beds: 111

2018 HAI DATA REPORT

PROCESS MEASURES

Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	65.9	94.0	Lower



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
NONE			

DATA NOTES:

- Specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CLABSI, CAUTI, or CLIP, because they do not have ICU, nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries.
- New Hampshire’s five rehabilitation and two psychiatric hospitals are only required to report influenza vaccination rates for patients and staff.

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection
 SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty
 SCIP: Surgical care improvement project CLIP: Central line insertion practices




ENCOMPASS REHABILITATION HOSPITAL

Concord, NH
 Corporate
 # of Admissions: 905
 # of Beds: 50

2018 HAI DATA REPORT

PROCESS MEASURES

Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	92.6	94.0	Similar



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
NONE			

DATA NOTES:

- Specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CLABSI, CAUTI, or CLIP, because they do not have ICU, nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries.
- New Hampshire's five rehabilitation and two psychiatric hospitals are only required to report influenza vaccination rates for patients and staff.

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection
 SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty
 SCIP: Surgical care improvement project CLIP: Central line insertion practices




NEW HAMPSHIRE HOSPITAL

Concord, NH
 State-operated
 # of Admissions: 1,473
 # of Beds: 168

2018 HAI DATA REPORT

PROCESS MEASURES

Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	78.9	94.0	Lower



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
NONE			

DATA NOTES:

- Specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CLABSI, CAUTI, or CLIP, because they do not have ICU, nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries.
- New Hampshire’s five rehabilitation and two psychiatric hospitals are only required to report influenza vaccination rates for patients and staff.

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection
 SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty
 SCIP: Surgical care improvement project CLIP: Central line insertion practices



NORTHEAST REHABILITATION HOSPITAL, THE ELLIOT

Manchester, NH

Network


of Admissions: 307

of Beds: 15

2018 HAI DATA REPORT

PROCESS MEASURES

Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	98.5	94.0	Similar



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination

DATA NOTES:

- Specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CLABSI, CAUTI, or CLIP, because they do not have ICUs nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries.
- New Hampshire’s five rehabilitation and two psychiatric hospitals are only required to report influenza vaccination rates for patients and staff.

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection
 SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty
 SCIP: Surgical care improvement project CLIP: Central line insertion practices



NORTHEAST REHABILITATION HOSPITAL, PEASE

Portsmouth, NH

Network


of Admissions: 832

of Beds: 33

2018 HAI DATA REPORT

PROCESS MEASURES

Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	99.6	94.0	Higher



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination

DATA NOTES:

- Specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CLABSI, CAUTI, or CLIP, because they do not have ICU, nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries.
- New Hampshire's five rehabilitation and two psychiatric hospitals are only required to report influenza vaccination rates for patients and staff.

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection
 SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty
 SCIP: Surgical care improvement project CLIP: Central line insertion practices



NORTHEAST REHABILITATION HOSPITAL, SALEM

Salem, NH

Network


of Admissions: 1,523

of Beds: 67

2018 HAI DATA REPORT

PROCESS MEASURES

Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	99.4	94.0	Higher



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination

DATA NOTES:

- Specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CLABSI, CAUTI, or CLIP, because they do not have ICU, nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries.
- New Hampshire's five rehabilitation and two psychiatric hospitals are only required to report influenza vaccination rates for patients and staff.

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection
 SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty
 SCIP: Surgical care improvement project CLIP: Central line insertion practices



NORTHEAST REHABILITATION HOSPITAL, SNHMC

Nashua, NH

Network


of Admissions: 394

of Beds: 20

2018 HAI DATA REPORT

PROCESS MEASURES

Measure	Percent Vaccinated	State Coverage	Comparison to State Coverage
HCP Influenza Vaccination	99.0	94.0	Higher



INFLUENZA VACCINATION POLICIES, 2018-2019 INFLUENZA SEASON

Policy	Exemptions Allowed in Policy*	Requirements for Unvaccinated HCP With Accepted Exemption	Consequences for Unvaccinated HCP Without Accepted Exemption
YES	Medical, Religious	Wear a mask	Progressive discipline, potentially including termination

DATA NOTES:

- Specialty hospitals (rehabilitation and psychiatric hospitals) are not required to report CLABSI, CAUTI, or CLIP, because they do not have ICU, nor SSI and surgical antimicrobial prophylaxis administration data, because they do not perform surgeries.
- New Hampshire’s five rehabilitation and two psychiatric hospitals are only required to report influenza vaccination rates for patients and staff.

HAI: Healthcare-associated infection CLABSI: Central line-associated bloodstream infection CAUTI: Catheter-associated urinary tract infection
 SSI: Surgical site infection COLO: Colon procedure HYST: Abdominal hysterectomy CABG: Coronary artery bypass graft KPRO: Knee arthroplasty
 SCIP: Surgical care improvement project CLIP: Central line insertion practices

APPENDIX 1: Technical Notes

1. The majority of data in this report were extracted from NHSN on 7/8/2019; additional influenza vaccination data were extracted from other data sources on the same date. Changes or new infections reported by hospitals after this date may not be reflected in this report.
2. Rate data were appropriately risk-adjusted according to standard NHSN recommendations. Rates were only presented if appropriately risk-adjusted as follows:
 - a. CLABSI: rate data must be broken down and aggregated only by the same type of unit.
 - b. CAUTI: rate data must be broken down and aggregated only by the same type of unit
 - c. CLIP: currently there are no CDC recommendations for risk-adjusting CLIP data.
 - d. SSI: In accordance with CDC recommendations and changes to NHSN methodology beginning in 2010, rates are no longer presented.
3. Rates for any grouping were not presented if data were insufficient to generate a stable rate.
 - a. CLABSI: there must be at least 50 central line days in the denominator to present a rate.
 - b. CAUTI: there must be at least 50 catheter days in the denominator to present a rate.
 - c. CLIP: there must be at least 20 insertions in the denominator to present a rate.
 - d. SSI: in accordance with CDC recommendations and changes to NHSN methodology beginning in 2010, rates are no longer presented.
4. SIR for any grouping were not presented if less than one infection was predicted.
5. All confidence intervals presented in this report are 95% confidence intervals. A confidence interval is a measure of certainty (usually with 95% confidence) of an estimate (such as a percentage). Because we can never obtain a hospital's true "population" data (e.g., all patients for all time), we use statistical procedures to "estimate" various measurements using "sample" data. Since estimates have "variability" we use 95% confidence limits to describe the variability around the estimate. The confidence interval gives us the range within which the TRUE value will fall 95% of the time, assuming that the sample data are reflective of the true population. If the confidence intervals for the two rates overlap, then it is reasonably possible that the REAL rates are not different from one another.
6. Statistical significance is affected by sample size. If a value is almost or just barely significant, just a few additional observations can push significance one way or the other.

Standardized Infection Ratios

7. Calculating a SIR: The SIR is the number of observed infections divided by the number of predicted infections based on most recent national data. In order to calculate an SIR, it is recommended that there be at least one predicted infection. See Appendix 3 for more information on the SIR.
8. Interpreting a SIR: The resulting SIR is a comparison between the number of observed infections and the number predicted.
 - a. An SIR of 1.0 means that exactly the same number of infections was observed as was predicted.
 - b. An SIR of less than one means that fewer infections were observed than was predicted (for example, SIR = 0.70 would be interpreted as 30% fewer infections observed than predicted).
 - c. An SIR of more than one means that fewer infections were observed than were predicted (for example, SIR = 1.30 would be interpreted as 30% more infections observed than predicted).
9. Calculating a corresponding confidence interval for a SIR: All hospital-specific SIR and corresponding confidence intervals in this report were generated directly by NHSN using statistical methods similar to those described in Liddell FD. Simple exact analysis of the standardized mortality ratio. *Journal of Epidemiology and Community Health*, 1984; 38:85-88.^x
10. Interpreting a SIR confidence interval: A confidence interval is a measure of certainty (usually with 95% confidence) of an estimate (such as a SIR). Confidence intervals can be used to assess whether differences in the number of observed and predicted infections is statistically significant (different or similar).
 - a. For confidence intervals that contain the value 1.0, the observed number of infections will be considered "Similar" to the predicted number of infections based on national data (e.g., 0.27–1.49).
 - b. For confidence intervals that are lower than and do not contain the value 1.0, the observed number of infections will be considered "Lower" than the predicted number of infections based on national data (e.g., 0.13–0.74).
 - c. For confidence intervals that are higher than and do not contain the value 1.0, the observed number of infections will be considered "Higher" than the predicted number of infections based on national data (e.g., 1.09–2.63).

Infection Rates

11. Calculating a CLABSI rate: CLABSI rates are presented as the number of infections per 1,000 central line days.

$$\text{CLABSI rate} = (\text{number of infections} / \text{number of central line days}) \times 1,000$$

12. Calculating a CAUTI rate: CAUTI rates are presented as the number of infections per 1,000 catheter days.

$$\text{CAUTI rate} = (\text{number of infections} / \text{number of catheter days}) \times 1,000$$

13. Interpreting a p-value: All hospital-specific rates and corresponding p-values in this report were generated directly by NHSN using Poisson statistical methods. State-level rates and corresponding p-values were calculated by DHHS using exact methods. A p-value provides a statistical comparison of two values in order to determine whether those values are statistically different or similar. In this report, p-values are used to assess whether hospital infection rates are similar or different to national infection rates. A p-value of <0.05 would indicate the hospital rate is significantly different than the national rate.

- a. If the p-value is ≥ 0.05 , then the hospital rate would be considered statistically “Similar” to the national rate.
- b. If the hospital rate is lower than the national rate and the p-value is <0.05, then the hospital rate would be considered significantly “Lower” than the national rate.
- c. If the hospital rate is higher than the national rate and the p-value is <0.05, then the hospital rate would be considered significantly “Higher” than the national rate.

Process Measure Percentages

14. Calculating a CLIP adherence percentage: CLIP adherence percentages are presented as the number of insertions that met the adherence criteria divided by the total number of insertions expressed as a percent.

$$\text{CLIP adherence (\%)} = (\text{number of insertions that met adherence criteria} / \text{total number of insertions}) \times 100$$

15. Calculating an influenza vaccination percentage: Influenza vaccination percentages are presented as the number of HCP vaccinated divided by the total number of HCP expressed as a percent.

$$\text{Influenza vaccination (\%)} = (\text{number of HCP vaccinated} / \text{total number of HCP}) \times 100$$

16. Calculating a surgical antimicrobial prophylaxis adherence percentage: Surgical antimicrobial prophylaxis adherence percentages are presented as the number of orders for which administration adhered to the measure (SCIP-1, SCIP-2, or SCIP-3) divided by the total number of orders expressed as a percent.

$$\text{Surgical antimicrobial prophylaxis adherence (\%)} = (\text{number of orders administered on time} / \text{total number of orders}) \times 100$$

17. Calculating a corresponding confidence interval for a CLIP adherence percentage: Confidence intervals calculated for CLIP data presented in this report are mid-p exact 95% confidence intervals, which were calculated using R statistical software program.

18. Calculating a corresponding confidence interval for an influenza vaccination percentage: Confidence intervals calculated for influenza vaccination data presented in this report are mid-p exact 95% confidence intervals, which were calculated using a statistical software

program. In prior reports, confidence intervals for influenza vaccination data were Wald normal approximation 95% confidence intervals, however the method of calculating these confidence intervals were changed due to the addition of several hospitals with small numbers of HCP.

19. Calculating a corresponding confidence interval for a surgical antimicrobial prophylaxis adherence percentage: Confidence intervals calculated for SCIP data presented in this report are Wald normal approximation 95% confidence intervals for national and State data, and mid-p exact 95% confidence intervals for hospital data, which were calculated using R statistical software program.

20. Interpreting a proportion confidence interval for central line insertion and vaccination data: A confidence interval is a measure of certainty (usually with 95% confidence) of an estimate (such as a percentage). Confidence intervals can be used to assess whether differences in the percentages observed for each group (for example, hospital versus State) is statistically significant.
 - a. Confidence intervals that overlap the State confidence interval are considered "Similar" to the overall State percentage.
 - b. Confidence intervals that are lower than and do not overlap the State confidence interval are considered "Lower" than the overall State percentage.
 - c. Confidence intervals that are higher than and do not overlap the State confidence interval are considered "Higher" than the overall State percentage.

APPENDIX 2: Influenza Vaccination Survey Questions, 2018-2019 Season

1. Background information (facility and survey respondent)
2. How many patients were admitted to your hospital between October 1, 2018 and March 31, 2019? Include all patients that were admitted to your facility during this period, even if they were admitted or moved during the influenza season.
 - 2a. Total number of patient admissions
 - 2b. Total number of patient admissions excluding readmissions
3. How many of the patients admitted to your facility between October 1, 2018 and March 31, 2019 received a seasonal influenza vaccination (at your facility or elsewhere) for the 2018-19 season? Influenza vaccine for a given influenza season may be available as early as July or August. Include all immunized patients that received the 2017-18 vaccine product, even if administered prior to October 1, 2018.
 - 3a. Total number of patients immunized against influenza for the 2018-19 season
 - 3b. Total number of patients not immunized against influenza for the 2018-19 season
4. How many of the patients admitted to your facility between October 1, 2018 and March 31, 2019 had ever received a pneumococcal disease vaccination (at your facility or elsewhere)?
5. How many HCP worked or volunteered in your facility for at least one working day between October 1, 2018 and March 31, 2019?
6. How many HCP received a seasonal influenza vaccination (at your facility or elsewhere) for the 2018-19 season? Influenza vaccine for a given influenza season may be available as early as July or August. Include all immunized HCP that received the 2018-19 vaccine product, even if administered prior to October 1, 2018.
 - 6a. Total number of HCP immunized against influenza for the 2018-19 season
 - 6b. Total number of HCP not immunized against influenza for the 2018-19 season
7. Of the HCP not immunized against influenza for the 2018-19 influenza season, how many HCP did not receive the seasonal influenza vaccine for each of the following reasons: medical contraindication, religious, other (e.g., personal/philosophical), unknown?
8. Does your facility have a seasonal influenza vaccination policy? Such a policy means that the facility requires all or some portion of HCPs working at that facility to receive a seasonal influenza vaccine. If NO, skip to item 13.
 - 8a. Yes, there is a policy currently in place
 - 8b. No, but we are considering a policy
 - 8c. No, and we are not considering a policy
 - 8d. Other
9. If your facility has a seasonal influenza vaccination policy, what reasons for exemption are acceptable (medical, religious, personal/philosophical, other)? Check all that apply.

10. If your facility has a seasonal influenza vaccination policy, what do you require of unvaccinated HCP with an acceptable reason for exemption (wear a mask, receive verbal and/or written education, other)? Check all that apply.
11. If your facility has a seasonal influenza vaccination policy, what are the potential consequences for unvaccinated HCP without an acceptable reason for exemption (wear a mask, progressive discipline potentially including termination, receive verbal and/or written education, other)? Check all that apply.
12. If your facility has a seasonal influenza vaccination policy, how many people were terminated, suspended, resigned, or dismissed as a result of noncompliance with the policy during the 2018-19 influenza season (terminated, temporarily suspended, resigned, dismissed permanently)?
13. Does your facility offer the high-dose influenza vaccine?
14. Please enter any comments you would like to share.

APPENDIX 3: Understanding the Relationship between Healthcare-Associated Infection Rates and Standardized Infection Ratio Comparison Metrics

HAI Elimination Metrics are very useful for performing evaluations.^{xi} Several metrics are based on the science employed in NHSN. While national aggregate CLABSI data are published in the annual NHSN reports, these rates must be stratified by types of locations to be risk-adjusted. This scientifically sound risk-adjustment strategy creates a practical challenge to summarizing this information nationally, regionally, or even for an individual healthcare facility. For instance, when comparing CLABSI rates, there may be quite a number of different types of locations for which a CLABSI rate could be reported. This raises the need for a way to combine CLABSI rate data across locations.

A SIR can be used as an indirect standardization method for summarizing HAI experience across any number of stratified groups of data. To illustrate the method for using an SIR as an HAI comparison metric, the following example data are displayed below:

Risk Group Stratifier	Observed CLABSI Rates			NHSN CLABSI Rates for 2017 (Standard Population)			
	Location Type	#CLABSI	#Central line-days	CLABSI rate*	#CLABSI	#Central line-days	CLABSI rate*
	ICU	170	100,000	1.7	1200	600,000	2.0
	WARD	58	58,000	1.0	600	400,000	1.5
$\text{SIR} = \frac{\text{observed}}{\text{expected}} = \frac{170 + 58}{100000 \times \left(\frac{2}{1000}\right) + 58,000 \times \left(\frac{1.5}{1000}\right)} = \frac{228}{200 + 87} = \frac{228}{287} = 0.79 \quad 95\% \text{ CI} = (0.628, 0.989)$							

*Defined as the number of CLABSI per 1000 central line days

In the table above, there are two strata to illustrate risk-adjustment by location type for which national data exist from NHSN. The SIR calculation is based on dividing the total number of observed CLABSI events by a “predicted” number using the CLABSI rates from the standard population. This “predicted” number is calculated by multiplying the national CLABSI rate from the standard population by the observed number of central line days for each stratum, which can also be understood as a prediction or projection. If the observed data represented a follow-up period, such as 2017, one would state that an SIR of 0.79 indicates that there was a 21% reduction in CLABSI overall for the nation, region, or facility.

The SIR concept and calculation is completely based on the underlying CLABSI rate data that exist across a potentially large group of strata. Thus, the SIR provides a single metric for performing comparisons rather than attempting to perform multiple comparisons across many strata which makes the task cumbersome.

The SIR concept and calculation can be applied equitably to other HAI metrics. This is especially true for HAI metrics for which national data are available and reasonably precise using a measurement system such as the NHSN. The SIR calculation methods differ in the risk group stratification only.

The SSI SIR uses improved risk adjustment calculated through logistic modeling. This allows for all available risk factors to be procedure specific. See the following logistic equation and SSI predictive risk factors that are used for calculating SSI SIR, respectively.

$$\text{logit}(p) = \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 = -5.448 + 0.520 (\text{Age} \leq 44^*) + 0.425 (\text{ASA } 3/4/5^*) + 0.501 (\text{Duration} > 100^*) + 1.069 (\text{Med school affiliation}^*)$$

**For these risk factors, if present = 1; if not = 0*

Procedure Code	SSI Predictive Risk Factors From SSI Logistic Models
CABG	Age, ASA, Duration, Gender, Hospital Bed Size
COLO	Age, Anesthesia, ASA, Duration, Endoscope, Medical School Affiliation, Hospital Bed Size, Wound Class
HYST	Age, Anesthesia, ASA, Duration, Endoscope, Hospital Bed Size
KPRO	Age, Anesthesia, ASA, Duration, Gender, Revision, Hospital Bed Size, Trauma

Detailed descriptions of the SIR in NHSN are available at: http://www.cdc.gov/nhsn/PDFs/Newsletters/NHSN_NL_OCT_2010SE_final.pdf.

There are clear advantages to reporting and comparing a single number for prevention assessment. In addition to the simplicity of the SIR concept and the advantages listed above, it is important to note another benefit of using an SIR comparison metric for HAI data. If there was need at any level of aggregation (national, regional, facility-wide, etc.) to combine the SIR values across mutually exclusive data one could do so. The below table demonstrates how the example data from the previous two metric settings could be summarized.

HAI Metric	Observed HAI			Predicted HAI		
	#CLABSI	#SSI [†]	#Combined HAI	#CLABSI	#SSI [†]	#Combined HAI
CLABSI 1	228			287		
SSI 1		636			853.8	
Combined HAI			228 + 636 = 864			287 + 853.8 = 1140.8
$\text{SIR} = \frac{\text{observed}}{\text{expected}} = \frac{228 + 636}{287 + 853.8} = \frac{864}{1140.8} = 0.76 \quad 95\% \text{ CI} = (0.673, 0.849)$						

APPENDIX 4: Preventing Healthcare-Associated Infections

What You Can Do to Prevent Healthcare-Associated Infections

There are several prevention tips you can follow all the time to reduce your chance of getting an infection or spreading your infection to others.

1. Clean your hands.

- Use soap and warm water. Rub your hands for at least 20 seconds. Rub your palms, fingernails, in between your fingers, and the backs of your hands.
- If your hands do not look dirty, you can clean them with alcohol-based hand rub. Rub the gel all over your hands, especially under your nails and between your fingers, until your hands are dry.
- Clean your hands before touching or eating food. Clean them after you use the bathroom, take out the trash, change a diaper, visit someone who is ill, or play with a pet.

2. Make sure healthcare providers clean their hands first, even if they wear gloves, before touching you or performing a procedure.

- Doctors, nurses, dentists, and other healthcare providers come into contact with many bacteria and viruses. If you do not see your healthcare provider wash their hands or use an alcohol-based hand rub before they treat you, ask them if they have cleaned their hands.
- Healthcare providers should wear clean gloves when they perform tasks such as taking throat cultures, pulling teeth, taking blood, touching wounds or body fluids, while suctioning tubes, and examining your mouth or genitalia. Don't be afraid to ask if they should wear gloves.

3. Cover your mouth and nose.

- Many diseases are spread through sneezes and coughs. When you sneeze or cough, the germs can travel three feet or more. Cover your mouth and nose to prevent the spread of infection to others.
- Use a tissue. Keep tissues handy at home, at work, and in your pocket. Be sure to throw away used tissues and clean your hands after coughing or sneezing.
- If you don't have a tissue, cover your mouth and nose with the bend of your elbow or hands. If you use your hands, clean them right away.

4. If you are sick, avoid close contact with others.

- If you are sick, stay away from other people or stay home. Don't shake hands or touch others.
- When you go for medical treatment, call ahead and ask if there is anything you can do to avoid infecting people in the waiting room.

5. Get shots to avoid disease and fight the spread of infection.

- Make sure that your vaccinations are current—even for adults. Check with your doctor about shots you may need.

6. If you are prescribed an antibiotic for an illness, take them exactly as directed by your doctor.

- Don't take half-doses or stop before you complete your prescribed course even if you feel better. Not taking them as directed can lead to infections that become resistant to antibiotics, making them more difficult to treat.

What You Can Do to Help Prevent a Catheter-Associated Bloodstream Infection

- Ask your doctors and nurses to explain why you need the catheter and how long you will have it.
- Ask your doctors and nurses what infection prevention methods they will use during the catheter insertion.
- Make sure that all doctors and nurses caring for you clean their hands with soap and water or an alcohol-based hand rub before and after caring for you. If you do not see your providers clean their hands, please ask them to do so.
- If the bandage comes off or becomes wet or dirty, tell your nurse or doctor immediately.
- Inform your nurse or doctor if the area around your catheter is sore or red.
- Do not let family and friends who visit touch the catheter or the tubing.
- Make sure family and friends clean their hands with soap and water or an alcohol-based hand rub before and after visiting you.
- Some patients are sent home from the hospital with a catheter in order to continue their treatment. If you go home with a catheter, your doctors and nurses will explain everything you need to know about taking care of your catheter.
 - Make sure you understand how to care for the catheter before leaving the hospital. For example, ask for instructions on showering or bathing with the catheter and how to change the catheter dressing.
 - Make sure you know who to contact if you have questions after you get home.
 - Make sure you wash your hands with soap and water or an alcohol-based hand rub before handling your catheter.
 - Watch for the signs and symptoms of catheter-associated bloodstream infection, such as soreness or redness at the catheter site or fever, and call your healthcare provider immediately if any occur.

What Hospitals Do to Prevent Catheter-Associated Bloodstream Infections

To prevent catheter-associated bloodstream infections, doctors and nurses will:

- Choose a vein where the catheter can be safely inserted and where risk for infection is small.
- Clean hands with soap and water or alcohol-based hand rub before putting in the catheter.
- Wear a mask, cap, sterile gown, and sterile gloves when putting in the catheter to keep it sterile. The patient will be covered with a sterile sheet.
- Clean the patient's skin with an antiseptic cleanser before putting in the catheter.

- Clean hands, wear gloves, and clean the catheter opening with an antiseptic solution before using the catheter to draw blood or give medications. Healthcare providers also clean their hands and wear gloves when changing the bandage that covers the area where the catheter enters the skin.
- Decide every day if the patient still needs to have the catheter. The catheter will be removed as soon as it is no longer needed.

What You Can Do to Help Prevent Catheter-Associated Urinary Tract Infections

- Ask doctors to explain why you need the catheter and how long you will have it.
- Make sure that your doctors and nurses caring for you clean their hands and use sterile gloves for catheter insertion.
- Make sure the tubing or bag is not on the floor. If it drops or is on the floor, ask for new tubing or bag.
- Ask doctors and nurses what infection prevention methods they will use during the catheter insertion.
- Ask your doctors and nurses if you still need the catheter each day.
- Always clean your hands before and after doing catheter care.
- Always keep your urine bag below the level of your bladder.
- Do not tug or pull on the tubing.

What Hospitals Do to Prevent Catheter-Associated Urinary Tract Infections

To prevent catheter-associated urinary tract infections, doctors and nurses will:

- Put in catheters only when necessary and are removed as soon as possible.
- Clean hands with soap and water or alcohol-based hand rub and put on sterile gloves before putting in the catheter.
- Clean the skin where the catheter will be inserted.
- Clean their hands before and after touching your catheter. If you do not see your providers clean their hands, please ask them to do so.
- Avoid disconnecting the catheter and drain tube.
- The catheter is secured to the leg to prevent pulling on the catheter.
- Avoid twisting or kinking the catheter.
- Keep the bag lower than the bladder.
- Empty the bag regularly.

What You Can Do to Help Prevent Surgical Site Infections

- Tell your doctor about other medical problems you may have. Health problems such as allergies, diabetes, and obesity could affect your surgery and your treatment.

- Quit smoking. Patients who smoke get more infections. Talk to your doctor about how you can quit before your surgery.
- Do not shave near where you will have surgery. Shaving with a razor can irritate your skin and make it easier to develop an infection.
- You may have some of your hair removed immediately before your surgery using electric clippers if the hair is in the same area where the procedure will occur, however you should not be shaved with a razor. Speak up if someone tries to shave you with a razor before surgery. Ask why you need to be shaved and talk with your surgeon if you have any concerns.
- Ask if you will get antibiotics before surgery.
- After your surgery, make sure that your healthcare providers clean their hands before examining you, either with soap and water or an alcohol-based hand rub. If you do not see your providers clean their hands, please ask them to do so.
- Family and friends who visit you should not touch the surgical wound or dressings and prevent pets from coming into contact with your wound.
- Family and friends should clean their hands with soap and water or an alcohol-based hand rub before and after visiting you. If you do not see them clean their hands, ask them to do so.
- Before you go home, your doctor or nurse should explain everything you need to know about taking care of your wound. Make sure you understand how to care for your wound before you leave the hospital. If you do develop an infection at the hospital, be sure to ask what type of infection you have, whether you need antibiotics for it, what steps you should take to prevent it from spreading, and make plans for follow up care for the infection.
- Always clean your hands before and after caring for your wound.
- Before you go home, make sure you know who to contact if you have questions or problems after you get home.
- If you have any symptoms of an infection, such as redness and pain at the surgery site, drainage, or fever, call your doctor immediately.

What Hospitals Do to Prevent Surgical Site Infections

To prevent surgical site infections, doctors, nurses, and other healthcare providers:

- Clean their hands and arms up to their elbows with an antiseptic agent before the surgery.
- Clean their hands with soap and water or an alcohol-based hand rub before and after caring for each patient.
- May remove some of your hair immediately before your surgery using electric clippers if the hair is in the same area where the procedure will occur. They should not shave you with a razor.
- Wear special hair covers, masks, gowns, and gloves during surgery to keep the surgery area clean.

- Give you antibiotics before your surgery starts. In most cases, you should get antibiotics within 60 minutes before the surgery starts and the antibiotics should be stopped within 24 hours after surgery.
- Clean the skin at the site of your surgery with a special soap that kills germs.

This information was adapted from materials developed by the Centers for Disease Control and Prevention (CDC), the Association for Professionals in Infection Control and Epidemiology (APIC), the Joint Commission, and Society of Healthcare Epidemiology of America (SHEA).

Other useful resources

Access the New Hampshire Healthcare-Associated Infections (HAI) Program website for public reports, guidelines, and other materials at: <http://www.dhhs.nh.gov/dphs/cdcs/hai/index.htm>.

For more information about HAI nationally and patient safety, visit the Centers for Disease Control and Prevention (CDC) website at: <http://www.cdc.gov/HAI/> and <http://www.cdc.gov/HAI/patientSafety/patient-safety.html>.


The Agency for Healthcare Quality and Research (AHRQ) has information for patients including care planning, diagnosis and treatment, and patient engagement. Visit their website at: <http://www.ahrq.gov/patients-consumers/index.html>.

The Society for Healthcare Epidemiology of America (SHEA) has several patient resources and guides. Visit their website at: <https://www.shea-online.org/index.php/practice-resources/patients>

The Association of Professionals in Infection Control and Epidemiology (APIC) have infographics, eCards, and a quiz about HAI. Visit their website to learn more: <http://consumers.site.apic.org/>.

To learn more about accreditation, certification and standards, visit the Joint Commission Website at: <http://www.jointcommission.org/>.

APPENDIX 5: Map of New Hampshire Hospitals, 2018



REFERENCES

- ⁱ The Centers for Disease Control and Prevention (CDC). Healthcare-Associated Infections Progress Report. March 33, 2016. Accessed online from: <http://www.cdc.gov/HAI/progress-report/index.html>.
- ⁱⁱ Magill SS, Edwards JR, Bamberg W, et al. Multistate Point-Prevalence Survey of Health Care–Associated Infections. *New England Journal of Medicine*. 2014; 370: 1198-208.
- ⁱⁱⁱ Klevens, RM, Edwards RJ, Richards CL, Jr, et al. Estimating health care-associated infections and deaths in U.S. Hospitals, 2002. *Public Health Reports*. 2007; 122(2):160-166. Accessed online from: http://www.cdc.gov/ncidod/dhqp/pdf/hicpac/infections_deaths.pdf
- ^{iv} Humphreys, H, Newcombe RG, Enstone J et al. Four Country Healthcare Associated Infection Prevalence Survey 2006: Risk Factor Analysis. *Journal of Hospital Infection* 2008; 69(3) 249-257.
- ^v Scott R, Douglas. The Direct Medical Costs of Healthcare-Associated Infections in US hospitals and the Benefits of Prevention. March 2009. Accessed online from: http://www.cdc.gov/ncidod/dhqp/pdf/Scott_CostPaper.pdf
- ^{vi} Linda McKibben, MD,^a Teresa Horan, MPH,^b Jerome I. Tokars. Guidance on Public Reporting of Healthcare-Associated Infections: Recommendations of the Healthcare Infection Control Practices Advisory Committee. *American Journal of Infection Control*. 2005;33:217-26. Accessed online from: <http://www.cdc.gov/ncidod/dhqp/pdf/hicpac/PublicReportingGuide.pdf>
- ^{vii} The Healthcare-Associated Infection Working Group of the Joint Public Policy Committee. Essentials of Public Reporting of Healthcare-Associated Infections: A Tool Kit. Accessed online from: http://www.cdc.gov/ncidod/dhqp/pdf/ar/06_107498_Essentials_Tool_Kit.pdf
- ^{viii} Dudeck MA, Edwards JR, Peterson KD, Mu Y, et al. National Healthcare Safety Network (NHSN) report: Data summary for 2013, issued March 2015. *American Journal of Infection Control*. 2015; 43:206-221. Accessed online from: <https://www.sciencedirect.com/science/article/pii/S0196655314013546>
- ^{ix} CDC. CDC. Pooled proportion of healthcare personnel reported as vaccinated in acute care hospital reporting to NHSN, by personnel group and state, 2017-2018 influenza season. Accessed online from: <https://www.cdc.gov/nhsn/pdfs/datastat/hcp-flu-vax-data-tables-hospitals-2018-508.pdf>
- ^x Liddell FD. Simple exact analysis of the standardised mortality ratio. *Journal of Epidemiology and Community Health*, 1984; 38:85-88.
- ^{xi} CDC. Template for State Healthcare Associated Infections Plans. Accessed online from: <http://www.cdc.gov/HAI/pdfs/stateplans/del.pdf>

Note: referenced according to citation order of appearance within the report.