




DISEASE	RECOMMENDED RX	DOSE/ROUTE	ALTERNATIVES
BACTERIAL VAGINOSIS	metronidazole oral ¹ OR metronidazole gel ¹ OR clindamycin cream ^{1,2}	500 mg orally 2x/day for 7 days 0.75%, one 5 g applicator intravaginally 1x/day for 5 days 2%, one 5 g applicator intravaginally at bedtime for 7 days	tinidazole 2 g orally 1x/day for 2 days OR tinidazole 1 g orally 1x/day for 5 days OR clindamycin 300 mg orally 2x/day for 7 days OR clindamycin ovules 100 mg intravaginally at bedtime for 3 days ²
Pregnancy	metronidazole oral ¹ OR metronidazole gel ¹ OR clindamycin cream ^{1,2}	500 mg orally 2x/day for 7 days 0.75%, one 5 g applicator intravaginally 1x/day for 5 days 2%, one 5 g applicator intravaginally at bedtime for 7 days	clindamycin 300 mg orally 2x/day for 7 days OR clindamycin ovules 100 mg intravaginally at bedtime for 3 days ²
CERVICITIS	azithromycin OR doxycycline ³	1 g orally in a single dose 100 mg orally 2x/day for 7 days ⁴	
CHLAMYDIA (CT) Uncomplicated genital/rectal/pharyngeal	azithromycin OR doxycycline ³	1 g orally in a single dose 100 mg orally 2x/day for 7 days	erythromycin base ⁵ 500 mg orally 4x/day for 7 days OR erythromycin ethylsuccinate ⁶ 800 mg orally 4x/day for 7 days OR levofloxacin ⁷ 500 mg orally 1x/day for 7 days OR ofloxacin ⁷ 300 mg orally 2x/day for 7 days
Pregnancy	azithromycin	1 g orally in a single dose	★ amoxicillin 500 mg orally 3x/day for 7 days OR erythromycin base ^{5,8} 500 mg orally 4x/day for 7 days OR erythromycin oddslot base 250 mg orally 4x/day for 14 days OR erythromycin ethylsuccinate 800 mg orally 4x/day for 7 days OR erythromycin ethylsuccinate 400 mg orally 4x/day for 14 days
EPIDIDYMITIS ^{9, 10} For acute epididymitis most likely caused by CT and GC	ceftriaxone PLUS doxycycline	250 mg IM in a single dose 100 mg orally 2x/day for 10 days	<h2 style="text-align: center;">SEXUALLY TRANSMITTED DISEASES:</h2> <h3 style="text-align: center;">SUMMARY OF 2015 CDC TREATMENT GUIDELINES</h3> <p style="text-align: center;">PREPARED BY THE NNPTC, NCSO, AND ASTDA, WITH SUPPORT FROM THE CDC DIVISION OF STD PREVENTION</p> <p style="text-align: center;">COMPLETE GUIDELINES CAN BE ORDERED AT WWW.CDC.GOV/STD/TREATMENT</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>NCSO National Coalition of STD Directors <i>Promoting Sexual Health Through STD Prevention</i></p> </div> <div style="text-align: center;">  <p>American Sexually Transmitted Diseases Association</p> </div> </div> <div style="display: flex; justify-content: space-around; align-items: center; margin-top: 20px;"> <div style="text-align: center;">  <p>National Network of STD Clinical Prevention Training Centers</p> </div> <div style="text-align: center;"> <p>STD CLINICAL CONSULTATION IS AVAILABLE IN 1-5 BUSINESS DAYS BY VISITING WWW.STDCCN.ORG</p> </div> </div>
For acute epididymitis most likely caused by CT and GC and enteric organisms (men who practice insertive anal sex)	ceftriaxone PLUS levofloxacin OR ofloxacin	250 mg IM in a single dose 500 mg orally 1x/day for 10 days 300 mg orally 2x/day for 10 days	
For acute epididymitis most likely caused by enteric organisms	levofloxacin OR ofloxacin	500 mg orally 1x/day for 10 days 300 mg orally 2x/day for 10 days	
GENITAL HERPES First clinical episode of genital herpes	acyclovir OR acyclovir OR valacyclovir OR famciclovir ¹¹	400 mg orally 3x/day for 7-10 days ¹² 200 mg orally 5x/day for 7-10 days ¹² 1 g orally 2 x/day for 7-10 days ¹² 250 mg orally 3x/day for 7-10 days ¹²	
Episodic therapy for recurrent genital herpes	acyclovir OR acyclovir OR acyclovir OR valacyclovir OR valacyclovir OR famciclovir ¹¹ OR famciclovir ¹¹ OR famciclovir ¹¹	400 mg orally 3x/day for 5 days 800 mg orally 2x/day for 5 days 800 mg orally 3x/day for 2 days 500 mg orally 2x/day for 3 days 1 g orally 1x/day for 5 days 125 mg orally 2x/day for 5 days 1 g orally 2x/day for 1 day 500 mg orally once, followed by 250 mg 2x/day for 2 days	
Suppressive therapy ¹³ for recurrent genital herpes	acyclovir OR valacyclovir OR valacyclovir OR famciclovir ¹¹	400 mg orally 2x/day 500 mg orally 1x/day 1 g orally 1x/day 250 mg orally 2x/day	
Recommended regimens for episodic infection in persons with HIV infection	acyclovir OR valacyclovir OR famciclovir ¹¹	400 mg orally 3x/day for 5-10 days 1 g orally 2x/day for 5-10 days 500 mg orally 2x/day for 5-10 days	
Suppressive therapy for pregnant women (start at 36 weeks gestation)	acyclovir OR valacyclovir	400 mg orally 3x/day 500 mg orally 2x/day	
Recommended regimens for daily suppressive therapy in persons with HIV infection	acyclovir OR valacyclovir OR famciclovir ¹¹	400-800 mg orally 2-3x/day 500 mg orally 2x/day 500 mg orally 2x/day	
GENITAL WARTS ¹⁴ (Human Papillomavirus) External genital and perianal warts	Patient Applied ★ imiquimod 3.75% or 5% cream ^{2,11} OR podofilox 0.5% solution or gel ¹⁴ OR sinecatechins 15% ointment ^{2,11} Provider Administered cryotherapy OR trichloroacetic acid OR bichloroacetic acid 80%-90% OR surgical removal	See complete CDC guidelines.	★ podophyllin resin 10%–25% in compound tincture of benzoin may be considered for provider-administered treatment if strict adherence to the recommendations for application. intralesional interferon OR photodynamic therapy OR topical cidofovir
GONORRHEA (GC) ¹⁵ Uncomplicated genital/rectal infections	Dual therapy with ceftriaxone PLUS azithromycin	250 mg IM in a single dose 1 g orally in a single dose	Dual therapy with cefixime ¹⁶ 400 mg orally PLUS azithromycin 1 g orally If cephalosporin allergy or IgE mediated penicillin allergy: gemifloxacin 320 mg orally PLUS azithromycin 2 g orally OR gentamicin 240 mg IM PLUS azithromycin 2 g orally
Pharyngeal ¹⁷	Dual therapy with ceftriaxone PLUS azithromycin	250 mg IM in a single dose 1 g orally in a single dose	If cephalosporin allergy or IgE mediated penicillin allergy, limited data exist on alternatives. See footnotes ¹⁸
Pregnancy	Dual therapy with ceftriaxone PLUS azithromycin	250 mg IM in a single dose 1 g orally in single dose	cefixime ¹⁶ 400 mg orally in single dose PLUS azithromycin 1g orally in single dose If cephalosporin allergy or IgE mediated penicillin allergy, consult with specialist, see footnotes ¹⁹

DISEASE	RECOMMENDED RX	DOSE/ROUTE	ALTERNATIVES
LYMPHOGRANULOMA VENEREUM	doxycycline ³	100 mg orally 2x/day for 21 days	erythromycin base 500 mg orally 4x/day for 21 days
NONGONOCOCCAL URETHRITIS (NGU)^{20, 21}	azithromycin OR doxycycline	1 g orally in a single dose 100 mg orally 2x/day for 7 days	erythromycin base ⁵ 500 mg orally 4x/day for 7 days OR erythromycin ethylsuccinate ⁶ 800 mg orally 4x/day for 7 days OR levofloxacin 500 mg 1x/day for 7 days OR ofloxacin 300 mg 2x/day for 7 days
PEDICULOSIS PUBIS	permethrin 1% cream rinse OR pyrethrins with piperonyl butoxide	Apply to affected area, wash off after 10 minutes Apply to affected area, wash off after 10 minutes	malathion 0.5% lotion, apply to affected area and wash off after 8-12 hours OR ivermectin 250 µg/kg, orally repeated in 2 weeks
PELVIC INFLAMMATORY DISEASE^{22, 23} Non-pregnant patients	Parenteral Regimens EITHER cefotetan OR cefoxitin PLUS doxycycline ³ OR clindamycin PLUS gentamicin Recommended Intramuscular/Oral Regimens EITHER ceftriaxone OR cefoxitin WITH probenecid PLUS doxycycline ³ WITH metronidazole if BV is present or cannot be ruled out	2 g IV every 12 hours 2 g IV every 6 hours 100 mg orally or IV every 12 hours 900 mg IV every 8 hours 2 mg/kg IV or IM followed by 1.5 mg/kg IV or IM for 8 hours 250 mg IM in a single dose 2 g IM in a single dose 1 g orally administered concurrently in a single dose 100 mg orally 2x/day for 14 days 500 mg orally 2x/day for 14 days	Parenteral Regimens ampicillin/sulbactam 3 g IV every 6 hours PLUS doxycycline 100 mg orally or IV every 12 hours The complete list of recommended regimens including regimens for pregnant women can be found in CDC's 2015 STD Treatment Guidelines
SCABIES	permethrin 5% cream OR ivermectin	Apply to all areas of body from neck down, wash off after 8-14 hours 200 µg/kg orally, repeated in 2 weeks	lindane 1% ^{24, 25} 1 oz. of lotion or 30 g of cream, applied thinly to all areas of the body from the neck down, wash off after 8 hours
SYPHILIS Primary, secondary, or early latent <1 year	benzathine penicillin G	2.4 million units IM in a single dose	doxycycline ³ 100 mg 2x/day for 14 days OR tetracycline ³ 500 mg orally 4x/day for 14 days
Latent >1 year, latent of unknown duration	benzathine penicillin G	2.4 million units IM in 3 doses each at 1 week intervals (7.2 million units total)	doxycycline ³ 100 mg 2x/day for 28 days OR tetracycline ³ 500 mg orally 4x/day for 28 days
Neurosyphilis and ocular ²⁶ syphilis	aqueous crystalline penicillin G	18–24 million units per day, administered as 3–4 million units IV every 4 hours or continuous infusion, for 10–14 days	procaine penicillin G 2.4 million units IM 1x daily PLUS probenecid 500 mg orally 4x/day, both for 10-14 days
Pregnancy Primary, secondary, or early latent <1 year	benzathine penicillin G	2.4 million units IM in a single dose	None ²⁷
Pregnancy Latent >1 year, latent of unknown duration	benzathine penicillin G	2.4 million units IM in 3 doses each at 1 week interval (7.2 million units total)	None ²⁷
Pregnancy Neurosyphilis and ocular syphilis	aqueous crystalline penicillin G	Same info as above under neurosyphilis	procaine penicillin G, 2.4 million units IM 1x/day
TRICHOMONIASIS	metronidazole ²⁹ OR tinidazole ²⁸	2 g orally in a single dose 2 g orally in a single dose	metronidazole ²⁸ 500 mg 2x/day for 7 days
Persistent or recurrent trichomoniasis	metronidazole ²⁹ If this regimen fails: metronidazole ²⁹ OR tinidazole ²⁸ If this regimen fails, susceptibility testing is recommended.	500mg orally 2x/day for 7 days 2 g orally for 7 days 2 g orally for 7 days	

- The recommended regimens are equally efficacious.
- These creams are oil-based and may weaken latex condoms and diaphragms. Refer to product labeling for further information.
- Should not be administered during pregnancy, lactation, or to children <8 years of age.
- If patient lives in community with high GC prevalence, or has risk factors (e.g. age <25 years, new partner, partner with concurrent sex partners, or sex partner with a STD), consider empiric treatment for GC.
- If patient cannot tolerate high-dose erythromycin base schedules, change to 250 mg 4x/day for 14 days.
- If patient cannot tolerate high-dose erythromycin ethylsuccinate schedules, change to 400 mg orally 4 times a day for 14 days.
- Contraindicated for pregnant or lactating women.
- Erythromycin estolate is contraindicated during pregnancy.
- Patients who do not respond to therapy (within 72 hours) should be re-evaluated.
- For patients with suspected sexually transmitted epididymitis, close follow-up is essential.
- No definitive information available on prenatal exposure.
- Treatment may be extended if healing is incomplete after 10 days of therapy.
- Consider discontinuation of treatment after one year to assess frequency of recurrence.
- Vaginal, cervical, urethral meatal, and anal warts may require referral to an appropriate specialist.
- CDC recommends that treatment for uncomplicated gonococcal infections of the cervix, urethra, and/or rectum should include dual therapy, i.e., both a cephalosporin (e.g., ceftriaxone) plus azithromycin.
- CDC recommends that cefixime in combination with azithromycin or doxycycline be used as an alternative when ceftriaxone is not available. Oral cephalosporins give lower and less-sustained bactericidal levels than ceftriaxone 250 mg; limited efficacy for treating pharyngeal GC.
- Only ceftriaxone is recommended for the treatment of pharyngeal infection. Providers should inquire about oral sexual exposure.
- Dual therapy with gemifloxacin 320 mg po plus azithromycin 2 g po or gentamicin 240 mg IM plus azithromycin 2 g po are potential alternatives. ID specialist consult may be prudent. Azithromycin monotherapy is no longer recommended due to resistance concerns and treatment failure reports. Pharyngeal GC patients treated with an alternative regimen should have a test of cure (with culture or nucleic acid amplification test) 14 days after treatment.

- Every effort should be made to use a recommended regimen. Test-of-cure follow-up (preferably by NAAT) 3-4 weeks after completion of therapy is recommended in pregnancy. In case of allergy to both alternative and recommended regimens, consult with the STD Clinical Consultation Network at www.stdccn.org.
- Mycoplasma genitalium is the most common cause of recurrent/persistent NGU. Men initially treated with doxycycline should be treated with azithromycin 1g orally. Men who fail a regimen of azithromycin should be treated with moxifloxacin 400mg orally 1x/day for 7 days (effective against M. genitalium)."
- In areas of high trichomonas prevalence, men who have sex with women should also be treated with metronidazole OR tinidazole 2 g orally in a single dose. MSM are unlikely to benefit from the addition of nitroimidazoles.
- Testing for gonorrhea and chlamydia is recommended because a specific diagnosis may improve compliance and partner management because these infections are reportable by state law.
- Evaluate for bacterial vaginosis and trichomonas. Even if bacterial vaginosis is present or cannot be ruled out, treatment with parenteral therapy for the first 24 hours followed by 14 days of therapy is sufficient as described above. If trichomonas is also present, treat per above guidelines.
- Contraindicated for pregnancy or lactating women, or children <10 years of age.
- Do not use after a bath; should not be used by persons who have extensive dermatitis.
- Some specialists recommend 2.4 million units of benzathine penicillin G once weekly for up to 3 weeks after completion of neurosyphilis treatment.
- Pregnant patients allergic to penicillin should be treated with penicillin after desensitization.
- Randomized controlled trials comparing single 2 g doses of metronidazole and tinidazole suggest that tinidazole is equivalent to, or superior to, metronidazole in achieving parasitologic cure and resolution of symptoms.
- Pregnant patients can be treated with 2 g single dose.
- ★ Indicates update from the 2010 CDC Guidelines for the Treatment of Sexually Transmitted Diseases.

Download the STD Treatment Guidelines App in iTunes or Google Play