


New Hampshire Suicide Prevention



Annual Report 2020

This report was produced by NAMI New Hampshire, the State Suicide Prevention Council (SPC) and Youth Suicide Prevention Assembly (YSPA).

Any individual or organization may freely copy and distribute this report.
Electronic copies are available at www.TheConnectProgram.org/annual-reports.

Table of Contents

<u>Introduction</u>	3
What's New in This Year's Report?	3
<u>Primary Partners</u>	4
<u>Accomplishments of Suicide Prevention Efforts in NH</u>	7
State Suicide Prevention Council	7
Youth Suicide Prevention Assembly.....	9
The NH Suicide Survivor Network.....	10
Other Statewide Initiatives	12
Annual NH Suicide Prevention Conference.....	16
NH Grown National Initiatives	16
<u>2020 Data</u>	20
Introduction	20
Demographic Profile of New Hampshire	21
The Big Picture: Suicide in NH and Nationally	23
Youth and Young Adult Suicide in NH	27
Older Adult Suicide in NH	29
Suicide Across the Lifespan in NH	30
Suicide Behavior in NH: Gender Differences	36
Suicide in NH: Methods	50
Linking At-Risk Individuals with Help	59
Costs of Suicide and Suicidal Behavior	60
Military and Veterans.....	61
Suicide Rates in NH	64
<u>Reading Tables and Figures</u>	68
<u>Glossary of Terms</u>	70
<u>Reliability of Rates</u>	73
<u>Frequently Asked Questions about NH Suicide Data</u>	74
<u>Contacts and Meeting Information</u>	76
<u>Recognize the Warning Signs of Suicide</u>	78
<u>Mental Health and Suicide Prevention Resources</u>	79

Introduction

The 2020 Annual Suicide Prevention Report, which includes a summary of accomplishments and data, is the result of the collaborative work of many groups, committees, and organizations in NH who have dedicated time and resources to study the issue of suicide and to look at prevention and postvention across the lifespan.

The work of these groups in suicide prevention and postvention is reaching across the state and into communities, schools, organizations, and individual lives.

Evidence of this includes some of the following accomplishments from calendar year 2020:

- Work commencing on a new State Suicide Prevention Plan.
- Holding the first-ever virtual NH Suicide Prevention Conference in response to the COVID-19 pandemic.
- The continued work being done in the area of Zero Suicide by representatives from the health and behavioral health care fields, Department of Corrections, Department of Safety, and military organizations.

Many achievements will be described further throughout this report. It is critical to NH that we continue to build on the momentum and collective knowledge that has been gained in suicide prevention to strengthen capacity and sustainability in order to reduce risk of suicide for all NH residents and promote healing for all of those affected by suicide.

Knowing that it takes all of us working together with common passion and goals, thank you to everyone who has been involved in suicide prevention and postvention efforts in New Hampshire.

What's New in this Year's Report?

Some of the new highlights this year include:

- Updated NH Violent Death Reporting System data, including newly available data on sexual orientation.
- Highlights of activities that took place across the state.
- New examples of positive outcomes and testimonials related to suicide prevention work being done in NH. These examples are included as text boxes interspersed throughout the report.

Primary Partners

NAMI New Hampshire and The Connect Suicide Prevention Program

NAMI New Hampshire (National Alliance on Mental Illness), a grassroots organization of families, individuals living with mental illness, professionals and other members, is dedicated to improving the quality of life of persons of all ages affected by mental illness and suicide through education, support, and advocacy.

NAMI NH's Connect Suicide Prevention Program has been recognized as a best practice and model for a comprehensive, systemic approach. The community-based approach of the Connect Program focuses on education about early recognition (prevention); skills for responding to attempts, thoughts, and threats of suicide (intervention); and reducing risk and promoting healing after a suicide (postvention). NAMI NH and The Connect Program assist the State Suicide Prevention Council and the Youth Suicide Prevention Assembly with implementation of the NH Suicide Prevention Plan. Connect provides consultation, training, technical assistance, information, and resources regarding suicide prevention throughout the state. NH-specific data, news and events, information and resources, and supports to survivors are available on the Connect website at www.TheConnectProgram.org.

New Hampshire Office of the Chief Medical Examiner

The New Hampshire Office of the Chief Medical Examiner (OCME) is responsible for determining the cause and manner of all sudden, unexpected or unnatural deaths falling under its jurisdiction (RSA 611-B:11). This includes all suicide deaths occurring within the state of NH. As the central authority making these determinations, the OCME is in an ideal position to provide timely data on NH suicide deaths. For more than 15 years the OCME has partnered with YSPA, and more recently the SPC, to provide data and insight into the deaths affecting the state.

New Hampshire Violent Deaths Reporting System

In 2015, NH partnered with the Centers for Disease Control and Prevention (CDC) Injury Prevention Division and began a joint surveillance program, also known as the National Violent Death Reporting System (NVDRS), which is now applied in all fifty US states and Puerto Rico. The surveillance program in NH is known as the NH Violent Death Reporting System (NH-VDRS), which is supported by CDC NVDRS grant funding. NH-VDRS is housed in the NH Office of the Chief Medical Examiner (OCME) with administrative oversight by the NH Department of Health and Human Services Injury Prevention Program as the grant holder. The NH-NVDS program is tasked with compiling case level data on all violent deaths in NH, including suicides, homicides, all deaths involving firearms, and deaths resulting from legal intervention (such as law enforcement or war). The NH-VDRS program's work also entails disseminating information within NH and to the CDC Injury Prevention Division and other affiliates. Since its inception, NH-VDRS has endeavored to engage entities focusing on suicide in NH, including local suicide prevention service providers, suicide prevention advocates, law enforcement, lawmakers and other interested groups. These groups are making use of aggregate

data reported by NH-VDRS to enhance prevention efforts in the state. The NH-VDRS data in this report is made possible under Grant Award # 5NU17CE924939-02-00.

For information regarding NH-VDRS or to request data, contact:

- JoAnne Miles-Holmes, Injury Prevention Program Manager, NH-VDRS Principal Investigator, Maternal and Child Health Section, Division of Public Health Services, NH Department of Health and Human Services, JoAnne.MilesHolmes@dhhs.nh.gov.
- Djelloul Fourar-Laidi, Planning Analyst/NH-VDRS Planning Analyst-Data Systems, Office of the Chief Medical Examiner, NH Department of Justice, djelloul.fourar-laid@doj.nh.gov.
- Kim Fallon, Chief Forensic Investigator, Office of the Chief Medical Examiner, kim.fallon@doj.nh.gov.

State Suicide Prevention Council

The mission of the State Suicide Prevention Council (SPC) is to reduce the incidence of suicide in New Hampshire by accomplishing the goals of the NH Suicide Prevention Plan:

- * Raise public and professional awareness of suicide prevention;
- * Address the mental health and substance misuse needs of all residents;
- * Address the needs of those affected by suicide; and
- * Promote policy change.

The success and strength of the Council is a direct result of the collaboration that takes place within its membership and with other agencies/organizations, including public, private, local, state, federal, military, and civilian. Strong leadership and active participation come from the Council's subcommittees: Communication and Public Education; Data Collection and Analysis; Law Enforcement; Military and Veterans; Public Policy; Suicide Fatality Review; and the Survivors of Suicide Loss subcommittee.

As part of NH RSA 126-R, which legislatively established the Suicide Prevention Council, the Council is required to report on its progress annually, to both the Governor and the legislature. This report serves that purpose, as well as providing an annual update on the accomplishments of our collective achievements and data regarding suicide deaths and suicidal behavior in NH.

Youth Suicide Prevention Assembly

The Youth Suicide Prevention Assembly (YSPA) is dedicated to reducing the occurrence of suicide and suicidal behaviors among New Hampshire's youth and young adults up to the age of 24. This is accomplished through a coordinated approach to providing communities with current information regarding best practices in prevention, intervention, and postvention strategies and by promoting hope and safety in our communities and organizations.

YSPA is an ad hoc committee of individuals and organizations that meet monthly to review the most recent youth suicide deaths and attempts in order to develop strategies for preventing them.

Over the years, YSPA and its partners have been involved with a wide range of suicide prevention efforts in the state. These efforts include but are not limited to collecting and analyzing timely data on suicide deaths and attempts, collaborating on an annual educational conference, creating the original NH Suicide Prevention Plan and identifying the need for statewide protocols and training, which were developed through NAMI NH into The Connect Program. The Survivor of Suicide Loss packets that are sent to the next of kin of anyone who dies by suicide in New Hampshire got their start in YSPA before expanding to all ages.

Accomplishments of Suicide Prevention Efforts in NH

State Suicide Prevention Council

This year marked the 12th anniversary of NH's Suicide Prevention Council (SPC) since its legislative inception in 2008. As part of a revision of the NH Suicide Prevention Plan (<https://www.dhhs.nh.gov/dphs/bchs/spc>) in 2016, the concept of a Zero Suicide approach was adopted by the SPC. In 2020 the SPC began a new revision process on the State Plan with the intention of issuing a revised version in 2021. The Zero Suicide approach was again integrated into the revision process. More information about Zero Suicide is available from <http://zerosuicide.sprc.org/>. A selection of accomplishments in NH related to Zero Suicide are featured below and on page 8. During 2020, the SPC, its subcommittees, and other stakeholders in the state have looked at ways of implementing the goals outlined in the revised Plan.

Much of the work of the SPC is done at the subcommittee level. Some of the subcommittee activities occurring in 2020 to move forward the goals of the NH Suicide Prevention Plan included:

Communications Subcommittee

- Coordinated a press event in September for Suicide Prevention Awareness Week.
- Worked with the Public News Service on writing and publishing stories around prevention efforts in the state. The stories take into account the media recommendations for reporting on suicide (<http://reportingonsuicide.org/>) as well as the National Action Alliance's Framework for Successful Messaging: <http://suicidepreventionmessaging.org/>.

Data Collection and Analysis Subcommittee

- Worked with multiple statewide partners to compile and analyze data covering calendar year 2019. The data were then included in the 2019 NH Suicide Prevention Annual Report and distributed statewide.
- Collaborated with the Analyst for the NH Violent Death Reporting System (NH-VDRS) to expand the use of NH-VDRS data in the NH Annual Suicide Prevention Report.

Military and Veterans Subcommittee

Military and Veterans Subcommittee members represent a variety of military & civilian groups and public & private sectors and regions of New Hampshire. This multi-discipline and expansive network has helped to enhance a common understanding and approach to suicide prevention efforts in NH for Service Members, Veterans, and their Families (SMVF).

The following outcomes and accomplishments resulted from the work of the Military and Veterans Subcommittee in 2020:

- The Committee continues to be actively involved with the *Governor's Challenge to Prevent Suicide among Service Members, Veterans and their Families* which is funded

through the Substance Abuse and Mental Health Services Administration (SAMHSA). This initiative continues to provide members of the Committee with a variety of technical assistance and free training related to best practices to reduce the rate of suicide in the military-connected community.

- Committee members promoted military cultural competency and suicide prevention awareness among community providers by presenting at various conferences and workshops throughout 2020 including the Halias Symposium, SAMHSA Academies, NH Behavioral Health Summit, Veterans Diversity Conference, and others.
- The Committee supported Northeast Passage in its application for a federal grant to develop a Wellness Series for Veterans in the Lakes Region. The funds were approved, and Northeast Passage began providing a variety of recreational and wellness workshops for Veterans in the Lakes Region area even throughout the time of the pandemic. Some activities were provided virtually, but many were and continue to be provided in person at various locations in the area.
- The NH Department of Military Affairs and Veterans Services website now hosts a page – Suicide Prevention Efforts – (<https://www.dmav.snh.gov/about-us/councils-and-committees/suicide-prevention-efforts>)—where the Committee’s efforts are described, and meeting minutes are publicly available.

Survivor of Suicide Loss Subcommittee

- Began a monthly coffee chat in December of 2020. The coffee chat offers Survivors of Suicide Loss access to a variety of resources as well as an opportunity to support fellow survivors through their loss and healing journeys.
- Provided support and technical assistance to NH Loss Survivor Support Groups and promoted the American Foundation for Suicide Prevention (AFSP) International SOSL Day.
- Ensured Loss Survivor participation in community events through targeted outreach.

As the council looks to continue its work, there is a desire to increase active membership on its subcommittees. This is particularly true for representatives from the field of substance use disorders as the council looks at the relationship between substance misuse and suicide prevention. The council also recognizes the role public health departments play in this work and the importance of their perspective for future collaborations. The public private partnerships developed in subcommittees should continue to expand and enhance the impact of the work being done by the council. Contact* any of the committee chairs if you have an initiative you would like to put forward related to suicide prevention efforts throughout the state.

The council continues to collaborate with the Department of Health and Human Services (DHHS) for statewide leadership and support as it looks to continue its work in promoting evidence-based initiatives and refining and expanding the state plan to ensure the very best outcomes for NH citizens.

**If you would like to join any of the Suicide Prevention Council Subcommittees, please contact the designated committee chair. The committee meeting schedule has been included on pages 76-77 of this report.*

The Youth Suicide Prevention Assembly (YSPA)

YSPA is a grassroots organization comprised of individuals interested in learning more about how to prevent all suicides, but especially those that occur among individuals age twenty-four and under. YSPA supports the State Suicide Prevention Plan by promoting a greater awareness of youth/young adult suicide risk factors, protective factors, and warning signs. YSPA encourages the development and maintenance of professional networks, and the use of natural supports to lessen the risk of suicide and promote support and postvention activities in the event of a suicide death.

YSPA, as with the rest of the world, faced some challenges on how to adapt to COVID-19. The April meeting was canceled as part of a statewide response to the “Stay at Home” Governor’s order in March. The co-chairs held a conference call and decided to address the challenges as follows:

- A message, including photos of the co-chairs to make it more personal, was sent to the mailing list acknowledging the struggles that they and the individuals they serve were likely experiencing, including higher anxiety.
- A document about the importance of home safety (in response to reports of more firearms being purchased, increase in anti-anxiety medications being prescribed, and the increased social isolation) was sent with this message. The document can be found at <https://theconnectprogram.org/YSPA-HSG>.

YSPA meetings resumed in May 2020 via Zoom. Case reviews were not part of the meetings due to confidentiality concern related to the virtual aspect of the meetings. The meetings allowed the opportunity to counter the (at the time) prevailing message that suicides were increasing because of the pandemic. New Hampshire is fortunate to have access to “real time” data; there was no indication then that this was the case.

Speakers, or specific topics, have always been part of YSPA. This did not change because of COVID-19. Speakers/topics for 2020 YSPA meetings included the following: a presentation by a representative from the Northern New England Poison Control Center, the annual “retreat” when the Mission Statement and brochure are reviewed, input is solicited from the YSPA membership (the “Assembly” of YSPA) as to meeting structure, data and case reviews and other aspects of the meetings. The annual data summary for the prior year was presented outlining common themes for YSPA-age deaths (mostly males, mostly in the 19-24 range). A riveting NAMI *In Our Own Voice* speaker, Jace Troie, captivated the August attendees. Two presentations that had been postponed in the spring were rescheduled for the fall. One was about “First Responder Mental Health” presented by a Derry Police Officer (and relative of an YSPA attendee); the other was about First Episode Psychosis.

YSPA co-chairs faced a difficult decision as to whether or how to resume the meeting case reviews. Consideration was made as to how the information obtained from the reviews is helpful in possibly preventing future suicides, yet the virtual format could raise issues of confidentiality and lack of support. Structures were developed to ensure that meeting attendees followed strict confidentiality guidelines and that the omnipresent “support person” for meetings was able to virtually provide support.

Case reviews resumed in December 2020 following an 8-month gap.

YSPA membership continues to be diverse with regular membership representing behavioral health, substance use, all levels of education, law enforcement, LGBTQ+ groups, public health, social service agencies and persons with lived experience. Virtual meetings have allowed for attendees from all parts of the state to attend, which has been one benefit. The Youth Suicide Prevention Assembly (YSPA) continues to meet the second Thursday of every month in Concord, NH.

For more information on YSPA, please contact Elizabeth Fenner-Lukaitis: Elizabeth.V.Fenner-Lukaitis@dhhs.nh.gov or Elaine de Mello: edemello@naminh.org.

The NH Suicide Survivor Network

In 2020 Survivors of Suicide Loss (SOSL) continued in their efforts of building capacity and establishing groups throughout NH. More and more loss survivors are finding comforting support in their healing journey and continue to mentor each other in facilitating and co-facilitating these groups by providing a safe environment to share their experience of suicide loss. The year began with 15 groups already in motion and attendees growing in numbers. As a result of the COVID-19 pandemic, many of these groups were forced to temporarily halt in-person meetings. In order to continue to support loss survivors, many of the groups began holding virtual meetings. These support groups continue to meet on a weekly, bi-weekly, and monthly basis. When it is again safe to do so, they will resume in-person meetings. Talk of teen (ages 14-18) and young adult (ages 18-25) Survivor of Suicide Loss peer support groups, while delayed by the pandemic, has continued to progress.

Positive Outcomes and Testimonials

Both Sides of the Door - Law Enforcement Investing in Loss Survivors!

Several Loss Survivors have experienced an extremely difficult situation at the scene of a suicide death in their home. Loss Survivors are in complete shock and disbelief upon finding out about this tragedy – accompanied by feelings of grief, sadness, and devastation. The last thing they want is to be separated from their family and their loved one they just lost to suicide.

Through the chaos of a suicide death, most often Loss Survivors aren't given any information during the investigation and Loss Survivors are led to feel like a suspect in their own home and loved one's death.

Goffstown's law enforcement is one step ahead of this for Loss Survivors. Their goal is to "invest" in Loss Survivors, recognizing the importance of treating Loss Survivors with the utmost respect and compassion at the scene of a suicide death and thereafter. With their police department chaplain, they work together to make this unimaginably tragic situation run as smoothly as it can.

Many law enforcement departments in NH also have something like this in place – through the Laconia Police and the Partnership for Public Health in this region a protocol for unattended death/death notification has been put together for all law enforcement to help remind them of what can be done and said to Loss Survivors at the time of a suicide death and an unattended death as well. These two examples help to make a tragic situation such as a suicide death go a little more smoothly – Loss Survivors are better understood, and law enforcement recognizes the importance of compassionate approach.

An ever-growing number of Loss Survivor Speakers continued to share their personal stories and experiences of suicide loss to help educate the public and provide healing and support, within their communities and throughout the state. In 2020, many of these presentations took place as part of virtual events, such as the NH Suicide Prevention Conference. The NH Survivors of Suicide Loss Resource Packet was updated and disseminated through the NH Office of the Chief Medical Examiner to the next of kin of all those who died by suicide. The book “Healing the Hurt Spirit: Daily affirmations for people who have lost a loved one to suicide,” authored by a NH survivor, continues to be available to new loss survivors. An online survey is also provided to solicit feedback on the folder and provide additional avenues to connect loss survivors to help.

Viewings of the American Foundation for Suicide Prevention (AFSP) International Survivors of Suicide Loss Day (ISOSLD or Survivor Day) are normally held at multiple sites throughout NH on the Saturday before Thanksgiving. In 2020 the event was fully virtual and resulted in 78 individuals registering to participate remotely.

The annual NH Survivor of Suicide Loss Newsletters were distributed throughout the state, with hard copies made available at trainings, loss survivor speaking presentations, the State Suicide Prevention Conference, health fairs, libraries, hospitals, healthcare facilities, mental health centers, funeral homes, churches and faith-based organizations, and in the Survivors of Suicide Loss Resource Packet. The newsletter was also distributed electronically to many email lists.

More and more loss survivors in NH are becoming involved in advocacy and fundraising efforts for various local and national suicide prevention organizations and initiatives. NH loss survivors volunteered over 1,000 hours of time by hosting support groups, sharing loss survivor resources at community events, speaking publicly about their loss, or displaying the quilts that were lovingly crafted by NH Survivors of Suicide Loss in memory of their loved ones lost to suicide.

The NH State Suicide Prevention Council continues to include the Survivors of Suicide Loss Subcommittee on the council, and to include loss survivors on the membership of the Council and its other subcommittees. Feedback from the NH loss survivor network clearly indicates great interest by loss survivors in expressing their voice, building capacity of support groups, expanding the International Survivors of Suicide Teleconference Day, and being involved in more advocacy and public speaking events.

This committee encourages new members to join and attend their monthly conference calls.

Positive Outcomes and Testimonials

“The resources for survivors are critical and every effort must be made to keep and improve their availability. Many survivors would not be functioning, healing or grieving if it were not for these programs. For a situation which is not understood by a large percentage of society, support and education still remain a priority.”

A New Hampshire Survivor of Suicide Loss

Attempt Survivor Initiative:

An Attempt Survivor Committee was formed to look at resources and support for individuals in NH who have attempted suicide. The committee had representation from persons with lived experience (loss and attempt survivors), staff from New Hampshire Hospital (NHH), NAMI NH, the Office of Consumer Affairs, and Peer Support Centers in NH. In the course of the committee's work, models for attempt survivor support groups were researched and the committee developed a manual to provide guidance around leading support groups. Currently there are attempt survivor groups in Keene and Berlin.

Other Statewide Initiatives

AFSP (American Foundation for Suicide Prevention)

The NH Chapter made tremendous efforts to increase support for Survivors of Suicide Loss in 2020.

Led by Board Member Ally Sharpe, the chapter continues to offer Healing Conversations to survivors across the state. In an effort to increase capacity and promotion, the chapter secured 3 additional volunteers and is vetting several more. In the final few months of the year, the chapter has consistently received 4-6 requests per month, with each visit completed within 10 days. Following each visit, Ally conducts a debriefing with the volunteers so that the chapter is able to support all involved in the visit in the best way possible.

With social distancing making it impossible to host in-person Survivor Day events, we were unable to offer 8 programs around the state. In an effort to offer a quality program, AFSP NH partnered with NH Healthy Families, the VA, and Debbie Baird from the NH Suicide Prevention Council to host a statewide virtual event. This collaborative effort resulted in 52 of 78 registrants participating online. Prior to the virtual event, every registered attendee was sent a resource packet that included brochures and materials from AFSP, the VA, NAMI NH, and Partnership for Public Health. The packet also included a luminaria and electronic candle to be lit at the end of the program. Loss survivors were encouraged to send in a picture of their loved one for the Memoriam Slideshow presented at the beginning of the program. Participants watched an AFSP documentary and were offered small group discussions in Zoom breakout rooms by loss. The chapter has been able to increase collaborations to support loss survivors with Area Director Shamera Simpson's new role as Co-Chair of the NH Suicide Prevention Council's Survivors of Suicide Loss Subcommittee. In this role, Shamera established a monthly coffee chat to allow an opportunity for survivors to connect with other survivors and to various resources from around the state.

In 2020, AFSP NH hosted its first State Advocacy Day. The program offered opening remarks from NH Senator David Watters and an agenda that included a background on AFSP, Advocacy 101, current state legislative issues, and how to take action. Over 45 constituents and partner organizations attended. The NH Chapter has 204 registered field advocates with 54% having taken action.

Finally, the NH Chapter was chosen as a Charity of Choice by Townsquare Media as part of the Year of Service Award. The award allowed the chapter to receive close to \$19,000 in free radio advertisements for a 3-month period. The Chapter used a portion of this promotion to focus on Loss and Healing. During the month of September, over 100 radio spots aired on 2 local radio stations as well as 5,376 ad impressions online promoting both Healing Conversations and ISOSLD. During the second half of September, there were 871 views on the Healing Conversations website and 1,263 views on the ISOSLD page.

Connor's Climb Foundation

Connor's Climb Foundation is a New Hampshire based nonprofit with the mission to provide suicide prevention education to NH youth and communities. Despite the challenges faced with COVID-19, the foundation hosted 17 suicide prevention events across the state with 2,426 attendees. Sixteen of these events were focused on the SOS Signs of Suicide® Prevention Program, which is a nationally recognized, evidence-based school suicide prevention program. Connor's Climb provided funding to implement the SOS program with students, faculty, and parents to 30 schools throughout the year. Despite the foundation's largest awareness event, Connor's Climb Annual 5K and Family Walk, being postponed and later held virtually, 311 individuals participated. Senate Bill 282, now LS 193-J, an act relative to suicide prevention education, went into effect in July of 2020, and remains a key priority for the foundation.

New Hampshire Nexus Project 2.0 – Garret Lee Smith (GLS) Grant Funding

Suicide remains the second leading cause of death for 10- to 44-year-olds here in New Hampshire and nationwide. The GLS New Hampshire Nexus Project 2.0 (GLS NHPN 2.0) is a cross-systems, collaborative approach to reducing suicide incidents among youth by improving pathways to care and offering comprehensive training to provide youth serving organizations with the resources to identify, screen, refer, and treat at-risk youth.

GLS NHPN 2.0 is focused on youth/young adults ages 10-24 in the Capital, Carroll County, and North Country Regional Public Health Networks of New Hampshire. Over the course of the five-year project period, project staff and key partners are working in collaboration to enhance care coordination infrastructure, suicide risk recognition and response, and statewide capacity for suicide prevention and postvention response.

Project partners include NAMI NH, the Behavioral Health Improvement Institute, Headrest, Northern Human Services, Riverbend Community Mental Health, Granite United Way, North Country Health Consortium, NHTI, and White Mountains Community College. Additionally, regional implementation teams established early in the project are comprised of key stakeholders across multiple community sectors working together to build and sustain capacity and infrastructure around implementation of best practices for suicide prevention and postvention for high-risk youth and addressing overall access to care issues at a local and systemic level.

GLS Year One Training Opportunities:

Despite navigating challenges presented by the pandemic, year one provided a variety of GLS region specific and statewide training opportunities. *Connect eLearning* options offered an online, self-paced training experience to over 2,400 New Hampshire individuals.

Connect Suicide Prevention Training and Planning Sessions held in October provided a two-day interactive training process for over 35 local stakeholders on national best practices on suicide prevention and intervention, as well as a mind-mapping process to facilitate the creation of a community-wide suicide prevention plan.

Research demonstrates that how media report on suicide may contribute to increased risk of suicide contagion. Our statewide media training provided an opportunity for local media, educators, members of law enforcement, business, public health officials, and municipal leaders to review action steps and guidelines in communicating about and responding to traumatic incidents such as suicide to promote healing and reduce risk for vulnerable populations.

In partnership with Elaine Frank, we conducted a *CALM (Counseling on Access to Lethal Means) Train-the-Trainer* for 12 individuals, including two master trainers.

Training opportunities for our young adult population (18-24) included an *In Our Own Voice* training and *Connect Young Adult Leader Suicide Prevention* training for a total of 7 individuals.

GLS Care Liaisons:

Care Liaisons based at Riverbend Community Mental Health and Northern Human Services are working with youth and young adults 10 to 24 identified as high risk for suicide in their respective regions. Their services include implementation of a suicide care pathway to reduce risk, as well as facilitating stabilization and recovery following a high-risk incident or period utilizing evidence-based approaches for up to 90 days. This brief intervention provides an opportunity for education, as well as to strengthen the individual's paid and natural support system.

Project Outreach

The Behavioral Health Summit is designed to bring together non-profit leaders, health care providers, clinical and licensed professionals, and public policy advocates in the mental health and substance use disorder community. This year the GLS PC, NAMI NH Director of Suicide Prevention Services, & Riverbend Community Mental Health Center staff provided a workshop that examined the Care Liaison role in facilitating pathways to care for youth/young adults at high risk for suicide. It provided an opportunity to highlight this brief intervention and introduce the GLS work happening regionally and statewide.

The 17th Annual NH Suicide Prevention Conference provided an opportunity for the GLS PC to engage individuals in the mental health field and provide a workshop elevating the voices of individuals with lived experience. It also provided another opportunity to communicate the GLS work happening regionally and statewide.

For more information on the GLS NHNP 2.0 contact Susan Ward at sward@naminh.org

Zero Suicide Examples of Efforts in New Hampshire

Hosted by Exeter Hospital, providers around NH came together in November 2017 for a Zero Suicide Academy to kick off strategic efforts to prevent suicide in systems across the state. Since that time many initiatives have been implemented in facilities and organizations in NH. For instance, many emergency departments, health care facilities and mental health centers have been implementing universal screening for suicide risk. While this summary is not all inclusive, some of these efforts in 2020 included the following:

The Foundation for Healthy Communities' Behavioral Health Clinical Learning

Collaborative is a statewide, essential initiative that engages and supports clinicians and staff from hospital emergency departments (EDs), community mental health centers (CMHCs) and other stakeholders to better engage and treat patients in a mental health crisis in the ED setting while they await referral to the most appropriate level of care. The Collaborative is funded by the Endowment for Health and New Hampshire Charitable Foundation.

Strategies created thus far in the Collaborative demonstrate our ability to coordinate and mobilize our partners in creating and connecting with appropriate resources. Examples include: the *Suicide Screening & Intervention Toolkit for New Hampshire Emergency Departments* using the Columbia-Suicide Severity Rating Scale to help ED staff members develop the confidence and know-how to screen quickly for suicidality and access clinical decision support tools to help determine next steps in care; a *Trauma-Informed Care (TIC) Pilot Project* where two hospital/community mental health center teams participated in an accelerated program to study and analyze TIC principles and design initiatives to improve safety and treatment outcomes within their ED settings; a *Behavioral Health Inpatient Admissions Guide* to assist ED and CMHC staff in understanding the admissions process to all voluntary and designated receiving facilities; Patient engagement initiatives including discussions with presenters from NAMI NH's *In Our Own Voice* program who shared their experiences and perspectives as patients boarding in EDs and provided recommendations for care. Collaborative members have connected with and receive ongoing updates from statewide Peer Support Agencies, Crisis Respite Centers, and Step-Up/Step-Down programs.

The Mental Health Center of Greater Manchester (MHCGM) has reported Zero Suicide related accomplishments including decreasing suicide deaths by 44%, training 80% of staff in suicide prevention, and assessing suicide risk in 100% of active clients. MHCGM has posted an extensive outcome story to the Zero Suicide website that can be viewed at <https://zerosuicide.edc.org/evidence/outcome-story/mental-health-center-greater-manchester>

The NH Army National Guard (NHARNG) The NHARNG provided 2 Applied Suicide Intervention Skills Training (ASIST) workshops in 2020, training over 22 key gatekeepers of the military and non-military personnel. For Suicide Prevention Month the NHARNG hosted an awareness building Virtual Suicide Prevention 5k Run/Walk. Over 80 military and community members participated.

Annual NH Suicide Prevention Conference

The NH Annual Suicide Prevention Conference was held virtually on November 12 and 13, 2020.

Leaders from multiple organizations, state agencies, and people with lived experience welcomed attendees in an earnest fashion that brought immediate engagement and trust that advocacy, resources, and support are growing, in spite of the worldwide pandemic. Hope was inspired as attendees joined in the challenge to keep working on prevention and overcoming social isolation while maintaining physical distance. This is a vital factor across the age span.

Concepts of varying levels of mental wellness were presented, in the context of lived experience from several perspectives, to help individuals set goals to achieve and maintain personal wellness.

The conference experience was filled with thoughtful and inspiring speakers, and many resources. Even break time between workshops offered very impressive 2020 data on thousands of individuals trained in suicide prevention, awareness and fundraising events, and support groups for survivors of loss.

NH Grown National Initiatives

Connect

NAMI NH's Connect Suicide Prevention and Postvention program continued to provide training and consultation to organizations, schools and communities across NH and around the U.S., providing evidence-based strategies in responding to individuals at risk for suicide and promoting healing and reducing risk after a suicide death. Due to the pandemic, there was a rapid shift to offering programs virtually in the spring of 2020, and by fall, most of the Connect programs were available virtually in addition to the Connect E-Learning self-paced modules that had been developed prior to the pandemic.

In NH in 2020, there were 1,773 participants trained in Connect Prevention and 279 in Connect Postvention. 16 trainers were trained from schools and mental health centers to help sustain their suicide prevention efforts in their respective organizations.

Positive Outcomes and Testimonials

"This Conference saved my life"

Feedback from an attendee at the Annual NH Suicide Prevention Conference

Positive Outcomes and Testimonials

"I feel more confident if a clinic patient is suicidal, I now know the steps to take to keep a suicidal patient safe. Thank you."

"I thoroughly enjoyed this training as it is a topic that hits close to home and makes me more acutely aware of ways to help and share resources for friends, family, staff, patients, etc."

"I am a triage nurse and some of the most stressful calls for me are the calls from individuals expressing thoughts of self-harm. This information will be a huge help to me! I will also pass it on to others!"

"I really liked that we addressed the 'elephant in the room' and that suicide is more complicated and it takes a village to save someone's life."

Feedback shared by Connect Suicide Prevention Training participants.

At the NH Police Academy, 155 new recruits also received training from Connect staff in suicide prevention and postvention as a standard part of their training curriculum, and the Department of Corrections also began implementing mental health and suicide prevention training for new recruits through NAMI NH.

Two schools in NH also implemented Connect's Youth Leader program, training 82 high school youth to partner with adults to lead this program for peers and teachers in their school and communities as a strong and vibrant protective factor. One young adult leader training was held during the pandemic. Both youth leader and young adult trainings were designed carefully to ensure safety and connectedness for these vulnerable populations.

Staff in the Connect program assisted individuals, schools, and communities in their journeys after a suicide with over 61 hours of postvention support and technical assistance to communities and organizations in NH.

Many schools and organizations took advantage of the self-paced Connect E-Learning Suicide Prevention training for gatekeepers, school personnel, health care, and mental health providers with 4,058 seats distributed in NH during the year.

The Connect Program staff also provided virtual training and consultation throughout the country in 2020 in numerous states, including Florida, Alaska, Ohio, Oregon, South Carolina, Connecticut, and Colorado.

Furthering suicide prevention efforts in NH came with the opportunity to apply for a federal SAMHSA grant, called the Garrett Lee Smith (GLS) grant, focused on suicide prevention for young people through the age of 24. NAMI NH was identified by the state to be the applicant for this grant in 2019, and in early 2020, NAMI NH received notification that its application was approved for a five-year award to focus on young people through the age of 24, with a concentration in the Northern region to include Carroll and Coos Counties, and the Capital Region. The work for this grant started in mid-January, 2020. A summary of the progress achieved with this grant during 2020 is provided on pages 13-14.

Counseling on Access to Lethal Means - CALM

CALM (Counseling on Access to Lethal Means) is a national best practice that was developed here in NH in 2006. Since then, it has been offered as an in-person workshop and Train the Trainer program, several online versions and, since COVID, as a virtual training. As of 2020, there are 10 Master Trainers in NH, NC, MO, and TX who are able to train CALM trainers. CALM was developed to assist health and mental health providers in having

Positive Outcomes and Testimonials

“I feel [the CALM training] was very valuable. I feel that this training will help improve my skills.”

“The data helped me to challenge many of my false perceptions.”

“Excellent and helpful for staff to educate patients and families on prevention and actions to take.”

Feedback shared by 2018 CALM training participants.

effective, respectful conversations about reducing access to lethal means (primarily firearms and medications) for those at risk for suicide.

A shorter version of CALM is being rolled out and evaluated to bring this information to the broader community, with particular emphasis on gun owners. CALM continues to be evaluated to demonstrate and improve its effectiveness.

National Guard and Veterans Affairs Initiatives

Work continued through 2020 to implement a statewide resource coordination center for Service Members, Veterans, and their Families (SMVF) so that SMVF will benefit from better coordinated referrals and more timely access to appropriate services. The Military & Veterans Subcommittee partnered with NH Department of Health & Human Services—Bureau of Elderly & Adult Services to enhance the state’s Servicelink contracts with regards to training and referral procedures for SMVF callers. The new contracts were effective July 1, 2020, and committee members met with the DHHS Administrator and Servicelink Managers to begin more specific work in this area. As these objectives are implemented, staff across the state working in the resource & referral Servicelink organizations will have the training, knowledge, tools, and resources necessary to best serve and connect SMVF in their times of need.

Members of the Military & Veterans Subcommittee worked to bring *Together with Veterans* (TWV), a community-based suicide prevention program for rural Veterans funded by the Veterans Administrative Office of Rural Health to the Lakes Region of New Hampshire. TWV involves partnering with rural Veterans and their communities to implement community-based suicide prevention. It is a 3-year program that provides \$100,000 for communities to utilize in support of their work. The group in New Hampshire identified an official name for themselves—Lakes Region Veterans Coalition—and created a website to promote their work. LRVC also established a Buddy Check Coffee program which provides Veterans with a chance to meet for a coffee, virtually or in person, by providing participants with a free gift card to a coffee shop for connecting with a buddy or being matched to talk with a Veteran they had not yet met.

In 2019, members from the Military & Veterans Subcommittee collaborated with the NH National Guard to bring the Star Behavioral Health Providers training (SBHP) to New Hampshire. This national program provides clinical providers with a high quality, free 3-tier training and free CEUs. In 2020, the Committee collaborated with the NH National Guard and community partners to promote the training series to various social services sectors and professionals. In 2020, a total of 306 hours of training were provided by national SBHP trainers to 31 clinical providers from eight counties in the fields of social work and behavioral health. Three of the 31 providers are now listed on the SBHP searchable, national registry.

The NH Gun Shop Project

The Gun Shop Project began 2020 with an effort to follow up on the mailing to NH Gun Shops by visiting free-standing retailers to further engage them in the project to educate their employees and customers on suicide prevention with a focus on firearms, the leading method for suicide in NH and most other states. This effort was suspended due to COVID-19. The NH

Firearm Safety Coalition (NHFSC), however, has been working on a contract with the American Academy of Pediatrics to develop an online training module for pediatricians and other primary care providers on how to approach the issue of firearm suicide with gun-owning families. This project should be completed in 2021.

Have you found this report to be useful?

Please share your feedback through the survey linked below so that this report can be even better in the future.

<https://www.surveymonkey.com/r/V99FMBP>

2020 Data

SPC/YSPA Data Subcommittee Membership Representation 2020-2021

Injury Prevention Center at CHaD
NAMI New Hampshire (National Alliance on Mental Illness)
New Hampshire Army National Guard
State of New Hampshire Department of Corrections
State of New Hampshire Department of Health and Human Services
State of New Hampshire Office of the Chief Medical Examiner

Introduction

The data presented in this report are the result of collaboration among a variety of organizations and people. The data were compiled by two major groups for suicide prevention in New Hampshire, the YSPA and the SPC. YSPA and SPC merged data efforts, combining historical expertise with emerging methods. YSPA has been collecting and analyzing data about youth and young adult suicide deaths and behavior over the last 20+ years and first created this report format in 2003. The SPC has been analyzing and planning for data capacity improvements for the last 11 years. Key areas of interest and concern for suicidal behavior in NH are included in this report. A data interpretation and chart reading section has been included at the end of the report.

While each suicide is a separate act, only aggregate data is presented in this report. Aggregate data helps inform which populations and age groups are most at risk, reveals points of particular vulnerability, and thus helps guide prevention and intervention efforts, and identify where to direct program funding. It also protects the privacy of individuals and their families. We respectfully acknowledge that the numbers referred to in this report represent lives tragically lost, leaving many behind who are profoundly affected by these deaths.

In previous years this report included death data from two primary sources; Vital Records data (official death records for NH residents) for the State of NH obtained from Health Statistics and Data Management (HSDM), Division of Public Health Services, NH DHHS; and Office of the Chief Medical Examiner (OCME) for the State of NH. As of the 2019 NH Suicide Prevention Annual Report, the primary source of death data has been changed to NH's implementation of the National Violence Death Reporting System (NH-VDRS).

NH DHHS collaborates with the NH Department of Justice (DOJ) on implementation of the NH-VDRS¹ under the auspices of the OCME. The CDC currently includes all fifty states and Puerto Rico in the NVDRS project. NVDRS is a de-identified secure database system used by all US states. NH-VDRS utilizes the system to collect data on violent deaths in NH. Violent deaths include suicides, homicides, firearms accidents, and other violent deaths.

¹ Disclosure: NH-NCDRS funding is from the Centers for Disease Control and Prevention Cooperative Agreement Number 6 NU17CE002610-04-02.

Suicide death demographic data is collected from the NH Division of Vital Records Administration death certificate database on all suicide victims who died in the state of New Hampshire. NH residents who died in other states are included in the NH-NVDRS statistics in the state where they died. NH-VDRS abstracted data comes from Assistant Deputy Medical Examiner (ADME) investigation reports, toxicology, and autopsies reports, all of which are located in the Medical Examiner's office. Another abstracted data resource is law enforcement reports, which include state, local, and sheriff departments.

NH-VDRS is required to report the outcomes of the data on violent deaths as defined by CDC grant requirements. The analysis as provided is focused on direct outcomes and does not engage in policy analysis. Any policy analysis based on the NH-VDRS provided data included in this report was done by the NH Suicide Prevention Council Data Subcommittee.

Additional data sources were used for specific purposes throughout this report that may have varying methods of collection. All of the Tables and Figures in this report include citation for the data source to prevent confusion. Different data sources also vary regarding how quickly the information is made available and how often it is collected/reported. The time periods reported for each source are indicated with the corresponding Table or Figure.

Demographic profile of New Hampshire

Comparing New Hampshire to the US

Tables 1 through 6 below present NH and US demographic characteristics, as well as indicators of substance use and mental health. NH is a small state, with just over 1.3 million residents (US Census, 2020). Overall, NH is relatively homogeneous in terms of race and ethnicity. It has above average ratings for economic factors and education. NH is also above the US average for alcohol and illegal drug use, with the 2nd highest rate in the US for alcohol use in the past month² and the 12th highest rate for marijuana use in the past month (National Survey on Drug Use and Health, 2018-2019).

Table 1


Race/Ethnicity.

	New Hampshire	United States
Race		
White	92.9%	76.3%
Black	1.9%	13.4%
American Indian/Alaskan Native	0.3%	1.3%
Asian	3.1%	5.9%
Native Hawaiian and Other Pacific Islander	< 0.1%	0.2%
Persons Reporting Two or More Races	1.8%	2.8%
Ethnicity		
Persons of Hispanic or Latino Origin	4.2%	18.5%

Data Source: US Census Bureau 2020

² "Past month" refers to the 30 days prior to the administration of the National Survey on Drug Use and Health.

Figure 1
NH and US Race/Ethnicity.



Data Source: US Census Bureau 2020

Table 2

Age.

	New Hampshire	United States
Under 18	1.85%	22.1%
18 to 24	8.9%	9.1%
25 to 44	24.5%	26.8%
45 to 64	28.8%	25.1%
65 to 74	11.6%	9.9%
75 and Up	7.6%	7.0%

Data Source: US Census Bureau 2020

Table 3

Economic Factors.

	New Hampshire	United States
Unemployed Residents	3.0%	4.5%
Persons Below Poverty Level	7.3%	12.3%
Persons Without Health Insurance	6.3%	9.2%
Per Capita Income (Yearly)	\$41,241	\$35,672
Median Household Income	\$77,933	\$65,712
Median Home Value (Owner Occupied)	\$281,400	\$240,500

Data Source: US Census Bureau American Community Survey 2019

Table 4

Education – Individuals Age 25 and Older.

	New Hampshire	United States
Less Than High School Graduate	6.7%	11.4%
High School Graduate or Associates Degree	55.8%	55.5%
Bachelor's Degree or Higher	37.6%	33.1%

Data Source: US Census Bureau American Community Survey 2019

Table 5

Substance Use – Individuals Age 12 and Older.

	New Hampshire	United States
Marijuana Use – Past Month	14.05%	10.80%
Alcohol Use – Past Month	62.54%	50.92%
Tobacco Use – Past Month	19.95%	21.28%

Data Source: National Survey on Drug Use and Health, 2018-2019

Table 6

Mental Health Indicators – Individuals Age 18 and Older.

	New Hampshire	United States
Serious Mental Illness – Past Year	4.85%	4.91%
Major Depressive Episode – Past Year	8.27%	7.51%
Had Thoughts of Suicide – Past Year	4.68%	4.58%


Data Source: National Survey on Drug Use and Health, 2018-2019

The Big Picture: Suicide in NH and Nationally

The Tables and Figures below depict various suicide related data. Some are specific to NH while others compare NH and national statistics.

Figure 2 (pg. 24) presents the suicide rate in NH and the US for the past ten years. The rate in NH has varied from year to year, due to its small size, while the US rate has remained more consistent year to year. Even though the NH rate has varied, until 2014 there had been no statistically significant differences from one year to the next since at least the year 2000. 2010 was the first year in recent history where there was a statistically significant difference compared to any other recent year. The 2010-2012 suicide rates are significantly greater than the rates for 2000, 2002, and 2004. This appears to be consistent with changes in the rates of suicide nationally. In 2014 there was a spike in the NH rate that brought it significantly above the rates prior to 2010. This increase was not seen in other states or for the US as a whole in 2014. The increase starting in 2014 continued through 2018, though not statistically significant from year to year. The rate decreased in 2019, though it was again not a statistically significant change from the preceding year.

Figure 2
Crude Suicide Death Rates per 100,000 in NH by Year 2010-2019.



Data Source: 2010-2019 – CDC WISQARS

Table 7 (pg. 25) displays the 10 leading causes of death for people of different age groups in NH. From 2015-2019, suicide among those aged 10-34 was the second leading cause of death in NH and nationally. Suicide rates for individuals age 10-34 during 2015-2019 were behind only deaths due to unintentional injury. Within that age group, a substantial number of unintentional injuries in NH include motor vehicle crashes and unintentional overdose deaths. Suicide among individuals of all ages was the 8th leading cause of death in NH, and the 10th leading cause of death nationally.

Table 7
10 Leading Causes of Death, New Hampshire, by Age Group, 2015 – 2019.

Rank	Age Groups										All Ages
	<1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65+	
1	Short Gestation 33	Unintentional Injury 10	Malignant Neoplasms ---	Malignant Neoplasms 11	Unintentional Injury 321	Unintentional Injury 818	Unintentional Injury 642	Malignant Neoplasms 866	Malignant Neoplasms 2,622	Heart Disease 11,250	Malignant Neoplasms 13,938
2	Congenital Anomalies 29	Congenital Anomalies ---	Unintentional Injury ---	Suicide 11	Suicide 152	Suicide 182	Malignant Neoplasms 211	Unintentional Injury 608	Heart Disease 1,395	Malignant Neoplasms 10,137	Heart Disease 13,416
3	Maternal Pregnancy Comp. 15	Homicide ---	Homicide ---	Unintentional Injury ---	Heart Disease 20	Malignant Neoplasms 62	Suicide 195	Heart Disease 577	Unintentional Injury 484	Chronic Low. Respiratory Disease 3,136	Unintentional Injury 4,469
4	Placenta Cord Membranes 15	Influenza & Pneumonia ---	Benign Neoplasms ---	Congenital Anomalies ---	Malignant Neoplasms 18	Heart Disease 49	Heart Disease 118	Suicide 272	Chronic Low. Respiratory Disease 397	Alzheimer's Disease 2,267	Chronic Low. Respiratory Disease 3,641
5	Respiratory Distress 10	Malignant Neoplasms ---	Congenital Anomalies ---	Benign Neoplasms ---	Homicide 14	Liver Disease 15	Liver Disease 68	Liver Disease 169	Diabetes Mellitus 277	Cerebro- vascular 2,203	Cerebro- vascular 2,486
6	Circulatory System Disease ---	Diabetes Mellitus ---	Heart Disease ---	Chronic Low. Respiratory Disease ---	Chronic Low. Respiratory Disease ---	Homicide 13	Diabetes Mellitus 31	Diabetes Mellitus 101	Liver Disease 277	Unintentional Injury 1,566	Alzheimer's Disease 2,296
7	Intrauterine Hypoxia ---	Heart Disease ---	Acute Bronchitis ---	Influenza & Pneumonia ---	Congenital Anomalies ---	Cerebro- vascular 11	Homicide 25	Chronic Low. Respiratory Disease 81	Suicide 223	Diabetes Mellitus 1,226	Diabetes Mellitus 1,646
8	Bacterial Sepsis ---	---	Diseases Of Appendix ---	Nephritis ---	Cerebro- vascular ---	Congenital Anomalies ---	Cerebro- vascular 17	Cerebro- vascular 73	Cerebro- vascular 175	Influenza & Pneumonia 1,070	Suicide 1,271
9	Unintentional Injury ---	---	---	Pneumonitis ---	Diabetes Mellitus ---	Diabetes Mellitus ---	Chronic Low. Respiratory Disease 16	Septicemia 36	Nephritis 84	Nephritis 810	Influenza & Pneumonia 1,167
10	SIDS ---	---	---	---	Four Tied ---	Two Tied ---	Congenital Anomalies 11	Congenital Anomalies 24	Septicemia 83	Parkinson's Disease 786	Nephritis 926


Produced By: Office of Statistics and Programming, National Center for Injury Prevention and Control,
 Centers for Disease Control and Prevention

Data Source: National Center for Health Statistics, National Vital Statistics System

---Note: Beginning with 2008 data, the CDC has suppressed state-level counts for categories with fewer than ten deaths

The vast majority of violent deaths in NH are suicides. For every homicide in NH, there are approximately 11 suicides. This ratio is in sharp contrast to national statistics, which show approximately 3 suicides for every homicide. For every suicide death in NH and nationally, there are approximately 4 deaths classified as unintentional injuries (CDC WISQARS, 2015-2019). Overall, suicide constitutes a larger proportion of all traumatic deaths in NH than in the US as a whole. The breakdown of violent deaths³ in NH by gender is presented below in **Figure 3**.

Figure 3
Males die of violent deaths of all manners at rates greater than those for females.



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

The most effective way to compare NH to the US is to look at suicide death rates. **Table 8** (below) presents NH and US suicide death rates by age group.

Table 8
Crude Suicide Death Rates per 100,000 in NH & US, by age group, 2015-2019.

	ALL AGES	YOUTH 10 TO 17	YOUNG ADULTS 18 TO 24	YOUTH AND YOUNG ADULTS 10 TO 24
NH	18.86	4.60	21.06	12.87
US	14.30	4.90	15.66	10.05

	AGES 25 TO 39	AGES 40 TO 59	AGES 60 TO 74	AGES 75 AND OVER
NH	22.61	26.70	18.00	22.40
US	17.19	19.69	16.50	18.69

Data Source: CDC WISQARS

³ Violence deaths include suicide, homicide, and any firearm related death regardless of intent.

Adults age 40 to 59 had the highest suicide rates of all age groups identified above (26.70 NH, 19.69 US) from 2015-2019 in both NH and the US. There is a substantial increase in the rates from youth (ages 10-17) to young adults (ages 18-24) revealing the transition from middle/late adolescence to late adolescence/early adulthood as a particularly vulnerable time for death by suicide.

Youth and Young Adult Suicide in NH

Between 2015 and 2019, 156 NH youth and young adults aged 10-24 have lost their lives to suicide. Males are much more likely to die by suicide in NH (80%) and nationwide (79%). Hanging and firearms were the most frequently used methods in NH among youth and young adults during this period, with firearms being used with a slightly higher frequency. Nationally, a greater proportion of youth and young adults who die by suicide use firearms.

Table 9 (pg. 30) presents the number of youth and young adult deaths by year. This year-by-year data has been plotted in **Figures 4 and 5** (pg. 28). There are a relatively small number of deaths in this age group that can fluctuate from year to year. The rates presented on the chart of deaths over rolling three-year intervals shown in **Figure 49** (pg. 65) helps to smooth out small year to year fluctuations, and also addresses population increases by presenting rates per 100,000.

Positive Outcomes and Testimonials


A student and his mother were sent to a NH emergency department one spring morning for an emergency suicide assessment based on requirements of the School District Suicide Intervention Protocol. The student had expressed suicidal warning signs. The School Resource Officer and a member of the Response Team, both known by the family, joined them at the hospital.

During the process the student's mother shared that her son had been asking for permission to take his father's rifle and go out into the woods near their home. The mother had denied his request and explained her safety concerns to him.

"There was a simultaneous shiver that went through each of us when we registered the great relief of intervening with an emergency assessment before a suicide attempt...especially with such a potentially lethal plan."


The student was able to share his feelings and a comprehensive follow up plan was created. The student and his mother learned about the resources available to help them both.

Figure 4
Suicide Deaths of NH Youth Ages 10-24
2015 - 2019



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 5
While NH male youth suicide deaths increased 2015-2019, female youth rates have remained relatively stable.




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Older Adult Suicide in NH

In light of the rapidly expanding number and proportion of older adults in New Hampshire's population, suicide in older adults is a growing public health concern. Added to the changing demographics is the rising prevalence of mental illness and substance disorders. Untreated mental illness such as depression is a significant risk factor for suicide among all ages, but it is particularly of concern in later life as older adults with depression or other mental health conditions receive treatment at markedly lower rates than the rest of the population.⁴

Another concern is the rate of attempts to completed suicides for older adults. The lethality rate in people over 65 years of age is markedly higher in comparison to other age groups. While there is one death for every 36 attempts in the general population, there is one death for every four attempts in individuals over 65. One related factor is that aged individuals may be physically fraailer than younger individuals and are therefore less likely to survive self-injurious acts. A second is that older adults tend to be more isolated than younger people, making detection or timely intervention less likely. A third factor is the lethality of means; compared to other age groups, adults over 65 are more likely to use firearms as a means of suicide.



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program


⁴ Conwell Y. Suicide and suicide prevention in later life. Focus 2013; 11(1): 39–47.

http://www.aoa.gov/AoA_Programs/HPW/Behavioral/docs2/Suicide%20Prevention%20Webinar%20Presentation%20Slides2.pdf#page=14

Figure 7

The number of male and female older adult deaths are relatively stable from year to year.

**Suicide Deaths by Gender - NH Older Adults
(Ages 65+)
2015 - 2019**



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Suicide Across the Lifespan in NH


Table 9 presents the number of suicide deaths in NH by year, by gender, and selected age groups. These counts include both NH residents and out-of-state residents who died by suicide in NH. When comparing year to year, there is a noticeable increase in the number deaths from 2015 to 2019. The proportion of deaths by gender and age group remained relatively consistent from one period to the next. The number of deaths by year have been plotted in **Figure 8** (pg. 31) and **Figure 9** (pg. 31).

Table 9
NH All Ages Suicide Death Trend, by Gender, Age Group and Method, 2015-2019.

Year	Total	Male	Female	≤ 24	25-64	65+
2015	228	164	64	19	174	35
2016	237	179	58	28	156	53
2017	263	205	58	38	182	43
2018	275	219	56	36	188	51
2019	259	207	52	35	171	53
Total	1262	974	288	156	871	235
Percent of Total	100%	77%	23%	12%	69%	19%


Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 8
NH Suicide Deaths - All
2015 - 2019



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 9
NH Suicide Deaths by Gender - All Ages
2015 - 2019




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 10 (pg. 32) and **Figure 11** (pg. 33), respectively, display NH suicide deaths and suicide death rates for all ages by age groups and gender from 2015-2019. Rates are expressed as the number of suicide deaths per 100,000 people. Displayed together, these charts reveal how death rates correct for differences in the size of each age group. While the highest number of suicide deaths occur in the 45 to 59 year-old age groups, the highest rates, or those at the greatest risk, are males over the age of 85. This is followed by males between the ages of 50 and 54. This second high-risk group is younger than has been seen in past years, where individuals in their 70's generally exhibited higher rates of suicide than individuals in their 40's and 50's.

Figure 10

The highest numbers of suicides deaths are seen in males and females in the 40 and 50-year-old age groups.

**NH Resident Suicide Deaths Counts by Age Group
2015 - 2019**



*Note: Counts/Rates for categories with fewer than six deaths have been suppressed.

Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program


Suicide death rates are also important in determining vulnerable age groups and age-related transitions. The suicide death rate in males rises rapidly from ages 10-14 to 15-19 and then again from ages 15-19 to 20-24, pointing to a rise in vulnerability during the transitions from early adolescence to middle adolescence and then middle adolescence to late adolescence/early adulthood. Similarly, suicide rates among elderly males increase substantially at 85 years compared to the younger age groups, indicating another vulnerable time of life for men. As mentioned above, there has been a recent increase in the suicide rates among individuals between the ages of 45 and 59. This may indicate an additional transition period where individuals are vulnerable.

Quick Facts/Talking Points

- Males in NH die by suicide at a rate that is four times the rate for females (CDC WISQARS, 2019).
- Although males are more likely than females to die by suicide, females report attempting suicide at nearly twice the rate of males (NH YRBS, 2019)
- Suicides are preventable (WHO. Suicide prevention: a global imperative. Geneva, Switzerland: WHO Press; 2014).

Figure 11

NH Resident Suicide Deaths Rates by Age Group
(Rates per 100,000)
2015 - 2019



*Note: Counts/Rates for categories with fewer than six deaths have been suppressed.

Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program


Geographic Distribution of Suicide in NH

The numbers and rates of suicide in NH are not evenly distributed throughout the state. **Figure 12** (pg. 34) shows youth and young adult suicide rates by county in NH. **Figure 13** (pg. 34) presents this data for NH residents of all ages⁵. The county suicide death rate chart indicates geographical locations that may be particularly vulnerable to suicide (youth and young adult and/or all ages). Due to small numbers, most of these differences are not statistically significant. However, the all ages rate (**Figure 13** – pg. 34) for Rockingham County (all ages rate: 14.2 per 100,000) is significantly lower than the all ages suicide rates for Carroll County (all ages rate: 20.8 per 100,000) and Merrimack County (all ages rate: 20.1 per 100,000). It is also significantly below the overall NH rates (all ages rate: 17.5 per 100,000). The rates for Carroll County (all ages rate: 21.8 per 100,000), Cheshire County (all ages rate: 19.2 per 100,000) Coos County (all ages rate: 21.9 per 100,000), Hillsborough County (all ages rate: 17.9 per 100,000), Merrimack County (all ages rate: 20.1 per 100,000), and Sullivan County (all ages rate: 20.9 per 100,000) were significantly higher than the all ages US rate (all ages rate: 14.3 per 100,000). County limits are neither soundproof nor absolute. A suicide that occurs in one county can have a strong effect on neighboring counties, as well as across the state, due to the mobility of residents. **Figure 14** (pg. 35) presents the suicide rates for all ages from 2015 to 2019 as a NH map broken down by county.

⁵ County level analyses exclude any cases where location data were unknown or otherwise not available. Eight cases were excluded due to missing data for 2015-2019.

Figure 12


**NH Youth Suicide Crude Death Rates by County - Ages 10 - 24
2015 - 2019**



NH Rate Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program
US Rate Data Source: CDC WISQARS

Figure 13

**NH Resident Suicide Crude Death Rates by County - All Ages
2015 - 2019**



NH Rate Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program
US Rate Data Source: CDC WISQARS

Figure 14




Table 10 (below) further expands upon this county breakdown by presenting the percent of suicide deaths in each county by gender. In the majority of counties, the ratio is five male deaths for every one female death. The exceptions to this include Belknap, Hillsborough, Merrimack, and Rockingham Counties where the ratio is approximately three male deaths for every one female death. The ratio of males and females residing in each county is approximately one-to-one statewide.

Table 10
2015 - 2019
Percent of Suicide Death by Gender in NH Counties

	Percent of Female Deaths in County	Percent of Male Deaths in County	Male to Female Suicide Death Ratio in County	Percent of Female County Residents	Percent of Male County Residents	Male to Female Population Ratio
Belknap County	26%	74%	3:1	51%	49%	1:1
Carroll County	17%	83%	5:1	50%	50%	1:1
Cheshire County	18%	82%	5:1	51%	49%	1:1
Coos County	17%	83%	5:1	48%	52%	1:1
Grafton County	18%	82%	5:1	51%	49%	1:1
Hillsborough County	26%	74%	3:1	50%	50%	1:1
Merrimack County	28%	72%	3:1	51%	49%	1:1
Rockingham County	22%	78%	3:1	50%	50%	1:1
Strafford County	18%	82%	5:1	51%	49%	1:1
Sullivan County	18%	82%	5:1	51%	49%	1:1

Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Suicide Behavior in NH: Gender Differences – Attempts, Deaths, and Risk Factors


Youth and Gender

While males represent 80% of the youth and young adult suicides from 2015-2019, the fact that males *die* by suicide at a higher rate than females may largely be due to males using more lethal means. See **Figures 15** (pg. 37) and **16** (pg. 37). In fact, females *attempt* suicide at a higher rate than males. When examining how many NH youth and young adults ages 10-24 were hospitalized and then discharged for self-inflicted injuries in 2015-2019, it is shown that 67% of the 799 inpatient discharges represent females, while only 33% represent males. Likewise, the 2019 NH Youth Risk Behavior Survey (YRBS) reports approximately 1.6 times as many female youth attempt suicide as males each year (8.4 % of females and 5.3% of males). Emergency department (ED/ambulatory) data reveals a similar gender ratio, based on self-inflicted injury rates.⁶

⁶ Classifying an injury as self-inflicted is another way of stating that the injury was an instance of deliberate self-harm. Not all self-inflicted injuries necessarily represent suicide attempts. However, analysis of these injuries is the best currently available proxy for estimating suicide attempts.

Figure 15
Four times as many male NH residents ages 10-24 died by suicide 2015-2019.


**NH Resident Suicide Deaths by Gender
- Ages 10-24
2015 - 2019**



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 16
For NH residents of all ages, three times as many males died by suicide than females during 2015-2019.

**NH Resident Suicide Deaths by Gender
- All Ages
2015 - 2019**



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program


Female youth are less likely to die by suicide, possibly resulting from less severe injuries during suicide attempts (self-inflicted injuries). However, female youth do attempt suicide more frequently than males – 1.2-2.0 times as often (**Figure 17** – below, **Figure 18** and **Figure 19** – pg. 39). This report refers to three types of data; Emergency Department Discharges, Inpatient Discharges, and individuals treated/transported by Emergency Medical Services (EMS).

Emergency Department (ED) data includes patients who came to the ED and stayed at the hospital for less than 24 hours (also called Ambulatory Discharges). Inpatient data refers to patients who were admitted to the hospital for more than 24 hours. If a patient goes to an ED and is admitted for inpatient services, they are removed from the count in the ED data and listed as inpatients. The hospital discharge data records the number of hospital visits, not the number of individual persons who went to the hospital for care. For example, if one patient went to the hospital three different times over the course of a year it would be counted as the same number of visits as three different patients who went to the hospital one time each over the course of one calendar year.

The EMS data presents the number of times individuals were treated and/or transported by an EMS provider where the individual had some type of self-inflicted injury. As with the hospital data, the EMS data looks at the number of visits/incidents, not unique individuals. The EMS data comes from a different source than the hospital data. Therefore, the cases are not de-duplicated between the two datasets (i.e., an individual may be counted in the hospital and EMS datasets for the same incident). The cases included in the EMS dataset are ones where the intent of the injury was listed as “self-inflicted”. This does not include incidents where an injury was deemed to be accidental. The NH Department of Safety, Division of Fire Standards and Training and Emergency Medical Services reports that pre-hospital care (EMS) was significantly impacted by COVID-19 in 2020. Self-harm/Suicide attempt calls were down approximately 8%, while the total number of incidents was down approximately 15%.

Figure 17
A greater percentage of female than male NH residents attempted suicide, as seen in inpatient and emergency department discharges related to self-inflicted injuries 2015-2019.

NH Inpatient Discharges for Self-Inflicted Injuries by Gender - Ages 10 - 24
2015 - 2019



NH Emergency Department Discharges for Self-Inflicted Injuries by Gender - Ages 10 - 24
2015 - 2019






Figure 18

**NH Inpatient Discharges for Self-Inflicted Injuries by Gender - All Ages
2015 - 2019**




**NH Emergency Department Discharges for Self-Inflicted Injuries by Gender - All Ages
2015 - 2019**




Data Source: NH Hospital Discharge Data by the NH DHHS Injury Prevention Program

Figure 19

**Self-Harm by Gender
All Ages, 2020**



**Self-Harm by Gender
Ages 0-24, 2020**



Data Source: New Hampshire Department of Safety, Division of Fire Standards and Training and Emergency Medical Services


History of Suicide Attempts and Intent Disclosure

The vast majority of individuals who died by suicide in NH have no reported history of prior suicide attempts or disclosure of suicidal intent. Females who died by suicide in NH were approximately twice as likely as males to be known to have a prior history of suicide attempts. Females who died by suicide in NH were also three times as likely as males to have previously disclosed their suicidal intent (**Figure 20** – below).

Disclosure of suicidal intent and prior suicide attempt(s) are significant risk factors for suicide. If you are concerned about an individual with these or other risk factors, connect them with appropriate resources such as the National Suicide Prevention Lifeline – 1-800-273-8255 or a local mental health professional. **If you are concerned that there is imminent risk, call 911.**

Figure 20

**Suicide Deaths in NH - History of Suicide and Suicide Intent Disclosure by Sex
2015 - 2019**




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

NH Behavioral Risk Factor Surveillance System (BRFSS)

The Behavioral Risk Factor Surveillance System (BRFSS) is a survey conducted with a representative sample of state residents. The survey includes the question “Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?” Although this is not a perfect proxy measure for depression, it gives one a general sense of the percentage of NH residents that may be experiencing a depressed mood. The results from this item are included in **Figure 21** (below).

Figure 21
NH BRFSS – Number of Days Mental Health Was Not Good - NH Residents Age 18 and Over.



Data Source: BRFSS data prepared by NH DHHS BPHSI


Gender differences exist not only for suicide attempts and deaths, but also for help-seeking behavior. A 2018 CDC report indicated that approximately half of individuals who take their own life had a mental health condition; the most common diagnoses being depression, anxiety, and substance use disorders⁷. Yet a much smaller percentage were receiving treatment. In NH, over 43,600 people received treatment at one of the state’s ten Community Mental Health Centers (CMHC)⁸ each year. In 2020, this works out to approximately 1 out of every 31

⁷ Stone DM, Simon TR, Fowler KA, et al. Vital Signs: Trends in State Suicide Rates — United States, 1999–2016 and Circumstances Contributing to Suicide — 27 States, 2015. MMWR Morbidity Mortality Weekly Report 2018; 67:617–624. DOI: <http://dx.doi.org/10.15585/mmwr.mm6722a1>.

⁸ Community Mental Health Centers are private not-for-profit agencies that have contracted with the NH Department of Health and Human Services, Bureau of Behavioral Health, to provide publicly funded mental health services to individuals and families who meet certain criteria for services. More information on the centers is available from <http://www.dhhs.state.nh.us/dcbcs/bbh/centers.htm>

residents in the state. Of those individuals in treatment, approximately 55% of them were female and 45% were male. This is illustrated in **Figure 22** (below). Without additional data it is not possible to say how these numbers relate to the connection between these treatment figures and the greater number of suicide deaths among males and/or the greater number of suicide attempts reported among females.

Figure 22
Individuals receiving treatment at NH Community Mental Health Centers presented by age and gender.



Data Source: NH Bureau of Behavioral Health

Patients that cannot be treated in an outpatient setting, such as involuntary admissions due to potential suicide risk, will generally be admitted to New Hampshire Hospital, the NH state psychiatric hospital. In an average year there are approximately 1,315 admissions to New Hampshire Hospital (estimates based on New Hampshire Hospital admissions for fiscal years 2016 - 2020⁹). **Figure 23** (pg. 43) presents the total number of admissions per bed at New Hampshire Hospital.

Positive Outcomes and Testimonials


Suicide is preventable with the understanding we all must embrace: “treatment works”.

Support and early intervention is everyone’s job, as saving a life makes a world of difference for so many.

Maggie Pritchard
Executive Director, Lakes Region Mental Health
Former Vice-Chair, NH Suicide Prevention Council

⁹ The NH State Fiscal Year runs from July 1st of one calendar year through June 30th of the following calendar year (e.g., fiscal year 2020 ran from July 1st 2019 through June 30th 2020).

Figure 23
**New Hampshire Hospital: Number of Admissions
Per Hospital Bed By Fiscal Year**




Data Source: New Hampshire Hospital

Mental Health and Suicides in NH

Among the various risk factors for suicide in NH, depression and depressed mood¹⁰ figure prominently. From 2015 to 2019, over half of all individuals who died by suicide in NH were reported to have a depressed mood around the time of death. A slightly greater proportion of females than males were reported to have been experiencing a depressed mood (Figure 24 – below)

Figure 24
**NH Suicide Deaths by Sex and Depressed Mood
2015 - 2019**




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

¹⁰ Depressed mood is a field tracked in the NH-VDRS. Based on CDC criteria, depressed mood does not require a clinical diagnosis, and does not need to have been identified as a factor directly contributing to the death.

Figure 25 (below) addresses mental health diagnoses of individuals who died by suicide, where this information was available. The mental health diagnoses are based on evidence at the scene such as medications prescribed to the deceased, information confirming that the individual had a mental health provider (psychiatrist, mental health counselor, etc.), and reports from next of kin. A challenge with reports from next of kin is that they may not have up-to-date knowledge on their loved one's mental health treatment and condition. As a result, there are many suicide deaths where there is no data available related to mental health diagnosis. The availability of mental health diagnosis information in the NH-VDRS continues to improve as death scene investigators expand their documentation of mental health issues. The categories in the graph below are based on counts in a data field that represents the primary mental health diagnosis. Some decedents may have had more than one mental health diagnosis.

Figure 25
Suicide Deaths in NH by Mental Health Diagnosis
2015 - 2019



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program


Age, Gender and Self-inflicted Injury

When the rates for NH resident inpatient hospitalizations/discharges and emergency department use for self-inflicted injuries from 2015-2019 are examined by gender and age group, the variability can be seen (**Figures 26** and **27** – pgs. 45-46). As above, these data refer to number of visits; therefore, individuals may be counted multiple times if they were admitted or seen more than once during the year.

Female NH residents have a higher overall rate of inpatient hospitalizations/discharges for self-inflicted injuries until the ages 85+ where the male rate exceeds the rate for females. For females aged 15-24, the rate of those being discharged from inpatient care (**Figure 26** – pg. 45) is 223/100,000, nearly four times the rate for males of the same age. The peak age for females and males is between 24 and 34 for self-inflicted injuries requiring hospitalizations. Again, ED usage rates, depicted in **Figure 27** (pg. 46), point to females aged 15-24 as a population particularly


vulnerable to self-injury and/or suicide attempts, with females in this group exhibiting a rate over 542/100,000, about 87 times the suicide death rate for this population. Males also peak in self-injury around this age group with the male rates for ages 15 to 24 being over 272/100,000. Although male rates peak around this age group, their rates are much lower than those for females. Also of note, the total number of youth and young adult (ages 15-24) ED visits (3,568) is 5 times greater than the number of inpatient discharges for this population. Because less severe injuries are more common among self-inflicted youth injuries, there are many more attempts than deaths. This data reinforces that the transition from middle adolescence to late adolescence/early adulthood is a time of great risk for suicidal thinking, self-harm and suicide attempts. EMS data (**Figure 28 – pg. 46**), which includes individuals treated and/or transported by Emergency Medical Services for a self-inflicted injury, presents a similar picture to the hospital data in terms of high-risk age groups. Females age 15 to 24 present the highest rates of self-inflicted injuries. In most other age groups, male rates exceed the rates for females in the EMS data.

Figure 26
NH female residents ages 15-24 and 25-34 show the highest rates of suicide attempts, higher than males of any age group.




Data Source: NH Hospital Discharge Data prepared by the NH DHHS Injury Prevention Program

Figure 27
NH female residents ages 15-24 show the highest rates of suicide attempts, with male rates also peaking at this age.



Data Source: NH Hospital Discharge Data prepared by the NH DHHS Injury Prevention Program

Figure 28
NH female residents ages 15-24 show the highest rates of suicide attempts followed closely by males and females in the 25-34 age group.



Data Source: New Hampshire Department of Safety, Division of Fire Standards and Training and Emergency Medical Services

According to inpatient and emergency department (ED) discharge data across all ages in NH, there are approximately 10 suicide attempts for every suicide death. This number does not include attempts that go unreported, unrecognized, or without a hospital or ED visit which required medical intervention. Further, the rates of attempts for young people and females create an even greater ratio of suicide attempts to deaths. Based solely on hospital and emergency department self-injury data, it can be estimated that over 1,080 youth and young adults (age 24 and under) attempt suicide each year in NH.

In contrast to the above data, which are based on cases where medical intervention is required, the results of the YRBS presents data collected from high school aged youth by self-report. In 2019, nearly 7 percent of high school students completing the YRBS reported having attempted suicide at least one time over the previous year. Based on the YRBS figures, this works out to over 3,800 high school age youth in NH who may attempt suicide each year. The YRBS reports may account for attempts not included in hospital self-injury data. This could be the case for any attempts with relatively non-lethal means where medical assistance was not sought. Of particular concern for this data is the likelihood that in many of these cases, the youth have never sought help or disclosed the attempt to any adult. It is also possible that self-reports exaggerate the incidence of suicide attempts among high school age youth.

While the great majority of self-inflicted injuries¹¹ are not fatal, because of the larger incidence they may directly and indirectly affect a greater number of people than fatalities. A significant risk factor for suicide is a previous attempt: in one study 21-33% of people who die by suicide have made a previous attempt (Shaffer & Gould, 1987). Therefore, any suicide attempt, regardless of its lethality, must be taken seriously. If not addressed, it could be followed by additional attempts. Therefore, once an individual has made an attempt, secondary prevention is necessary.

Positive Outcomes and Testimonials


Following a SAMHSA grant awarded in 2018, the NH State Police have begun implementing the Crisis Intervention Team (CIT) Program coordinated by NAMI NH. This training has resulted in Troopers being able to identify individuals experiencing a mental health crisis and/or individuals at risk for suicide, defuse situations, and direct individuals to mental health services where they might have otherwise been arrested.

¹¹ Classifying an injury as self-inflicted is another way of stating that the injury was an instance of deliberate self-harm. Not all self-inflicted injuries necessarily represent suicide attempts. Analysis of these injuries, however, is the best currently available proxy for approximating suicide attempts.

Additional Demographic Characteristics of Individuals in NH Who Died by Suicide

Additional demographic factors may play a role in suicide. **Figure 29** (below) presents the marital status of individuals who died by suicide in NH between 2015 and 2019. The data in **Figure 29** differs substantially from the overall breakdown by marital status for NH with fewer married individuals and more divorced individuals dying by suicide. In the NH population, approximately 50% of individuals are married and approximately 12% are divorced, while only 30% of individuals who died by suicide in NH were married and over 22% were divorced.

Figure 29
Suicides in NH by Sex and Marital Status
2015 - 2019




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Educational attainment may also play a role in suicide. The prevalence of suicides in NH is greatest among individuals who had educational levels of high school or less than high school (**Figure 30 – pg. 49**) and substantially lower among individuals with college degrees. Among adults in NH, over 37% have a bachelor's degree or higher (**Table 4 – pg. 23**), while only 18% of male and 22% of female suicide deaths in NH are by individuals with an equivalent educational level. Nationally, higher levels of education are generally correlated with higher income and lower levels of unemployment¹². The larger number of suicide deaths among individuals with education levels of high school or less could indicate a greater prevalence of employment or financial stressors among this group. Job and financial stressors were frequently identified among individuals in NH who died by suicide (**Figure 43 – pg. 58**).

¹² <https://www.bls.gov/careeroutlook/2021/data-on-display/education-pays.htm>


Figure 30
**Suicide Deaths in NH by Sex
and Educational Levels**
2015 - 2019



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Prior to the implementation of NH-VDRS, data on the sexual orientation of individuals who died in NH was not readily available. As seen in **Figure 31** (below), the data related to this is still limited for individuals who died by suicide, with sexual orientation being unknown in over 40% of cases. The large number of cases where this information is unknown may be due to a number of factors, including the stringent NH-VDRS requirements for reporting sexual orientation, and stigma around revealing sexual orientation.

Figure 31
Suicide Deaths in NH by Sexual Orientation
2015-2019



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Suicide Risk While Incarcerated - NH Department of Corrections

The New Hampshire Department of Corrections reported that there have been no deaths by suicide in any of their facilities during 2019. There are approximately 2,500 residents (includes all facilities), with 385 individuals admitted to Department of Corrections facilities in 2019. Every resident (includes new or returning) receives a comprehensive behavioral health screening which includes an assessment for risk of suicide. Individuals assessed as being at risk for suicide are placed on a 24-hour observation level that includes continuing assessment by mental health professionals. After they are discharged from this level of care, they receive appropriate follow up services for a time-period based on their individual needs.

Ongoing training includes quarterly suicide prevention trainings conducted for the corrections officers on the Special Housing Unit (SHU), the Reception and Diagnostics Unit (R & D), and the Secure Psychiatric Unit. All new Department of Corrections employees, both uniformed and non-uniformed, receive four hours of suicide prevention and mental health training as part of the comprehensive orientation program and security officers newly employed at the Secure Psychiatric Unit receive an additional 16 hours of training on behavioral health issues.

Suicide in NH: Methods

The gender difference in suicide deaths/attempts may be explained in part by the fact that males, in general, use more lethal means. Of NH male youth and young adults who died by suicide between 2015 and 2019, 57% used firearms compared to 10% of females (**Figure 33 – pg. 51**). This gender disparity in firearm use persists as residents enter their late 20's, 30's, and 40's with the proportion of male and female deaths from firearms decreasing equally for both genders. The proportion of firearm deaths increases sharply at age 65 for males, with 68% of the suicide deaths in that age group involving a firearm. In NH, the vast majority of all deaths involving a firearm are suicide. This can be seen in the **Figure 32** below.

Figure 32
In 2019, approximately 89% of all NH deaths involving a firearm were suicides.




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Suicide attempt methods have varying lethality. **Figure 34** (pg. 52) compares firearms, hanging, poisoning, and cutting/piercing in terms of the percentage of various outcomes (emergency department visit, inpatient admission, or death) for each method. Over 80% of self-injuries using a firearm result in death. Among youth and young adults, suicide is often a highly impulsive act and poor impulse control is one of the risk factors for suicide. Therefore, intervention efforts that reduce access to firearms and other highly lethal means may be effective to reduce suicide among those at risk for suicide, particularly for those who are more likely to be impulsive. Firearms remain the most used method of suicide throughout the lifespan in NH. **Figure 35** (pg. 52) indicates that self-inflicted drug overdoses/poisonings are treated/transported by EMS at several times the rate of most other mechanisms. The one exception being self-inflicted cutting/piercing injuries which are treated/transported at a substantially higher rate than other mechanisms. Hospital data (**Figure 37** – pg. 53) does not show this same proportion of cut/pierce injuries indicating that EMS providers may treat self-inflicted cut/pierce injuries without the need to transport the individual to a hospital, or that individuals are more likely to contact EMS for a cut/pierce injury and be transported to a hospital by other means for things such as a poisoning. It may also indicate that EMS providers are more likely to report that a cut/pierce injury as being self-inflicted than they are with other injury types. The use of suffocation as a suicide method peaks in early adolescence and decreases steadily throughout the lifespan (**Figure 33** – below).


Figure 33

**Methods Used in Suicide Deaths by Gender and Age Group
2015 - 2019**




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 34



Data Source: NH-VDRS (deaths) and NH Hospital Discharge (emergency and inpatient visits)
Data prepared by the NH DHHS Injury Prevention Program


Figure 35
Percent of method of self-inflicted injuries treated/transported by EMS from 2020.



*This chart is based on the field "Trauma Mechanism of Injury". This field is not available for all incidents. The field may also include a response of "Not Recorded" or "Not Applicable". For the purpose of this report, only incidents with a reported mechanism of injury were included above (approximately 36% of all incidents in 2020).


Data Source: New Hampshire Department of Safety, Division of Fire Standards and Training and Emergency Medical Services

Figure 36
Suicide methods used in NH vary by age group, as seen in 2015-2019.



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program


Figure 37
Poisoning is the most frequent method of suicide attempt,
as seen in hospital discharge data 2015-2019.



Data Source: NH-VDRS (deaths) and NH Hospital Discharge (emergency and inpatient visits)
Data prepared by the NH DHHS Injury Prevention Program

Although suicide attempts employing poison do not account for as many deaths in NH as firearms or hangings, intentional poisonings account for the overwhelming majority of inpatient and ED admissions for suicide attempts (**Figure 34** – pg. 52). **Figure 38** (below) depicts the prevalence of the most common substances used in suspected suicide attempts and self-harm-related exposures in NH as collected by the NNEPC. The top two substances in 2020 were antidepressants and non-opioid analgesics without sedatives (e.g., aspirin or Tylenol).¹³ A recent trend noted by the NNEPC is an increase in the use of cardiovascular medications which can have severe clinical effects. Based on the NNEPC Annual Report covering July 1, 2018 – June 30, 2019, Poison Center staff responded to 8,677 NH human exposure cases during that one-year period. Of those cases approximately 16% were identified as suicide attempts.

Figure 38
Antidepressants have been the top substance used in suspected NH suicide attempts from 2016-2020.
NH Substances Used in Suspected Suicide Attempts
NNEPC Substance Abuse Surveillance and Reporting System, 2016-2020



Data Source: Northern New England Poison Center


Increasing Accidental Poisoning and Drug-Related Death Rates – Cause for Concern

As seen in **Figure 39** (pg. 55), the accidental poisoning and drug-related death rates in NH and the US as a whole have steadily increased from 2010 to 2019. During this time the US rate has

¹³ The suspected suicide attempt cases presented were determined by self-report or the report of an individual acting on behalf of the patient (e.g., a health care professional), or a NNEPC staff assessment. For more information on the NNEPC Annual Report, contact Colin Smith - SMITHC12@mmc.org.

increased by approximately 73% while the NH rate has increased more than 145%. Although it is not possible to determine an exact number, it is likely that these accidental poisoning and drug-related deaths include suicide deaths where there was not enough evidence for the Medical Examiner to classify them as such. This trend is a cause for concern as both a potential increase in poisoning and drug-related suicide deaths, and as a potential indicator of increased risk-taking behavior.

Figure 39
Poisoning/Drug-related death rates in NH increase by more than 145% from 2010 to 2019.




Data Source: CDC WISQARS

Alcohol and Drug Use and Suicide

Alcohol was found to be present in 27% of all NH suicide deaths from 2015 to 2019. Alcohol was found in a greater percentage of male deaths (32% of deaths) than female deaths (16% of deaths). When looking at the presence of alcohol by cause of death (**Figure 40** – pg. 56), it was found to most often be present in firearm deaths for males (55% of male firearm deaths) and drug/poisoning deaths for females (49% of female drug/poisoning deaths). Alcohol is often not the only substance used by individuals who die by suicide in NH. In cases where alcohol and/or drugs were detected, approximately 18% of cases had both alcohol and one or more drugs present (**Figure 41** – pg. 56).

Figure 40


**Suicide Deaths in NH with Alcohol as a Risk Factor by Sex and Cause of Death
2015 - 2019**



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 41


**Suicide Deaths in NH
Alcohol Present with Drugs
2015-2019**



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

The results of toxicological reports include testing various specimens from suicide victims, at various points of the investigation or autopsy. **Figure 42** (pg. 57) depicts the categories of the most commonly found substances from toxicology reports. The most frequently detected substances were benzodiazepines and antidepressants among females, and alcohol and marijuana among males. The figure is based on a total count of the number of times a substance was found in a positive test. Some decedents tested positive for multiple substances and are therefore counted in multiple categories. Individuals who tested positive in this compilation of substance(s) used may or may not have died of such substance(s). Cause of death is presented in **Figure 33** (pg. 51).

Figure 42
**Top Five Substances with a Positive Test Result
in NH Suicide Deaths by Sex**
2015 - 2019




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Co-Occurring Factors and Suicide

Suicide is most often the result of a number of co-occurring risk factors. **Figure 43** (pg. 58) identifies three risk factors that are tracked in the NH-VDRS – problems with alcohol, employment, and finances¹⁴. Approximately 40% of individuals who died by suicide from 2015 to 2019 experienced at least one of those three risk factors. When looking at the intersection of those risk factors, nearly 67% of individuals who were experiencing job problems around the time of death were also experiencing financial problems. It is not surprising to find such a strong correlation between employment and finances as one can have a direct relationship on the other. However, as these risk factors compound, individuals may feel that they are under ever increasing levels of stress.

¹⁴ The NH-VDRS does not have access to detailed data on financial hardship or other such documents, only observations made during field investigations were compiled in ADME reports. Financial hardship may include any one of the following: loss of income, foreclosure on estate/business or, loss of business. NH-VDRS also does not have access to detailed employment data. Job troubles were documented based on OCME field investigations, death certificate statements, by funeral home directors, and declarations by next of kin.

Figure 43
**Suicide Deaths in NH by Sex and
Co-Occurring Factors**
2015 - 2019




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Suicide Notes

In just over 37% of NH suicide deaths from 2015 to 2019, individuals left some form of note behind (**Figure 44 – below**). Females being more likely to have left a note (46% of female deaths) than males (35% of male deaths). These notes vary in format, content, and intent. Individuals may leave instructions for their loved ones on how to resolve financial, estate, burial, and other affairs; complaints/obstacles that they faced; or planning/details that the deceased went through leading up to the death. For the individuals left behind after a suicide death, a note will rarely ever give a satisfactory answer to why their loved one died by suicide.

Figure 44
**Suicide Deaths in NH, Found Note Left
by Sex 2015 - 2019**




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Linking At-Risk Individuals with Help

Crisis lines, such as the National Suicide Prevention Lifeline (NSPL) are vital to suicide prevention efforts in NH and nationally. Nationally, the NSPL receives approximately 2 million calls per year. In 2020, nearly 6,000 of those calls, or roughly 500 per month, were received by the NH NSPL call center (see **Figure 45** – below). These calls indicate that individuals in the state who are at risk for suicide are reaching out for help. The large volume of calls may also indicate decreased stigma around help seeking for mental health and/or suicide.



Figure 45
NH NSPL call center responded to an average of 500 calls per month in 2020.



Data Source: National Suicide Prevention Lifeline

Positive Outcomes and Testimonials

We received a call on our general mailbox from a woman that was directed to our Hotline Manager. The woman stated that she had been calling the Hotline for a long time whenever she is feeling down and everyone has been so great and supportive. "They make me feel better. I have just gotten to the point where I can't even reach out to my church pastor. I called tonight and was feeling very low and I don't remember who I talked to, but she was wonderful. Keep doing what you're doing, you train your people well. You people do good work. Thank you for what you do."

We also recently received a call from a gentleman who had just gotten off the phone with one of our hotline counselors who wanted to share his experience. He said he had called that morning because he was contemplating taking his life and that the Hotline counselor he spoke to had a "very calming and caring voice and listened to me for several minutes, with no judgement whatsoever. He then gave me some places I could call to get further help." He stated that our counselor gave him hope for the day.

Two examples of the feedback received by Headrest, the only accredited hotline in NH that receives calls placed to the National Suicide Prevention Hotline. Headrest receives the majority of calls originating from individuals in NH.

Costs of Suicide and Suicidal Behavior

There were between 35,538 and 47,685 years of potential life lost¹⁵ to suicide from 2015-2019 in NH (CDC WISQARS, 2019). The most obvious cost of suicide is the loss of individuals and their potential contribution to their loved ones and to society. For each suicide death, there are many survivors of suicide loss (the family and close friends of someone who died by suicide) who are then at higher risk for depression and suicide themselves. In addition, many others are affected, including those who provide emergency care to the victims and others who feel they should have seen the warning signs and prevented the death.

Nationally, suicide attempts treated in emergency departments and hospitals represented an estimated \$3.9 billion in health care costs in 2010. This does not include the costs associated with mental health services on an inpatient or outpatient basis (CDC WISQARS, 2019). In NH, suicide deaths where the individual received treatment in a hospital or emergency department and subsequently died resulted in an estimated \$500,000 in medical expenses in 2010 (CDC WISQARS, 2019). Harder to measure is the cost to employers of lower or lost productivity due to suicide attempts or deaths by employees or their loved ones. An estimate provided by the CDC indicates that there is an average work loss cost of \$1.1 million for each suicide death in NH (CDC WISQARS, 2019).

¹⁵ Years of potential life lost (YPLL) is a measure of the extent of premature mortality in a population. This estimate is based on the approximate age at death as well as the number of people who died in that age group in a given year.

Military and Veterans

The NH National Guard


From 2016 through 2020 the NH Army National Guard recorded a total of 44 suicide related incidents of varying levels of severity (ideation, plan in place, attempt, or death), with the majority being ideation or having a plan in place. Of these incidents, 11% were from individuals under the age of 22, 30% were age 22-26, 11% were age 27-31, 7% were age 32-36, 18% were age 37-41, and 11% were ages 42-46. The remaining 11% were age 47 and above (total may not equal 100% due to rounding). Thirty-nine percent of the incidents were by non-deployed personnel, veterans, or dependents of National Guard personnel. Of the incidents recorded, 98% were by males and 2% were by females (males may be disproportionately represented among the NH National Guard compared with the general population).

NH Veterans Served by the Veterans Administration (VA)

The VA provides care to many of the Veterans in the State of NH. During the 2020 Federal Fiscal Year (October 1, 2019 – September 30, 2020), the VA provided care to 24,776 individuals in NH. The percentage of these individuals treated for depression, post-traumatic stress disorder (PTSD), traumatic brain injuries (TBI), and substance use disorder is presented in **Figure 46** below.

Figure 46

Percentage of NH Veterans treated at the VA with depression, PTSD, TBI, or substance abuse as their primary or secondary diagnosis Federal Fiscal Year 2020



Data Source: Veterans Administration


Suicide among Veterans in New Hampshire¹⁶:

Of the individuals who died by suicide in NH from 2015 to 2019, 17% were identified as having current or prior military service (**Figure 47 and Table 11 – below**). The use of the term military service is for all those who served in the armed services of the United States or are still serving. The data sources available to NH-VDRS do not distinguish between individuals who are currently active and those who have been discharged. Veterans made up approximately 8% of the NH population as of 2018¹⁷. With veterans accounting for 17% of the individuals who died by suicide in the state, this may indicate a high-risk group dying at a greater than expected rate.

Figure 47

Suicide Deaths in NH by Sex and Military Status

2015 - 2019



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Table 11
Suicide Deaths in NH by Sex and Military Status
2015 - 2019

NH Suicides - Military Status	Male (Count)	Male (percent)	Female (Count)	Female (percent)
Served in the US Armed Forces	206	21.15%	*	1.39%
Did not serve in the US Armed Forces	756	77.62%	282	97.92%
US Armed Forces Service Unknown	12	1.23%	*	0.69%

Note:

*Counts/Rates for categories with fewer than six deaths have been suppressed.

Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

¹⁶NH-VDRS collects data on veterans only from standard surveillance data sources. The data collection is based on medical examiner data, death certificates, and law enforcement reports. There is no data used that is sourced from any branch of the military.

¹⁷ Veteran population data by state available from https://www.va.gov/vetdata/veteran_population.asp

Military Service and Cause of Death:

Individuals in NH who die by suicide that have served in the military are substantially more likely to use a firearm than civilians (Figure 48 – below, and Table 12 – pg. 64). This difference is evident in males with 52% of individuals with no military service using firearm compared with 71% of males with military service using a firearm. The difference is even more significant among females with military service. Among females with no military service just 23% used firearms, while 75% of females with military service used a firearm.

Figure 48

Suicide Deaths in NH of Individuals Who Served in the US Armed Forces by Sex and Cause of Death 2015 - 2019




Table 12
**Suicide Deaths in NH of Individuals Who
Served in the US Armed Forces by Sex and Cause of Death
2015 - 2019**

NH Suicides of Individuals Who Served in the US Armed Forces - Cause of Death	Lethal Means	Male (Count)	Male (percent)	Female (Count)	Female (percent)
Military Service	Cutting / Piercing	*	2.43%	*	25.00%
	Drugs / Poisoning	22	10.68%	-	0.00%
	Firearms	147	71.36%	*	75.00%
	Hanging / Asphyxiation	30	14.56%	-	0.00%
	Other	*	0.97%	-	0.00%
No Military Service	Cutting/ Piercing	17	2.21%	*	0.35%
	Drugs/ Poisoning	89	11.59%	115	40.49%
	Firearms	399	51.95%	66	23.24%
	Hanging/ Asphyxiation	243	31.64%	86	30.28%
	Other	20	2.60%	16	5.63%

Note:

*Counts/Rates for categories with fewer than six deaths have been suppressed.

-Indicates a category with no cases to report.


Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Suicide Rates in NH

Until 2010, data had indicated that rates of youth and young adult suicide and suicidality overall in NH were flat or on a downward trend. It is nearly impossible to firmly establish causality for such trends. Statewide collaborative prevention efforts, including the work of YSPA, the SPC, implementation of NH's Suicide Prevention Plan, GLS funding through SAMHSA, suicide prevention programs implemented in the state, and the work of many community partners likely played a role in that downward trend. Even though rates have recently increased, the value of prevention efforts should not be discounted. Without the continued work of these and other individuals and organizations, a greater increase in NH suicide rates may have occurred.


Figure 49 (pg. 65) presents NH suicide death rates for youth and young adults aged 10-24 in rolling three-year intervals from 2015 to 2019 and **Figure 50** (pg. 65) presents the same information for individuals of all ages. NH-VDRS data is currently limited to 2015-2019. As new data becomes available these figures will be expanded and used to identify trends in NH rates over time and compare them with national trends.

Figure 49
Suicide rates among 10-24 year old NH residents have increased from 2015-2019.



Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 50
NH Resident Suicide Death Rates for Rolling
3-Year Intervals - All Ages
2015-2019




Data Source: NH-VDRS data prepared by the NH DHHS Injury Prevention Program

Figure 51 (below) presents the results of the NH YRBS from 2011, 2013, 2015, 2017, and 2019. The percentage of high school youth in NH who felt sad or hopeless for 2+ weeks in the past year and the percentage of youth who seriously considered a suicide attempt in the past year have both increased between 2011 and 2019. In 2019, 1 in 5 youth surveyed reported having seriously considered attempting suicide in the past year, while 1 in 14 reported actually having made an attempt.

Figure 51

Self-reported depression and suicidal ideation among high school youth increased from 2011 to 2019.



Data Source: NH YRBS Results, NH Department of Education

The NH YRBS item addressing whether students have made a suicide plan in the past year was not asked from 2013-2017. This was removed due to the similarity to the question asking whether youth had seriously considered a suicide attempt during the past year. At the time the removal of this question allowed for the addition of a question addressing non-suicidal self-inflicted injuries (e.g., cutting or burning oneself without the intent of dying). Beginning in 2019 the question was again included in the survey. The results of the question on self-inflicted injuries indicate that 19.8% of NH high school age youth (12.6% of males and 27.3% of females) report intentionally hurting themselves without the intent to die during the past year (NH YRBS, 2019).

Positive Outcomes and Testimonials

Safe Messaging and Media Guidelines:

Work has been done continuously across the state to educate the public and media about safe messaging, a national best practice standard (www.sprc.org/library/SafeMessagingfinal.pdf). Safe messaging has become part of the standard for statewide and regional meetings, part of suicide prevention trainings, a guide for health promotion materials, and essentially part of the culture in NH. Media Guidelines have been disseminated to media outlets across the state, and journalism students in several universities in NH have received training in the Media Guidelines and how to safely write about suicide. The Communications/Media Sub-Committee of the SPC provides feedback to media outlets and suicide prevention experts in the state to guide public information that is produced through consultation, media contributions and feedback. The results of these efforts became evident after the tragic death of Robin Williams. Rather than sensationalizing this highly publicized tragedy, many media outlets across NH interviewed local representatives in the mental health and suicide prevention field. “Not only did the media in our state reach out to partner with key stakeholders to create responsible follow up articles, but all of the people interviewed provided the same consistent messages of hope and help for those struggling with mental illness and resources for those in crisis. It was clear that everyone, independent of each other, was reading off of the same page.”

Elaine de Mello
Supervisor of Training and Prevention Services
NAMI New Hampshire

Reading Tables and Figures

This section is intended to assist the reader in interpreting the various charts included in the report. The four topics covered in this section include types of charts; common parts of a chart; frequently used scales in charts; and interpreting the information presented in a chart. These topics contain information that applies primarily to the charts included in this report, but much of the information can also be applied elsewhere.

Types of Charts

- **Line Chart:** A line chart presents a series of connected observations in order. For example, the line chart in **Figure 4** of this report shows the number of youth and young adult suicides over a 5-year span in NH.
- **Pie Chart:** A pie chart gives the percent values for the individual parts of a whole using a circle that is divided into wedges. For example, a pie chart (**Figure 15**) of this report shows the percent of male and female youths and young adults in NH that died by suicide from 2015 to 2019.
- **Bar Chart:** A bar chart shows the values for one or more categories using rectangular boxes with height representing the value (greater height being a larger value and lesser height being a smaller value). For example, two bar charts (**Figures 10 and 11**) in this report show the number of suicide deaths by age group in NH from 2015 to 2019 and the rate of suicide deaths by age group in NH from 2015 to 2019.

Common Parts of a Chart

- **Title:** The title will generally be found at the top of the chart and should describe the data that are being presented. Depending on the chart this may list the variables and/or the time period. Also, all charts in this report list the data source used.
- **Scales/Labels:** The scales/labels are generally found on the bottom and left side of the chart. The scale/label on the bottom shows what is being measured on the x-axis (horizontal axis) and the scale/label on the left side shows what is being measured on the y-axis (vertical axis). For example, in **Figure 4**, the line chart of youth suicides in NH over the past four years has a different scale on each axis. On the x-axis (the bottom) are years which range from 2015 to 2019. On the y-axis (the side) the scale is the number of youth suicides, which ranges from 0 to 40.
- **Legend/Key:** Some charts include a legend/key to explain what different colors, shapes, dotted/solid lines mean. The location of this may vary depending on the type of chart and where space is available on the page.
- **Error Bars/Confidence Intervals:** Error bars/confidence intervals represent the range that the actual value may fall within. There is some degree of uncertainty when calculating values such as rates due to statistical error (captured by the confidence intervals) and data quality issues (which there is no real way to estimate). The width of the error bar/confidence interval indicates the level of uncertainty. A wider bar denotes more uncertainty and may indicate more data is needed. A smaller bar indicates a greater level of confidence in the results. When error bars/confidence intervals overlap in a chart, one cannot state with certainty whether there is a significant difference between the

values. Error bars can be seen on several of the charts in this document, including the NH crude death rate chart (**Figure 13**). In that chart you can see that the error bar for Merrimack County does not overlap the bar for Rockingham County. From this we are able to determine that the rate of suicide in Merrimack County is significantly different from the rate in Rockingham County.

Frequently Used Scales

- **Standard:** What is being referred to here as standard is a numbered scale that gives the actual value of the variable(s) being presented in the chart (e.g., the number of youth and young adult suicides in a given year).
- **Rate:** A scale using a rate is saying how common something is in relation to a standard value. This report uses rates per 100,000. Therefore, a youth and young adult suicide rate of 10 would mean that there are likely to be 10 suicides by youth or young adults for every 100,000 youths or young adults in the population. Rates are approximations based on past data and do not guarantee the same trend will or will not continue.
- **Percent:** A scale using percent is expressing a certain proportion of the variable falls into one category (i.e., 25% of youth is equivalent to 25 out of 100 youth).

Interpreting Information from Charts

- Can different charts be compared? Yes, but only under certain circumstances. Different charts should only be compared if they were generated using the same dataset and related variables. Depending on the charts there may be other factors that prevent you from directly comparing them. When in doubt, attempt to contact the person who made the chart or someone with access to the data used to generate the chart.
- Data is generated in a variety of ways and therefore it is not always consistent. For example, in NH the OCME is charged with keeping records of all deaths that occur in the state, regardless of where the person lived. Thus, a Vermont resident who dies in a NH hospital would be included in OCME data. On the other hand, the Bureau of Vital Records collects data on the deaths of NH residents regardless of where the death occurs. So, a NH resident who dies in Massachusetts would be included in Vital Records statistics. Therefore, these two data sets will have small differences. Neither is wrong. They simply measure different things.

Glossary of Terms

Acronyms

American Foundation for Suicide Prevention	AFSP
Army National Guard	ARNG
Behavioral Risk Factor Surveillance System	BRFSS
Centers for Disease Control and Prevention	CDC
Crisis Intervention Team	CIT
Community Mental Health Center	CMHC
Counseling on Access to Lethal Means	CALM
Department of Health and Human Services	DHHS
Electronic Data Warehouse	EDW
Emergency Departments	ED
Garrett Lee Smith	GLS
Health Insurance Portability and Accountability Act	HIPAA
Health Statistics and Data Management	HSDM
International Classification of Diseases 10 th Revision	ICD-10
National Alliance on Mental Illness New Hampshire	NAMI NH
National Suicide Prevention Lifeline	NSPL
National Violent Death Reporting System	NVDRS
New Hampshire Violent Death Reporting System	NH-VDRS
Northern New England Poison Center	NNEPC
Office of Economic Planning	OEP
Office of the Chief Medical Examiner	OCME
Post-Traumatic Stress Disorder	PTSD
Substance Abuse and Mental Health Services Administration	SAMHSA
Suicide Prevention Council	SPC
Suicide Prevention Program	SPP
Suicide Prevention Resource Center	SPRC
Survivor of Suicide Loss	SOSL
Traumatic Brain Injury	TBI
Veterans Administration	VA
Web-based Injury Statistics Query and Reporting System	WISQARS
Youth Risk Behavior Survey	YRBS
Youth Suicide Prevention Assembly	YSPA

Age Adjustment and Rates

When possible, rates in this document are age-adjusted to the 2010 US standard population. This allows the comparison of rates among populations having different age distributions by standardizing the age-specific rates in each population to one standard population. Age-adjusted rates refer to the number of events that would be expected per 100,000 persons in a selected population if that population had the same age distribution as a standard population. Age-adjusted rates were calculated using the direct method as follows:

Where,

m = number of age groups

di = number of events in age group i

Pi = population in age group i

Si = proportion of the standard population in age group i

This is a weighted sum of Poisson random variables, with the weights being (Si /pi).

$$\hat{R} = \sum_{i=1}^m s_i (d_i/p_i) = \sum_{i=1}^m w_i d_i$$

Age Specific Rate/Crude Rates

The age-specific rate or crude rate is the number of individuals with the same health issue per year within a specific age group, divided by the estimated number of individuals of that age living in the same geographic area at the midpoint of the year.

Confidence Intervals (Ci)

The standard error can be used to evaluate statistically significant differences between two rates by calculating the confidence interval. If the interval produced for one rate does not overlap the interval for another, the probability that the rates are statistically different is 95% or higher.

The formula used is:

Where,

R=age-adjusted rate of one population

z = 1.96 for 95% confidence limits

SE= standard error as calculated below

$$R + z (SE)$$

A confidence interval is a range of values within which the true rate is expected to fall. If the confidence intervals of two groups (such as NH and the US) overlap, then any difference between the two rates is not statistically significant. All rates in this report are calculated at a 95% confidence level.

Data Collection

The BRFSS is a telephone survey conducted annually by the health departments of all 50 states, including NH. The survey is conducted with assistance from the federal CDC. The BRFSS is the largest continuously conducted telephone health survey in the world and is the primary source of information for states and the nation on the health-related behaviors of adults. The BRFSS has been conducted in NH since 1987. HSDM develops the annual questionnaire, plans survey protocol, locates financial support, and monitors data collection progress and quality with the

assistance of CDC. HSDM employs a contractor for telephone data collection. Survey data are submitted monthly to CDC by the contractor for cleaning and processing and then returned to HSDM for analysis and reporting.

Death Certificate Data is collected by the Department of Vital Records in NH and provided to the HSDM through a Memorandum of Understanding. Death Certificate Data is available to the HSDM through the state Electronic Data Warehouse (EDW), a secure data server.

Hospital Discharge Data for inpatient and emergency department care is compiled, and de-identified at the Maine Health Information Center, delivered to the Office of Medicaid Business and Policy for further cleaning, then available to the HSDM through the state EDW.

State and county population estimates for NH data are provided by HSDM, Bureau of Disease Control and Health Statistics, Division of Public Health Services, and NH DHHS. Population data are based on US Census data apportioned to towns using NH Office of Economic Planning (OEP) estimates and projections, and further apportioned to age groups and gender using Claritas Corporation estimates and projections to the town, age group, and gender levels. Data add up to US Census data at the county level between 1990 and 2005 but do not add to OEP or Claritas data at smaller geographic levels.

Data Confidentiality

The data provided in this report adheres to the NH DHHS “Guidelines for Release of Public Health Data” and the Health Insurance Portability and Accountability Act (HIPAA). Data are aggregated in-to groups large enough to prevent constructive identification of individuals who were discharged from hospitals or who are deceased.

Graphs

Graphs have varying scales depending on the range of the data displayed. Therefore, caution should be exercised when comparing such graphs.

Incidence

Incidence refers to the number or rate of new cases in a population. Incidence rate is the probability of developing a particular disease or injury occurring during a given period of time; the numerator is the number of new cases during the specified time period and the denominator is the population at risk during the period. Rates are age-adjusted to 2010 US standard population. Some of the rates also include age-specific rates. Rates based on 10 or fewer cases are not calculated, as they are not reliable.

Death Rate

Death rate is the number of deaths per 100,000 in a certain region in a certain time period and is based on International Classification of Diseases 10th Revision (ICD-10). Cause of death before 1999 was coded according to ICD-9; beginning with deaths in 1999, ICD-10 was used.

Reliability of Rates

Several important notes should be kept in mind when examining rates. Rates based on small numbers of events (e.g., less than 10 events) can show considerable variation. This limits the usefulness of these rates in comparisons and estimations of future occurrences. Unadjusted rates (age-specific or crude rates) are not reliable for drawing definitive conclusions when making comparisons because they do not take factors such as age distribution among populations into account. Age-adjusted rates offer a more refined measurement when comparing events over geographic areas or time periods. When a difference in rates appears to be significant, care should be exercised in attributing the difference to any particular factor or set of factors. Many variables may influence rate differences. Interpretation of a rate difference requires substantial data and exacting analysis.

Small Numbers

With very small counts, it is often difficult to distinguish between random fluctuation and meaningful change. According to the National Center for Health Statistics, considerable caution must be observed in interpreting the data when the number of events is small (perhaps less than 100) and the probability of such an event is small (such as being diagnosed with a rare disease). The limited number of years of data in the registry and the small population of the state require policies and procedures to prevent the unintentional identification of individuals. Data on rare events, and other variables that could potentially identify individuals, are not published.

Standard Errors

The standard errors of the rates were calculated using the following formula:

Where,

w_j = fraction of the standard population in age category

n_j = number of cases in that age category

p = person-years denominator

$$S.E. = \sqrt{\frac{w_j^2 n_j}{p_j^2}}$$

Frequently Asked Questions about NH Suicide Data

Q: Statistical significance of suicide deaths vs. significance in the community.

A: Statistical significance, which this document focuses on, is used to look at whether the change in the number of suicide deaths from one time period to another has truly increased/decreased, or whether the difference is potentially due to chance. In general, in NH a small number of additional deaths are unlikely to result in a statistically significant change. However, the significance of even a single death in a family or a community is tremendous. When discussing “significance” it is best to be clear about whether the focus is on measurable changes or the practical impact on a family or community.

Q: Have there been more suicide deaths in NH during “X” months of this year compared with previous years?


A: It is best to focus on data from a full year or multiple years rather than periods of just a few months. Over brief periods these numbers are too volatile to draw accurate conclusions from them.

Q: If there is an increase during part of a year does this mean that there will be a greater number of suicide deaths during the remainder of the year when compared with previous years?

A: Not necessarily. Even though there may have been a greater number of deaths during part of a given year, this does not indicate that there will be a greater number of deaths for the remainder of the year. Until the end of the year it is not possible to say whether the overall number of suicide deaths will be higher or lower than previous years.

Q: Has NH ever had a large change in suicide deaths from one year to the next?

A: As a small state, NH has a substantial degree of variability in the suicide deaths in a given year. It is not at all uncommon for the number (and rate) of suicide deaths in NH to vary by as much as 33% (up or down) from the previous year – see chart and table below. Significant differences are indicated by non-overlapping confidence intervals (the brackets overlaid on the bars in the chart). For example, the confidence intervals for 2013 do not overlap with the 2014 or 2017-2019 confidence intervals, meaning that the rates for 2014 and 2017-2019 were significantly higher than the rate for 2013.



Change in Rate per 100,000 from Year to Year	
2010-2011	14.89 to 15.02 (Up 1%)
2011-2012	15.02 to 15.25 (Up 2%)
2012-2013	15.25 to 13.95 (Down 9%)
2013-2014	13.95 to 18.52 (Up 33%)
2014-2015	18.52 to 17.06 (Down 8%)
2015-2016	17.06 to 18.18 (Up 7%)
2016-2017	18.18 to 19.65 (Up 8%)
2017-2018	19.65 to 20.61 (Up 5%)
2017-2018	20.61 to 18.75 (Down 9%)

Q: What are the differences between the Centers for Disease Control (CDC) data and NH data on suicide deaths?

A: The CDC data includes all deaths of NH residents regardless of whether they occurred in the state or elsewhere. The NH data comes directly from the Office of the Chief Medical Examiner (OCME) and includes all suicide deaths that have occurred in the state, even if the death was of a non-resident. Also, CDC data are often not released until 12-24 months after the end of a calendar year (e.g., 2019 data were released in early-2021). Preliminary NH data are available within months of a calendar year ending.

Q: What is the difference between a rate and a count?

A: A count simply shows the number of incidents that have taken place during a given period of time (e.g., 100 deaths in a one-year period). A rate is a way of showing the prevalence of something among the population. For example, saying that there are 10 deaths resulting from “x” per 100,000 means that in a given population approximately 10 out of every 100,000 individuals have been found to die as a result of “x”.

Q: Has “X” (e.g., the recession) caused the increase/decrease in the number of suicide deaths in a specific year?

A: Suicide is a complex issue, and it is not possible to say that a single factor is the direct cause of these deaths. For instance, from 2013 to 2014, the number of deaths were up over 33% followed by an 8% decrease from 2014 to 2015; we are still unable to identify the underlying cause of these fluctuations and whether any of those deaths are attributable to the same cause.

Q: How do the number of suicide deaths compare to other causes of death in the state?

A: 10 Leading Causes of Death, New Hampshire, by Age Group, 2015 – 2019

Rank	Age Groups										All Ages
	<1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65+	
1	Short Gestation 33	Unintentional Injury 10	Malignant Neoplasms ...	Malignant Neoplasms 11	Unintentional Injury 321	Unintentional Injury 818	Unintentional Injury 642	Malignant Neoplasms 866	Malignant Neoplasms 2,622	Heart Disease 11,250	Malignant Neoplasms 13,938
2	Congenital Anomalies 29	Congenital Anomalies ...	Unintentional Injury ...	Suicide 11	Suicide 152	Suicide 182	Malignant Neoplasms 211	Unintentional Injury 608	Heart Disease 1,395	Malignant Neoplasms 10,137	Heart Disease 13,416
3	Maternal Pregnancy Comp. 15	Homicide ...	Homicide ...	Unintentional Injury ...	Heart Disease 20	Malignant Neoplasms 62	Suicide 195	Heart Disease 577	Unintentional Injury 484	Chronic Low Respiratory Disease 3,136	Unintentional Injury 4,469
4	Placenta Cord Membranes 15	Influenza & Pneumonia ...	Benign Neoplasms ...	Congenital Anomalies ...	Malignant Neoplasms 18	Heart Disease 49	Heart Disease 118	Suicide 272	Chronic Low Respiratory Disease 397	Alzheimer's Disease 2,267	Chronic Low Respiratory Disease 3,641
5	Respiratory Distress 10	Malignant Neoplasms ...	Congenital Anomalies ...	Benign Neoplasms ...	Homicide 14	Liver Disease 15	Liver Disease 68	Liver Disease 169	Diabetes Mellitus 277	Cerebro-vascular 2,203	Cerebro-vascular 2,486
6	Circulatory System Disease ...	Diabetes Mellitus ...	Heart Disease ...	Chronic Low Respiratory Disease ...	Chronic Low Respiratory Disease ...	Homicide 13	Diabetes Mellitus 31	Diabetes Mellitus 101	Liver Disease 277	Unintentional Injury 1,566	Alzheimer's Disease 2,296
7	Intrauterine Hypoxia ...	Heart Disease ...	Acute Bronchitis ...	Influenza & Pneumonia ...	Congenital Anomalies ...	Cerebro-vascular 11	Homicide 25	Chronic Low Respiratory Disease 81	Suicide 223	Diabetes Mellitus 1,226	Diabetes Mellitus 1,646
8	Bacterial Sepsis	Diseases Of Appendix ...	Nephritis ...	Cerebro-vascular ...	Congenital Anomalies ...	Cerebro-vascular 17	Cerebro-vascular 73	Cerebro-vascular 175	Influenza & Pneumonia 1,070	Suicide 1,271
9	Unintentional Injury	Pneumonitis ...	Diabetes Mellitus ...	Diabetes Mellitus ...	Chronic Low Respiratory Disease 16	Septicemia 36	Nephritis 84	Nephritis 810	Influenza & Pneumonia 1,167
10	SIDS	Four Tied ...	Two Tied ...	Congenital Anomalies 11	Congenital Anomalies 24	Septicemia 83	Parkinson's Disease 786	Nephritis 926

Data Source: CDC WISQARS, 2015-2019

--Note: Beginning with 2008 data, the CDC has suppressed state-level counts for categories with fewer than ten deaths

Contacts and Meeting Information

**Please note that the 2020 and 2021 meeting schedules and formats may vary due to COVID-19.
Contact the identified individual(s) below to confirm meeting the details if you would like to attend.**

State Suicide Prevention Council

Chair: Russell Conte – Russell.Conte@dos.nh.gov

Vice Chairs: Candice Porter - candice.porter@connorsclimb.org

Mary Forsythe-Taber – mft@mih4u.org

Meets 4th Monday – Every **other** month 10:00 am – 12:00 pm

Philbrook Adult Transitional Housing, 121 South Fruit Street, Concord/Video conference

Youth Suicide Prevention Assembly

Primary Contact: Elaine de Mello – edemello@naminh.org

Meets 2nd Thursday of the month 10:00 – 12:30 am

Currently meeting via video conference. Future in person meeting location to be determined.

Suicide Prevention Council Subcommittees

Communications & Public Education

Chair: Mary Forsythe-Taber – mft@mih4u.org

Meets 2nd Wednesday of the month 10:00 am – 12:00 pm

All meetings held via video conference – contact the chair if interested in attending

Data Collection & Analysis

Chair: Patrick Roberts – proberts@naminh.org

Meets 4th Wednesday of the Feb., May, Aug., and Oct. 9:30 – 11:30 am

NAMI NH, 85 North State Street, Concord/Video conference

Law Enforcement

Chair: Trooper Seth Gahr

Meeting schedule to be determined

Military & Veterans

Co-Chairs: Amy Cook – Amy.Cook@nh.gov

Beth Alves - Beth.Alves@va.gov

Meets 1st Wednesday of the Month 2:00 – 3:30 pm

Location varies, please contact co-chairs for location information.

Public Policy

Co-Chairs: Candace Porter - candice.porter@connorsclimb.org
Debbie Robinson - deborah.a.robinson@dhhs.nh.gov

Contact the co-chairs for meeting dates/location.

State Suicide Prevention Conference Meetings

Primary Contact: Elaine de Mello – edemello@naminh.org

Contact Elaine de Mello for current meeting schedule and location

Suicide Fatality Review

Chair: Dr. Paul Brown

Attendance is by invitation only

Survivors of Suicide Loss

Co-Chairs: Deb Baird – dbaird0688@gmail.com
Shamera Simpson – ssimpson@afsp.org

Meets 1st Tuesday of the Month 10:00 am – 11:30 am
All meetings held via video conference

Recognize the Warning Signs for Suicide to Save Lives!

Sometimes it can be difficult to tell warning signs from “normal” behavior especially in adolescents. Ask yourself, *is the behavior I am seeing very different for this particular person?* Also, recognize that sometimes those who are depressed can appear angry, irritable, and/or hostile in addition to withdrawn and quiet.

Warning signs:

- Talking about or threatening to hurt or kill oneself
- Seeking firearms, drugs, or other lethal means for killing oneself
- Talking or writing about death, dying, or suicide
- Direct Statements or Less Direct Statements of Suicidal Intent: (Examples: “I’m just going to end it all” or “Everything would be easier if I wasn’t around.”)
- Feeling hopeless
- Feeling rage or uncontrollable anger or seeking revenge
- Feeling trapped - like there’s no way out
- Dramatic mood changes
- Seeing no reason for living or having no sense of purpose in life
- Acting reckless or engaging in risky activities
- Increasing alcohol or drug use
- Withdrawing from friends, family, and society
- Feeling anxious or agitated
- Being unable to sleep, or sleeping all the time

For a more complete list of warning signs and more information on suicide prevention, please consult the *Connect* website at www.theconnectprogram.org and click on Resources.

If you see warning signs and/or are otherwise worried about this person:

Connect with Your Loved One, Connect Them to Help

- 1) Ask directly about their suicidal feelings. Talking about suicide is the first step to preventing suicide!
- 2) Let them know you care.
- 3) Keep them away from anything that may cause harm such as guns, pills, ropes, knives, vehicles.
- 4) Stay with them until a parent or professional is involved.
- 5) Offer a message of hope - Let them know you will assist them in getting help.
- 6) Connect them with help:
 - * National Suicide Prevention Lifeline (24/7) **1-800-273-TALK (8255)** (press “1” for veterans)
 - * The Lifeline also offers text-based chat through their website: www.suicidepreventionlifeline.org/
 - * Head rest – For teens and adults (24/7) **1-800-639-6095** or your local mental health center
 - * For an emergency, **dial 911.**



Mental Health and Suicide Prevention Resources

General Resources:

Local Resources

Community Mental Health Centers: <http://www.dhhs.state.nh.us/dcbcs/bbh/centers.htm>
Disaster Behavioral Health Response Teams: <http://www.dhhs.nh.gov/esu/dbhrtnh.htm>
NAMI New Hampshire: www.NAMINH.org, 603-225-5359

Gay, Lesbian Bisexual, and Transgender (GLBT) Resources

Fenway Peer Listening Line: 1-800-399-PEER www.fenwayhealth.org
GLBT National Hotline (M-F 4-12 pm; Sat. 12-5 pm): 1-888-843-4564 www.glnh.org
GLBT National Youth Talkline (M-F 8-12 pm): 1-800-246-PRIDE (7743)
Email: youth@GLBTNationalHelpCenter.org
SPRC Library: www.sprc.org/library_resources/sprc
Trevor Helpline (24/7): 1-866-4u-TREVOR (488-7386) www.thetrevorproject.org

Military Resources

Military One Source: www.militaryonesource.mil
Tragedy Assistance Program for Survivors (TAPS): www.taps.org
US Department of Veterans Affairs: www.va.gov
Veterans Crisis Line: 1-800-273-8255 (press 1 after connecting)

National Organizations

American Association of Suicidology: www.suicidology.org
American Foundation for Suicide Prevention: www.afsp.org
National Action Alliance for Suicide Prevention: theactionalliance.org
National Alliance on Mental Illness: www.nami.org
Suicide Prevention Resource Center: www.sprc.org

Older Adults

NH Fact Sheet on Suicide and Aging: bit.ly/2nuLd5O
SPRC Older Adult Suicide Prevention Resources: www.sprc.org/populations/older-adults

Substance Abuse and Mental Health Services Administration (SAMHSA)

Obtaining Prevention Materials:
Visit their website: store.samhsa.gov/ (includes downloadable materials)
Call: 1-877-SAMHSA-7 (1-877-726-4727) or Email: samhsainfo@samhsa.hhs.gov
Treatment Provider Locator:
SAMHSA maintains a searchable list of mental health and substance use disorder providers.
You can use it to find a local provider by going to www.samhsa.gov/find-treatment

Resources for Survivors of Suicide Loss / Individuals Bereaved by Suicide:

National Helplines

Compassionate Friends: 1-877-696-0010

Friends for Survival: 1-800-646-7322

Websites

Alliance of Hope for Suicide Survivors: www.allianceofhope.org

American Foundation for Suicide Prevention: afsp.org

Compassionate Friends: www.compassionatefriends.org

The Connect Program: <https://theconnectprogram.org/find-support/coping-with-suicide-loss>

Friends for Survival: www.friendsforsurvival.org

Heartbeat: www.heartbeatsurvivorsafter-suicide.org

Parents, Family and Friends of Suicide Loss: www.pos-ffos.com

SAVE (Suicide Awareness Voices of Education): www.save.org/coping

Survivors of Suicide Loss: www.survivorsofsuicide.com

Discussion Forums

Help for People Left Behind: forums.grieving.com

Suicide's Survivors: bit.ly/legacy-suicidesurvivors

Booklets

Coping with the Loss of a Friend or Loved One: bit.ly/save-coping-withloss

Handbook for Survivors of Suicide: bit.ly/2lnGUsm

Hope and Healing after Suicide: bit.ly/2n0cxse

Resource and Healing Guide: bit.ly/2nyiEVg

Have you found this report to be useful?

Please share your feedback through the survey linked below so that this report can be even better in the future.

<https://www.surveymonkey.com/r/V99FMBP>