

Activities to Support Infant Oral Function

Bibliography

© Melissa Cole, MS, IBCLC

- AAP (n/d). Noise: a hazard for the fetus and newborn. American Academy of Pediatrics. Committee on Environmental Health. (1997). *Pediatrics*, 100(4), 724-727.
- AAP (n/d). Prevention and management of pain and stress in the neonate. American Academy of Pediatrics. Committee on Fetus and Newborn. Committee on Drugs. Section on Anesthesiology. Section on Surgery. Canadian Paediatric Society. Fetus and Newborn Committee. (2000). [Guideline Practice Guideline]. *Pediatrics*, 105(2), 454-461.
- AAP (n/d). The assessment and management of acute pain in infants, children, and adolescents. (2001). [Guideline Practice Guideline]. *Pediatrics*, 108(3), 793-797.
- Abdel Razeq, N. M., Akuma, A. O., & Jordan, S. (2016). Status of Neonatal Pain Assessment and Management in Jordan. *Pain Manag Nurs*, 17(4), 239-248. doi: 10.1016/j.pmn.2016.02.050
- Al Qahtani, A. M., & Ahmed, H. M. (2020). The Effect of Educational Program for New Mothers about Infant Abdominal Massage and Foot Reflexology for Decreasing Colic at Najran City. *Compr Child Adolesc Nurs*, 1-16. doi:10.1080/24694193.2020.1740827
- Allegaert, K., & van den Anker, J. N. (2016). Neonatal pain management: still in search for the Holy Grail. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Review]. *Int J Clin Pharmacol Ther*, 54(7), 514-523. doi: 10.5414/CP202561
- Allen, K. A. (2013). Music therapy in the NICU: is there evidence to support integration for procedural support? [Research Support, N.I.H., Extramural Review]. *Adv Neonatal Care*, 13(5), 349-352. doi: 10.1097/ANC.0b013e3182a0278b
- Alvares, D., Torsney, C., Beland, B., Reynolds, M., & Fitzgerald, M. (2000). Modelling the prolonged effects of neonatal pain. *Prog Brain Res*, 129, 365-373.
- Alves Cde, O., Duarte, E. D., Azevedo, V. M., Nascimento, G. R., & Tavares, T. S. (2011). [Use of sweet solutions for neonatal pain relief in premature newborns: an integrative review]. [Review]. *Revista gaucha de enfermagem / EENFUFGRS*, 32(4), 788-796.
- Arnold, A. P. (2009). *Rhythm and Touch: the Fundamentals of Craniosacral Therapy*. Berkeley, Calif.: North Atlantic Books.
- Arvedson, J. C., & Brodsky, L. (2002). *Pediatric swallowing and feeding: assessment and management* (2nd ed.). Australia; Albany, NY: Singular Thomson Learning.
- Avila-Alvarez, A., Carbajal, R., Courtois, E., Pertega-Diaz, S., Anand, K. J., & Muniz-Garcia, J. (2016). [Clinical assessment of pain in Spanish Neonatal Intensive Care Units]. [English abstract]. *Anales de pediatria*, 85(4), 181-188. doi: 10.1016/j.anpedi.2015.09.019
- Axelin, A., Salanterä, S., & Lehtonen, L. (2006). 'Facilitated tucking by parents' in pain management of preterm infants—a randomized crossover trial. *Early human development*, 82(4), 241-247. doi: <http://dx.doi.org/10.1016/j.earlhumdev.2005.09.012>
- Azevedo-Santos, I. F., Lima, L. V., & DeSantana, J. M. (2016). Influence of Neonatal Pain in Motor Development During Childhood. *Pain Med*, 17(6), 1204-1206. doi: 10.1093/pm/pnv059
- Baccei, M. L. (2016). Rewiring of Developing Spinal Nociceptive Circuits by Neonatal Injury and Its Implications for Pediatric Chronic Pain. [Review]. *Children (Basel)*, 3(3). doi: 10.3390/children3030016
- Badr, L. K., Abdallah, B., & Kahale, L. (2015). A Meta-Analysis of Preterm Infant Massage: An Ancient Practice With Contemporary Applications. *MCN. The American journal of maternal child nursing*, 40(6), 344-358. doi:10.1097/nmc.0000000000000177
- Ballard, J. L., Auer, C. E., & Khoury, J. C. (2002). Ankyloglossia: assessment, incidence, and effect of frenuloplasty on the breastfeeding dyad. [Comparative Study Research Support, Non-U.S. Gov't]. *Pediatrics*, 110(5), e63.
- Barbul, A., Lazarou, S. A., Efron, D. T., Wasserkrug, H. L., & Efron, G. (1990). Arginine enhances wound healing and lymphocyte immune responses in humans. [Clinical Trial Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Surgery*, 108(2), 331-336; discussion 336-337.
- Beckman, D. (2013). *Beckman Oral Motor Assessment and Intervention*. Paper presented at the Beckman Oral Motor Assessment and Intervention, Seattle, Washington.
- Bellieni, C. V., Buonocore, G., Nenci, A., Franci, N., Cordelli, D. M., & Bagnoli, F. (2001). Sensorial saturation: an effective analgesic tool for heel-prick in preterm infants: a prospective randomized trial. [Clinical Trial Randomized Controlled Trial]. *Biology of the neonate*, 80(1), 15-18.

- Belliemi, C. V., Buonocore, G., Nenci, A., Franci, N., Cordelli, D. M., & Bagnoli, F. (2001). Sensorial saturation: an effective analgesic tool for heel-prick in preterm infants: a prospective randomized trial. [Clinical Trial Randomized Controlled Trial]. *Biology of the neonate*, 80(1), 15-18.
- Bergman, J. (2010). *Hold Your Premie*. Cape Town, South Africa: New Voices publishing.
- Bong, C. L., Hilliard, J., & Seefelder, C. (2009). Severe methemoglobinemia from topical benzocaine 7.5% (baby orajel) use for teething pain in a toddler. [Case Reports]. *Clinical pediatrics*, 48(2), 209-211. doi: 10.1177/0009922808324491
- Bovey, A., Noble, R., & Noble, M. (1999). Orofacial exercises for babies with breastfeeding problems? *Breastfeeding review: professional publication of the Nursing Mothers' Association of Australia*, 7(1), 23-28. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10197368>
- Brown, K. L., & Phillips, T. J. (2010). Nutrition and wound healing. *Clin Dermatol*, 28(4), 432-439. doi: 10.1016/j.clindermatol.2010.03.028
- Buchheit, T., Van de Ven, T., & Shaw, A. (2012). Epigenetics and the transition from acute to chronic pain. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, Non-P.H.S. Review]. *Pain Med*, 13(11), 1474-1490. doi: 10.1111/j.1526-4637.2012.01488.x
- Buryk, M., Bloom, D., & Shope, T. (2011). Efficacy of neonatal release of ankyloglossia: a randomized trial. *Pediatrics*, 128(2), 280-288. doi: 10.1542/peds.2011-0077
- Butkevich, I. P., Mikhailenko, V. A., Vershinina, E. A., & Aloisi, A. M. (2016). Effects of neonatal pain, stress and their interrelation on pain sensitivity in later life in male rats. *Chin J Physiol*, 59(4), 225-231. doi: 10.4077/CJP.2016.BAE412
- Carbajal, R., Guedj, R., Rambaud, J., & Leger, P. L. (2016). Impact of a systematic neonatal pain and sedation protocol. [Editorial]. *Acta paediatrica*, 105(7), 734-735. doi: 10.1111/apa.13373
- Carmo Ede, C., Sanada, L. S., Machado, N. L., & Fazan, V. P. (2016). Does Pain in the Neonatal Period Influence Motor and Sensory Functions in a Similar Way for Males and Females During Post-Natal Development in Rats? *Pain Med*, 17(8), 1520-1529. doi: 10.1093/pm/pnv117
- Castro, M. (1993). *Homeopathy for Pregnancy, Birth, and Your Baby's First Year* (1st U.S. ed.). New York: St. Martin's Press.
- Christoffel, M. M., Castral, T. C., Dare, M. F., Montanholi, L. L., & Scochi, C. G. (2016). Knowledge of healthcare professionals on the evaluation and treatment of neonatal pain. *Revista brasileira de enfermagem*, 69(3), 552-558. doi: 10.1590/0034-7167.2016690319i
- Cho, A., Kelsberg, G., & Safranek, S. (2010). Clinical inquiries. When should you treat tongue-tie in a newborn? [Review]. *The Journal of family practice*, 59(12), 712a-b.
- Chopra, K., Tadisina, K. K., Patel, K. R., & Singh, D. P. (2013). Syndactyly repair. *Eplasty*, 13, ic51.
- Cignacco, E., Axelin, A., Stoffel, L., Sellam, G., Anand, K., & Engberg, S. (2010). Facilitated tucking as a non-pharmacological intervention for neonatal pain relief: is it clinically feasible? *Acta paediatrica*, 99(12), 1763-1765. doi: 10.1111/j.1651-2227.2010.01941.x
- Cignacco, E. L., Sellam, G., Stoffel, L., Gerull, R., Nelle, M., Anand, K. J., & Engberg, S. (2012). Oral sucrose and "facilitated tucking" for repeated pain relief in preterms: a randomized controlled trial. [Comparative Study Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Pediatrics*, 129(2), 299-308. doi: 10.1542/peds.2011-1879
- Cole, M., & Ghaheri, B. (2012). General Information about Frenulum Procedures for the Infant (pp. 2).
- Courtois, E., Cimerman, P., Dubuche, V., Goiset, M. F., Orfevre, C., Lagarde, A., Carbajal, R. (2016). The burden of venipuncture pain in neonatal intensive care units: EIPPAIN 2, a prospective observational study. [Research Support, Non-U.S. Gov't]. *International journal of nursing studies*, 57, 48-59. doi: 10.1016/j.ijnurstu.2016.01.014
- Courtois, E., Drouman, S., Magny, J. F., Merchaoui, Z., Durmeyer, X., Roussel, C., Carbajal, R. (2016). Epidemiology and neonatal pain management of heelsticks in intensive care units: EIPPAIN 2, a prospective observational study. *International journal of nursing studies*, 59, 79-88. doi: 10.1016/j.ijnurstu.2016.03.014
- da Costa, S. P., & van der Schans, C. P. (2008). The reliability of the Neonatal Oral-Motor Assessment Scale. *Acta paediatrica*, 97(1), 21-26. doi: 10.1111/j.1651-2227.2007.00577.x
- Dao, K. D., Shin, A. Y., Billings, A., Oberg, K. C., & Wood, V. E. (2004). Surgical treatment of congenital syndactyly of the hand. [Review]. *J Am Acad Orthop Surg*, 12(1), 39-48.
- Davis, R. H., Leitner, M. G., Russo, J. M., & Byrne, M. E. (1989). Wound healing. Oral and topical activity of Aloe vera. *J Am Podiatr Med Assoc*, 79(11), 559-562.
- de Melo, G. M., Lelis, A. L., de Moura, A. F., Cardoso, M. V., & da Silva, V. M. (2014). [Pain assessment scales in newborns: integrative review]. *Rev Paul Pediatr*, 32(4), 395-402. doi: 10.1016/j.rpped.2014.04.007

- de Weerth, C., van Hees, Y., & Buitelaar, J. K. (2003). Prenatal maternal cortisol levels and infant behavior during the first 5 months. [Research Support, Non-U S Gov't]. *Early human development*, *74*(2), 139-151.
- Derbyshire, S. W. (2006). Can fetuses feel pain? [Research Support, Non-U.S. Gov't]. *BMJ*, *332*(7546), 909-912. doi: 10.1136/bmj.332.7546.909
- Dezhdar, S., Jahanpour, F., Firouz Bakht, S., & Ostovar, A. (2016). The Effects of Kangaroo Mother Care and Swaddling on Venipuncture Pain in Premature Neonates: A Randomized Clinical Trial. *Iran Red Crescent Med J*, *18*(4), e29649. doi: 10.5812/ircmj.29649
- Diego, M. A., Field, T., Hernandez-Reif, M., Deeds, O., Ascencio, A., & Begert, G. (2007). Preterm infant massage elicits consistent increases in vagal activity and gastric motility that are associated with greater weight gain. [Randomized Controlled Trial Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't]. *Acta paediatrica*, *96*(11), 1588-1591. doi: 10.1111/j.1651-2227.2007.00476.x
- Dollberg, S., Botzer, E., Grunis, E., & Mimouni, F. B. (2006). Immediate nipple pain relief after frenotomy in breast-fed infants with ankyloglossia: a randomized, prospective study. [Randomized Controlled Trial]. *Journal of pediatric surgery*, *41*(9), 1598-1600. doi: 10.1016/j.jpedsurg.2006.05.024
- Dryden, S. V., Shoemaker, W. G., & Kim, J. H. (2013). Wound management and nutrition for optimal wound healing. *Atlas Oral Maxillofac Surg Clin North Am*, *21*(1), 37-47. doi: 10.1016/j.cxom.2012.12.008
- Duhn, L. J., & Medves, J. M. (2004). A systematic integrative review of infant pain assessment tools. [Research Support, Non-U.S. Gov't Review]. *Adv Neonatal Care*, *4*(3), 126-140.
- Dumas, P., Freidel, M., & Trepsat, F. (1977). [The labial scar of hare lip. Technic and usefulness of early and prolonged postoperative massage]. *Journal francais d'oto-rhino-laryngologie, audio-phonologie et chirurgie maxillo-faciale*, *26*(9), 679-681.
- Edmunds, J., Miles, S. C., & Fulbrook, P. (2011). Tongue-tie and breastfeeding: a review of the literature. [Review]. *Breastfeeding review: professional publication of the Nursing Mothers' Association of Australia*, *19*(1), 19-26.
- Edmunds, J. E., Fulbrook, P., & Miles, S. (2013). Understanding the experiences of mothers who are breastfeeding an infant with tongue-tie: a phenomenological study. *Journal of human lactation: official journal of International Lactation Consultant Association*, *29*(2), 190-195. doi: 10.1177/0890334413479174
- Efe, E., & Ozer, Z. C. (2007). The use of breast-feeding for pain relief during neonatal immunization injections. [Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Applied nursing research: ANR*, *20*(1), 10-16. doi: 10.1016/j.apnr.2005.10.005
- Eriksson, M., & Finnstrom, O. (2004). Can daily repeated doses of orally administered glucose induce tolerance when given for neonatal pain relief? [Clinical Trial Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Acta paediatrica*, *93*(2), 246-249.
- Fitzgerald, M. (1995). Developmental biology of inflammatory pain. [Review]. *British journal of anaesthesia*, *75*(2), 177-185.
- Forstadt, L., Cooper, S., & Andrews, S. M. (2015). Changing Medicine and Building Community: Maine's Adverse Childhood Experiences Momentum. *Perm J*, *19*(2), 92-95. doi: 10.7812/TPP/14-169
- Frantz, F. W., Bettinger, D. A., Haynes, J. H., Johnson, D. E., Harvey, K. M., Dalton, H. P., Cohen, I. K. (1993). Biology of fetal repair: the presence of bacteria in fetal wounds induces an adult-like healing response. [Research Support, U.S. Gov't, P.H.S.]. *Journal of pediatric surgery*, *28*(3), 428-433; discussion 433-424.
- Fronza, M., Heinzmann, B., Hamburger, M., Laufer, S., & Merfort, I. (2009). Determination of the wound healing effect of Calendula extracts using the scratch assay with 3T3 fibroblasts. [Research Support, Non-U.S. Gov't]. *J Ethnopharmacol*, *126*(3), 463-467. doi: 10.1016/j.jep.2009.09.014
- Genna, C. W. (2008). *Supporting sucking skills in breastfeeding infants*. Sudbury, Mass.: Jones and Bartlett Publishers.
- Gernsbacher, M. A., Sauer, E. A., Geye, H. M., Schweigert, E. K., & Hill Goldsmith, H. (2008). Infant and toddler oral- and manual-motor skills predict later speech fluency in autism. *J Child Psychol Psychiatry*, *49*(1), 43-50. doi:10.1111/j.1469-7610.2007.01820.x
- Ghaheri, B. A., Cole, M., Fausel, S. C., Chuop, M., & Mace, J. C. (2016). Breastfeeding improvement following tongue-tie and lip-tie release: A prospective cohort study. *The Laryngoscope*, n/a-n/a. doi: 10.1002/lary.26306
- Ghomi, H., Yadegari, F., Soleimani, F., Knoll, B. L., Noroozi, M., & Mazouri, A. (2019). The effects of premature infant oral motor intervention (PIOMI) on oral feeding of preterm infants: A randomized clinical trial. *International journal of pediatric otorhinolaryngology*, *120*, 202-209. doi:10.1016/j.ijporl.2019.02.005
- Gibbins, S., Stevens, B., Hodnett, E., Pinelli, J., Ohlsson, A., & Darlington, G. (2002). Efficacy and safety of sucrose for procedural pain relief in preterm and term neonates. [Clinical Trial Randomized Controlled Trial]. *Nursing research*, *51*(6), 375-382.

- Glynn, L. M., Davis, E. P., Schetter, C. D., Chicz-Demet, A., Hobel, C. J., & Sandman, C. A. (2007). Postnatal maternal cortisol levels predict temperament in healthy breastfed infants. [Research Support, N I H, Extramural]. *Early human development*, 83(10), 675-681.
- Goubet, N., Rattaz, C., Pierrat, V., Bullinger, A., & Lequien, P. (2003). Olfactory experience mediates response to pain in preterm newborns. [Clinical Trial]. *Developmental psychobiology*, 42(2), 171-180.
- Gray, C. (2010). *Craniosacral Therapy for Infants*. Paper presented at the Craniosacral Therapy for Infants, Portland, Oregon.
- Gray, L., Watt, L., & Blass, E. M. (2000). Skin-to-skin contact is analgesic in healthy newborns. [Clinical Trial Randomized Controlled Trial Research Support, Non-U S Gov't Research Support, U S Gov't, P H S]. *Pediatrics*, 105(1).
- Green, J., Darbyshire, P., Adams, A., & Jackson, D. (2016). It's agony for us as well: Neonatal nurses reflect on iatrogenic pain. *Nurs Ethics*, 23(2), 176-190. doi: 10.1177/0969733014558968
- Grunau, R. E. (2013). Neonatal pain in very preterm infants: long-term effects on brain, neurodevelopment and pain reactivity. *Rambam Maimonides Med J*, 4(4), e0025. doi: 10.5041/RMMJ.10132
- Halimaa, S. L. (2003). Pain management in nursing procedures on premature babies. [Review]. *Journal of advanced nursing*, 42(6), 587-597.
- Hall, R. W., & Anand, K. J. (2014). Pain management in newborns. [Review]. *Clinics in perinatology*, 41(4), 895-924. doi: 10.1016/j.clp.2014.08.010
- Hallam, M. J., Shirley, R., Dheansa, B., & Coker, O. (2009). Scar massage therapy: a new technique. [Case Reports Letter]. *J Wound Care*, 18(6), 258.
- Harrison, D., Beggs, S., & Stevens, B. (2012). Sucrose for procedural pain management in infants. [Research Support, Non-U.S. Gov't Review]. *Pediatrics*, 130(5), 918-925. doi: 10.1542/peds.2011-3848
- Harrison, D., Yamada, J., & Stevens, B. (2010). Strategies for the prevention and management of neonatal and infant pain. [Review]. *Current pain and headache reports*, 14(2), 113-123. doi: 10.1007/s11916-009-0091-0
- Hartley, K. A., Miller, C. S., & Gephart, S. M. (2015). Facilitated tucking to reduce pain in neonates: evidence for best practice. [Research Support, Non-U S Gov't Research Support, U S Gov't, P H S Review]. *Adv Neonatal Care*, 15(3), 201-208.
- Hatherly, P. (2004). *The Homeopathic Physician's Guide to Lactation*. Chapel Hill, Queensland: Luminoz Pty Ltd.
- Hazelbaker, A. K. (2010). *Tongue-Tie: Morphogenesis, Impact, Assessment and Treatment*: Aidan and Eva Press.
- Heaton, P., & Herd, D. (2000). Neonatal pain relief. [Comment Letter]. *Archives of disease in childhood. Fetal and neonatal edition*, 83(1), F77-78.
- Heaton, P., Herd, D., & Fernando, A. (2007). Pain relief for simple procedures in New Zealand neonatal units: practice change over six years. *Journal of paediatrics and child health*, 43(5), 394-397. doi: 10.1111/j.1440-1754.2007.01086.x
- Ho, Y. B., Lee, R. S., Chow, C. B., & Pang, M. Y. (2010). Impact of massage therapy on motor outcomes in very low-birthweight infants: randomized controlled pilot study. [Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Pediatr Int*, 52(3), 378-385. doi: 10.1111/j.1442-200X.2009.02964.x
- Hoffmann, D. (2003). *Medical herbalism: the science and practice of herbal medicine*. Rochester, Vt.: Healing Arts Press.
- Holsti, L., Zwicker, J. G., & Grunau, R. E. (2016). Comment on the Paper by van den Bosch et al. Entitled 'Prematurity, Opioid Exposure and Neonatal Pain: Do They Affect the Developing Brain': The Impact of Subtle Messaging. [Comment]. *Neonatology*, 109(2), 120-121. doi: 10.1159/000442081
- Howard, F. M., Howard, C. R., Fortune, K., Generelli, P., Zolnoun, D., & tenHooopen, C. (1998). A randomized, placebo-controlled comparison of EMLA and dorsal penile nerve block for pain relief during neonatal circumcision. *Prim Care Update Ob Gyns*, 5(4), 196.
- Howe, T. H., Sheu, C. F., Hsieh, Y. W., & Hsieh, C. L. (2007). Psychometric characteristics of the Neonatal Oral-Motor Assessment Scale in healthy preterm infants. *Developmental medicine and child neurology*, 49(12), 915-919. Retrieved from <https://onlinelibrary.wiley.com/doi/pdfdirect/10.1111/j.1469-8749.2007.00915.x?download=true>
- Jain, S., Kumar, P., & McMillan, D. D. (2006). Prior leg massage decreases pain responses to heel stick in preterm babies. [Randomized Controlled Trial]. *Journal of paediatrics and child health*, 42(9), 505-508. doi: 10.1111/j.1440-1754.2006.00912.x
- Johnston, C. C., Fillion, F., Snider, L., Majnemer, A., Limperopoulos, C., Walker, C. D., Boyer, K. (2002). Routine sucrose analgesia during the first week of life in neonates younger than 31 weeks' postconceptional age. [Clinical Trial Multicenter Study Randomized Controlled Trial Research Support, Non-U S Gov't]. *Pediatrics*, 110(3), 523-528.

- Juif, P. E., Salio, C., Zell, V., Melchior, M., Lacaud, A., Petit-Demouliere, N., . . . Poisbeau, P. (2016). Peripheral and central alterations affecting spinal nociceptive processing and pain at adulthood in rats exposed to neonatal maternal deprivation. *Eur J Neurosci*, *44*(3), 1952-1962. doi: 10.1111/ejn.13294
- Karabulut, R., Sonmez, K., Turkyilmaz, Z., Demirogullari, B., Ozen, I. O., Bagbanci, B., Basaklar, A. C. (2008). Ankyloglossia and effects on breast-feeding, speech problems and mechanical/social issues in children. [Comparative Study]. *B-ENT*, *4*(2), 81-85.
- Karaca Ciftci, E., Kardas Ozdemir, F., & Aydin, D. (2016). Effect of flick application on pain level and duration of crying during infant vaccination. *Ital J Pediatr*, *42*, 8. doi: 10.1186/s13052-016-0218-y
- Kaufman, E., & Lamster, I. B. (2000). Analysis of saliva for periodontal diagnosis--a review. [Review]. *J Clin Periodontol*, *27*(7), 453-465.
- Kemper, K. J., Vohra, S., Walls, R., Complementary, t. T. F. o., Medicine, A., the Provisional Section on Complementary, H., & Medicine, I. (2008). The Use of Complementary and Alternative Medicine in Pediatrics. *Pediatrics*, *122*(6), 1374-1386. doi: 10.1542/peds.2008-2173
- Klockars, T., & Pitkaranta, A. (2009). Pediatric tongue-tie division: indications, techniques and patient satisfaction. [Comparative Study Research Support, Non-U.S. Gov't]. *International journal of pediatric otorhinolaryngology*, *73*(10), 1399-1401. doi: 10.1016/j.ijporl.2009.07.004
- Ko, W. J., Na, Y. C., Suh, B. S., Kim, H. A., Heo, W. H., Choi, G. H., & Lee, S. U. (2013). The effects of topical agent (kelo-cote or contractubex) massage on the thickness of post-burn scar tissue formed in rats. *Arch Plast Surg*, *40*(6), 697-704. doi: 10.5999/aps.2013.40.6.697
- Kothari, S. Y., Dongara, A. R., Nimbalkar, S. M., Phatak, A. G., & Nimbalkar, A. S. (2016). Missed Opportunities for Sedation and Pain Management at a Level III Neonatal Intensive Care Unit, India. *Front Pediatr*, *4*, 7. doi: 10.3389/fped.2016.00007
- Koulivand, P. H., Khaleghi Ghadiri, M., & Gorji, A. (2013). Lavender and the nervous system. *Evid Based Complement Alternat Med*, *2013*, 681304. doi: 10.1155/2013/681304
- Kumar, M., & Kalke, E. (2012). Tongue-tie, breastfeeding difficulties and the role of Frenotomy. *Acta paediatrica*, *101*(7), 687-689. doi: 10.1111/j.1651-2227.2012.02661.x
- Kupietzky, A., & Botzer, E. (2005). Ankyloglossia in the infant and young child: clinical suggestions for diagnosis and management. *Pediatric dentistry*, *27*(1), 40-46.
- Lago, P., Garetti, E., Merazzi, D., Pieragostini, L., Ancora, G., Pirelli, A., & Bellieni, C. V. (2009). Guidelines for procedural pain in the newborn. [Practice Guideline Review]. *Acta paediatrica*, *98*(6), 932-939.
- Larjava, H. (2012). *Oral wound healing: cell biology and clinical management*. Chichester, West Sussex, UK: Wiley-Blackwell.
- Larsson, B. A. (2001). [Pain and pain relief during the neonatal period. Early pain experiences can result in negative late-effects]. [Review]. *Lakartidningen*, *98*(14), 1656-1662.
- Lauwers, J., & Swisher, A. (2021). *Counseling the nursing mother: a lactation consultant's guide* (Seventh edition. ed.). Burlington, MA: Jones & Bartlett Learning.
- Lee, J. H., Espinera, A. R., Chen, D., Choi, K. E., Caslin, A. Y., Won, S., Yu, S. P. (2016). Neonatal inflammatory pain and systemic inflammatory responses as possible environmental factors in the development of autism spectrum disorder of juvenile rats. *J Neuroinflammation*, *13*(1), 109. doi: 10.1186/s12974-016-0575-x
- Lee, Y., Rodriguez, C., & Dionne, R. A. (2005). The role of COX-2 in acute pain and the use of selective COX-2 inhibitors for acute pain relief. [Research Support, U S Gov't, P H S Review]. *Curr Pharm Des*, *11*(14), 1737-1755.
- Lee, H. G., & Eun, H. C. (1999). Differences between fibroblasts cultured from oral mucosa and normal skin: implication to wound healing. *J Dermatol Sci*, *21*(3), 176-182.
- Lee, N. (2011). *Complementary & Alternative Medicine in Breastfeeding Therapy*. Amarillo, Texas: Hale Publishing.
- Lehr, J., Masters, A., & Pollack, B. (2012). Benzocaine-induced methemoglobinemia in the pediatric population. [Review]. *J Pediatr Nurs*, *27*(5), 583-588. doi: 10.1016/j.pedn.2012.07.003
- Lessen, B. S. (2011). Effect of the premature infant oral motor intervention on feeding progression and length of stay in preterm infants. *Adv Neonatal Care*, *11*(2), 129-139. doi:10.1097/ANC.0b013e3182115a2a
- Lessen Knoll, B. S., Daramas, T., & Drake, V. (2019). Randomized Controlled Trial of a Prefeeding Oral Motor Therapy and Its Effect on Feeding Improvement in a Thai NICU. *Journal of obstetric, gynecologic, and neonatal nursing: JOGNN / NAACOG*, *48*(2), 176-188. doi:10.1016/j.jogn.2019.01.003
- Levene, M. (1995). Pain relief and sedation during neonatal intensive care. *European journal of pediatrics*, *154*(8 Suppl 3), S22-23.
- Li, F., Huang, Q., Chen, J., Peng, Y., Roop, D. R., Bedford, J. S., & Li, C. Y. (2010). Apoptotic cells activate the "phoenix rising" pathway to promote wound healing and tissue regeneration. [Research Support, N.I.H.,

- Extramural Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, Non-P.H.S.]. *Sci Signal*, 3(110), ra13. doi: 10.1126/scisignal.2000634
- Li, X. L., Liu, Y., Liu, M., Yang, C. Y., & Yang, Q. Z. (2020). Early Premature Infant Oral Motor Intervention Improved Oral Feeding and Prognosis by Promoting Neurodevelopment. *Am J Perinatol*, 37(6), 626-632. doi:10.1055/s-0039-1685448
- Liaw, J. J., Yang, L., Lee, C. M., Fan, H. C., Chang, Y. C., & Cheng, L. P. (2013). Effects of combined use of non-nutritive sucking, oral sucrose, and facilitated tucking on infant behavioural states across heel-stick procedures: a prospective, randomised controlled trial. [Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *International journal of nursing studies*, 50(7), 883-894. doi: 10.1016/j.ijnurstu.2012.08.021
- Liu, J., Li, Z., & Zhang, Q. (2016). Challenges in the pain assessment during neonatal transport. An update. *Minerva pediatrica*.
- Longoni, L., Provenzi, L., Cavallini, A., Sacchi, D., Scotto di Minico, G., & Borgatti, R. (2018). Predictors and outcomes of the Neonatal Oral Motor Assessment Scale (NOMAS) performance: a systematic review. *European journal of pediatrics*, 177(5), 665-673. doi:10.1007/s00431-018-3130-1
- Lynch, S. E., de Castilla, G. R., Williams, R. C., Kiritsy, C. P., Howell, T. H., Reddy, M. S., & Antoniades, H. N. (1991). The effects of short-term application of a combination of platelet-derived and insulin-like growth factors on periodontal wound healing. [Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Journal of periodontology*, 62(7), 458-467. doi: 10.1902/jop.1991.62.7.458
- Marfurt-Russenberger, K., Axelin, A., Kesselring, A., Franck, L. S., & Cignacco, E. (2016). The Experiences of Professionals Regarding Involvement of Parents in Neonatal Pain Management. *Journal of obstetric, gynecologic, and neonatal nursing: JOGNN / NAACOG*, 45(5), 671-683. doi: 10.1016/j.jogn.2016.04.011
- Marucha, P. T., Kiecolt-Glaser, J. K., & Favagehi, M. (1998). Mucosal wound healing is impaired by examination stress. *Psychosomatic Medicine*, 60(3), 362-365.
- Mathew, P. J., & Mathew, J. L. (2003). Assessment and management of pain in infants. [Review]. *Postgraduate medical journal*, 79(934), 438-443.
- McDaniel, J. C., Massey, K., & Nicolaou, A. (2011). Fish oil supplementation alters levels of lipid mediators of inflammation in microenvironment of acute human wounds. [Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Wound Repair Regen*, 19(2), 189-200. doi: 10.1111/j.1524-475X.2010.00659.x
- Menon, G., Anand, K. J., & McIntosh, N. (1998). Practical approach to analgesia and sedation in the neonatal intensive care unit. [Review]. *Seminars in perinatology*, 22(5), 417-424.
- Mills, S., & Bone, K. (2005). *The Essential Guide to Herbal Safety*. St. Louis, Mo.: Elsevier Churchill Livingstone.
- Monell, T. (2011). Living out the Past: Infant Surgery Prior to 1987. *Journal of Prenatal and Perinatal Psychology and Health*, 25(Three).
- Montirosso, R., Casini, E., Del Prete, A., Zanini, R., Bellu, R., & Borgatti, R. (2016). Neonatal developmental care in infant pain management and internalizing behaviours at 18 months in prematurely born children. *Eur J Pain*, 20(6), 1010-1021. doi: 10.1002/ejp.826
- Montirosso, R., Casini, E., Del Prete, A., Zanini, R., Bellu, R., & Borgatti, R. (2016). Neonatal developmental care in infant pain management and internalizing behaviours at 18 months in prematurely born children. *Eur J Pain*, 20(6), 1010-1021. doi: 10.1002/ejp.826
- Mooney-Leber, S. M., & Brummelte, S. (2016). Neonatal pain and reduced maternal care: Early-life stressors interacting to impact brain and behavioral development. [Review]. *Neuroscience*. doi: 10.1016/j.neuroscience.2016.05.001
- Moore, T. E. (2001). The M and M's of aloe vera--is it for dentistry? [Historical Article]. *J Okla Dent Assoc*, 91(3), 30-31, 36.
- Moos, D. D., & Cuddeford, J. D. (2007). Methemoglobinemia and benzocaine. [Review]. *Gastroenterol Nurs*, 30(5), 342-345; quiz 346-347. doi: 10.1097/01.SGA.0000296253.78457.f4
- Moskowitz, R. (1992). *Homeopathic Medicines for Pregnancy and Childbirth*. Berkeley, Calif.: North Atlantic Books; Homeopathic Educational Services.
- Nahidi, F., Gazerani, N., Yousefi, P., & Abadi, A. R. (2017). The Comparison of the Effects of Massaging and Rocking on Infantile Colic. *Iran J Nurs Midwifery Res*, 22(1), 67-71. doi:10.4103/ijnmr.IJNMR_31_13
- Nevin, K. G., & Rajamohan, T. (2010). Effect of topical application of virgin coconut oil on skin components and antioxidant status during dermal wound healing in young rats. *Skin pharmacology and physiology*, 23(6), 290-297.
- Ohlsson, A., & Shah, P. S. (2016). Paracetamol (acetaminophen) for prevention or treatment of pain in newborns. [Review]. *Cochrane database of systematic reviews*, 10, CD011219. doi: 10.1002/14651858.CD011219.pub3
- O'Neill, M. C., Riddell, R. P., Garfield, H., & Greenberg, S. (2016). Does Caregiver Behaviour Mediate the Relationship Between Cultural Individualism and Infant Pain at 12 Months of Age? *J Pain*. doi: 10.1016/j.jpain.2016.08.008

- Osborne, C. M., & Robbins, B. (2012). Oral sucrose with facilitated tucking is effective pain control for preterm infants. [Comment]. *The Journal of pediatrics*, *161*(1), 169-170. doi: 10.1016/j.jpeds.2012.04.050
- Pados, B. F., & McGlothen-Bell, K. (2019). Benefits of Infant Massage for Infants and Parents in the NICU. *Nurs Womens Health*, *23*(3), 265-271. doi:10.1016/j.nwh.2019.03.004
- Page, G. G. (2004). Are there long-term consequences of pain in newborn or very young infants? *The Journal of perinatal education: an ASPO/Lamaze publication*, *13*(3), 10-17. doi: 10.1624/105812404X1725
- Palmer, M. (2013). *NOMAS Certification Course*. Paper presented at the NOMAS Certification Course, Walnut Creek, California.
- Palomaa, A. K., Korhonen, A., & Polkki, T. (2016). Factors Influencing Parental Participation in Neonatal Pain Alleviation. *J Pediatr Nurs*, *31*(5), 519-527. doi: 10.1016/j.pedn.2016.05.004
- Palomaa, A. K., Korhonen, A., & Polkki, T. (2016). PO20 - Parental suggestions for developing pain management in neonatal intensive care units. *Nurs Child Young People*, *28*(4), 100. doi: 10.7748/ncyp.28.4.100.s99
- Parente, L. M., Lino Junior Rde, S., Tresvenzol, L. M., Vinaud, M. C., de Paula, J. R., & Paulo, N. M. (2012). Wound Healing and Anti-Inflammatory Effect in Animal Models of *Calendula officinalis* L. Growing in Brazil. *Evid Based Complement Alternat Med*, *2012*, 375671. doi: 10.1155/2012/375671
- Peirsman, E. a. N. (2006). *Craniosacral Therapy for Babies and Small Children*. Berkeley, California: North Atlantic books.
- Perrott, D. A., Piira, T., Goodenough, B., & Champion, G. D. (2004). Efficacy and safety of acetaminophen vs ibuprofen for treating children's pain or fever: a meta-analysis. [Comparative Study Meta-Analysis Research Support, Non-U S Gov't Review]. *Archives of pediatrics & adolescent medicine*, *158*(6), 521-526.
- Pillai Riddell, R., Fitzgerald, M., Slater, R., Stevens, B., Johnston, C., & Campbell-Yeo, M. (2016). Using only behaviours to assess infant pain: a painful compromise? *Pain*, *157*(8), 1579-1580. doi: 10.1097/j.pain.0000000000000598
- Pinelli, J., & Symington, A. (2001). Non-nutritive sucking for promoting physiologic stability and nutrition in preterm infants. [Review]. *Cochrane database of systematic reviews*, *3*.
- Pinelli, J., Symington, A., & Ciliska, D. Nonnutritive Sucking in High-Risk Infants: Benign Intervention or Legitimate Therapy? *Journal of Obstetric, Gynecologic & Neonatal Nursing*, *31*(5), 582-591. doi: 10.1111/j.1552-6909.2002.tb00084.x
- Polkki, T., Korhonen, A., Heikkinen, U., Palomaa, A. K., & Miettinen, S. (2016). PO19 - Developing a pain scale for infants in collaboration with clinical staff in the neonatal intensive care unit. *Nurs Child Young People*, *28*(4), 99-100. doi: 10.7748/ncyp.28.4.99.s98
- Pölkki, T., Korhonen, A., & Laukkala, H. Nurses' Expectations of Using Music for Premature Infants in Neonatal Intensive Care Unit. *Journal of Pediatric Nursing: Nursing Care of Children and Families*, *27*(4), e29-e37. doi: 10.1016/j.pedn.2011.06.001
- Porter, F. L., Grunau, R. E., & Anand, K. J. (1999). Long-term effects of pain in infants. [Review]. *Journal of developmental and behavioral pediatrics: JDBP*, *20*(4), 253-261.
- Preethi, K. C., & Kuttan, R. (2009). Wound healing activity of flower extract of *Calendula officinalis*. [Research Support, Non-U.S. Gov't]. *Journal of basic and clinical physiology and pharmacology*, *20*(1), 73-79.
- Prestes, A. C., Guinsburg, R., Balda, R. C., Marba, S. T., Rugolo, L. M., Pachi, P. R., & Bentlin, M. R. (2005). [The frequency of pharmacological pain relief in university neonatal intensive care units]. [Multicenter Study]. *Jornal de pediatria*, *81*(5), 405-410. doi: 10.2223/JPED.1392
- Procianoy, R. S., Mendes, E. W., & Silveira, R. C. (2010). Massage therapy improves neurodevelopment outcome at two years corrected age for very low birth weight infants. [Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Early human development*, *86*(1), 7-11. doi: 10.1016/j.earlhumdev.2009.12.001
- Provenzi, L., Giusti, L., Fumagalli, M., Tasca, H., Ciceri, F., Menozzi, G., Montiroso, R. (2016). Pain-related stress in the Neonatal Intensive Care Unit and salivary cortisol reactivity to socio-emotional stress in 3-month-old very preterm infants. *Psychoneuroendocrinology*, *72*, 161-165. doi: 10.1016/j.psyneuen.2016.07.010
- Ranger, M., & Grunau, R. E. (2014). Early repetitive pain in preterm infants in relation to the developing brain. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Review]. *Pain Manag*, *4*(1), 57-67. doi: 10.2217/pmt.13.61
- Reece-Stremtan, S., & Gray, L. (2016). ABM Clinical Protocol #23: Nonpharmacological Management of Procedure-Related Pain in the Breastfeeding Infant, Revised 2016. *Breastfeeding medicine: the official journal of the Academy of Breastfeeding Medicine*. doi: 10.1089/bfm.2016.29025.srs
- Rennix, C., Manjunatha, C. M., & Ighanesebhor, S. E. (2004). Pain relief during common neonatal procedures: a survey. [Letter]. *Archives of disease in childhood. Fetal and neonatal edition*, *89*(6), F563. doi: 10.1136/adc.2003.045351

- Ribeiro, L. M., Castral, T. C., Montanholi, L. L., Dare, M. F., Silva, A. C., Antonini, S. R., & Scochi, C. G. (2013). [Human milk for neonatal pain relief during ophthalmoscopy]. *Rev Esc Enferm USP*, 47(5), 1039-1045. doi: 10.1590/S0080-623420130000500005
- Riordan, J., Gross, A., Angeron, J., Krumwiede, B., & Melin, J. (2000). The effect of labor pain relief medication on neonatal suckling and breastfeeding duration. [Research Support, Non-U.S. Gov't]. *Journal of human lactation: official journal of International Lactation Consultant Association*, 16(1), 7-12.
- Rocha, A. D., Moreira, M. E., Pimenta, H. P., Ramos, J. R., & Lucena, S. L. (2007). A randomized study of the efficacy of sensory-motor-oral stimulation and non-nutritive sucking in very low birthweight infant. *Early human development*, 83(6), 385-388. doi:10.1016/j.earlhumdev.2006.08.003
- Rohan, A. J. (2016). Pain-associated stressor exposure and neuroendocrine values for premature infants in neonatal intensive care. *Developmental psychobiology*, 58(1), 60-70. doi: 10.1002/dev.21346
- Salo, T., Makela, M., Kylmaniemi, M., Autio-Harmanen, H., & Larjava, H. (1994). Expression of matrix metalloproteinase-2 and -9 during early human wound healing. [Research Support, Non-U.S. Gov't]. *Lab Invest*, 70(2), 176-182.
- Saniski, D. (2005). Neonatal pain relief protocols in their infancy. *Nebr Nurse*, 38(2), 24.
- Schlez, A., Litmanovitz, I., Bauer, S., Dolfin, T., Regev, R., & Arnon, S. (2011). Combining kangaroo care and live harp music therapy in the neonatal intensive care unit setting. [Comparative Study Randomized Controlled Trial]. *Isr Med Assoc J*, 13(6), 354-358.
- Schwaller, F., & Fitzgerald, M. (2014). The consequences of pain in early life: injury-induced plasticity in developing pain pathways. [Review]. *Eur J Neurosci*, 39(3), 344-352. doi: 10.1111/ejn.12414
- Seo, Y. S., Lee, J., & Ahn, H. Y. (2016). Effects of Kangaroo Care on Neonatal Pain in South Korea. *Journal of tropical pediatrics*, 62(3), 246-249. doi: 10.1093/tropej/fmv102
- Shah, P. S., Aliwalas, L. I., & Shah, V. (2006). Breastfeeding or breast milk for procedural pain in neonates. [Meta-Analysis Review]. *Cochrane database of systematic reviews*, 19(3).
- Shandley, S., Capilouto, G., Tamilya, E., Riley, D. M., Johnson, Y. R., & Papadelis, C. (2020). Abnormal Nutritive Sucking as an Indicator of Neonatal Brain Injury. *Front Pediatr*, 8, 599633. doi:10.3389/fped.2020.599633
- Shepherd, A. A. (2003). Nutrition for optimum wound healing. [Review]. *Nurs Stand*, 18(6), 55-58.
- Sherman, A. R., & Barkley, M. (2011). Nutrition and wound healing. *J Wound Care*, 20(8), 357-358, 360, 362-357.
- Shin, T. M., & Bordeaux, J. S. (2012). The role of massage in scar management: a literature review. [Review]. *Dermatol Surg*, 38(3), 414-423. doi: 10.1111/j.1524-4725.2011.02201.x
- Smith, C., & Goldman, R. D. (2012). Alternating acetaminophen and ibuprofen for pain in children. *Canadian family physician Medecin de famille canadien*, 58(6), 645-647.
- So, T. Y., & Farrington, E. (2008). Topical benzocaine-induced methemoglobinemia in the pediatric population. [Case Reports]. *Journal of pediatric health care: official publication of National Association of Pediatric Nurse Associates & Practitioners*, 22(6), 335-339; quiz 340-331. doi: 10.1016/j.pedhc.2008.08.008
- Spelman, K. (2011). Ecological Pharmacy: Molecular Biology to Systems Theory *Fundamentals of Complementary and Alternative Medicine* (4th ed., pp. 454-475). St. Louis, Mo.: Saunders/Elsevier.
- Standley, J. (2012). Music therapy research in the NICU: an updated meta-analysis. [Meta-Analysis Review]. *Neonatal network: NN*, 31(5), 311-316.
- Stevens, B., Yamada, J., & Ohlsson, A. (2004). Sucrose for analgesia in newborn infants undergoing painful procedures. [Review]. *Cochrane database of systematic reviews*, 3.
- Stillerman, E. (2009). C-section scar massage. *Midwifery today with international midwife* (92), 29, 64-25.
- Suter, V. G., & Bornstein, M. M. (2009). Ankyloglossia: facts and myths in diagnosis and treatment. [Review]. *Journal of periodontology*, 80(8), 1204-1219. doi: 10.1902/jop.2009.090086
- Szpaderska, A. M., Zuckerman, J. D., & DiPietro, L. A. (2003). Differential injury responses in oral mucosal and cutaneous wounds. [Comparative Study Research Support, U.S. Gov't, P.H.S.]. *Journal of dental research*, 82(8), 621-626.
- Tanideh, N., Tavakoli, P., Saghiri, M. A., Garcia-Godoy, F., Amanat, D., Tadbir, A. A., Tamadon, A. (2013). Healing acceleration in hamsters of oral mucositis induced by 5-fluorouracil with topical *Calendula officinalis*. [Comparative Study]. *Oral Surg Oral Med Oral Pathol Oral Radiol*, 115(3), 332-338. doi: 10.1016/j.oooo.2012.08.450
- Thabet, A. M., & Sayed, Z. A. (2021). Effectiveness of the Premature Infant Oral Motor Intervention on Feeding Performance, Duration of Hospital Stay, and Weight of Preterm Neonates in Neonatal Intensive Care Unit: Results From a Randomized Controlled Trial. *Dimens Crit Care Nurs*, 40(4), 257-265. doi:10.1097/DCC.0000000000000475

- Tohill, J., & McMorrow, O. (1990). Pain relief in neonatal intensive care. [Letter]. *Lancet*, 336(8714), 569.
- Tsai, S. W., Chen, C. H., & Lin, M. C. (2010). Prediction for developmental delay on Neonatal Oral Motor Assessment Scale in preterm infants without brain lesion. *Pediatr Int*, 52(1), 65-68. doi:10.1111/j.1442-200X.2009.02882.x
- Tsao, J. C., Evans, S., Meldrum, M., Altman, T., & Zeltzer, L. K. (2008). A Review of CAM for Procedural Pain in Infancy: Part I. Sucrose and Non-nutritive Sucking. *Evid Based Complement Alternat Med*, 5(4), 371-381. doi: 10.1093/ecam/nem084
- Tsao, J. C., Evans, S., Meldrum, M., Altman, T., & Zeltzer, L. K. (2008). A Review of CAM for Procedural Pain in Infancy: Part II. Other Interventions. *Evid Based Complement Alternat Med*, 5(4), 399-407. doi: 10.1093/ecam/nem089
- Tsao, J. C., & Zeltzer, L. K. (2005). Complementary and Alternative Medicine Approaches for Pediatric Pain: A Review of the State-of-the-science. *Evid Based Complement Alternat Med*, 2(2), 149-159. doi: 10.1093/ecam/neh092
- Vaglio, S. (2009). Chemical communication and mother-infant recognition. *Commun Integr Biol*, 2(3), 279-281.
- van den Hoogen, N. J., Tibboel, D., Honig, W. M., Hermes, D., Patijn, J., & Joosten, E. A. (2016). Neonatal paracetamol treatment reduces long-term nociceptive behaviour after neonatal procedural pain in rats. *Eur J Pain*, 20(8), 1309-1318. doi: 10.1002/ejp.855
- van Ganzewinkel, C. J., Been, J. V., Dieleman, J. P., Katgert, T., Boelen-van der Loo, T., van der Pal, S. M., Andriessen, P. (2016). Pain coping strategies: Neonatal intensive care unit survivors in adolescence. *Early human development*, 103, 27-32. doi: 10.1016/j.earlhumdev.2016.07.003
- Van Howe, R. S., & Svoboda, J. S. (2008). Neonatal pain relief and the Helsinki Declaration. *J Law Med Ethics*, 36(4), 803-823, 611. doi: 10.1111/j.1748-720X.2008.00339.x
- Vinall, J., & Grunau, R. E. (2014). Impact of repeated procedural pain-related stress in infants born very preterm. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Review]. *Pediatric research*, 75(5), 584-587. doi: 10.1038/pr.2014.16
- Vogel, S. (1986). Oral motor and feeding problems in the tube fed infant. *Occup Ther Health Care*, 3(2), 63-79. doi:10.1080/J003v03n02_06
- Walker, D. G. (1954). Nutrition in oral surgery and its relationship to wound healing and infection. *Oral surgery, oral medicine, and oral pathology*, 7(8), 797-812.
- Walls, A., Pierce, M., Wang, H., Steehler, A., Steehler, M., & Harley Jr, E. H. (2014). Parental perception of speech and tongue mobility in three-year olds after neonatal frenotomy. *International journal of pediatric otorhinolaryngology*, 78(1), 128-131. doi:http://dx.doi.org/10.1016/j.ijporl.2013.11.006
- Wambach, K., & Spencer, B. (2021). *Breastfeeding and human lactation* (Sixth edition. ed.). Burlington, Massachusetts: Jones & Bartlett Learning.
- Ward, M. J., Cronin, K. B., Renfro, P. D., Lowman, D. K., & Cooper, P. D. (2000). Oral motor feeding in the neonatal intensive care unit: exploring perceptions of parents and occupational therapists. *Occup Ther Health Care*, 12(2-3), 19-37. doi:10.1080/J003v12n02_02
- Warnock, F. F., Castral, T. C., Brant, R., Sekilian, M., Leite, A. M., Owens Sde, L., & Scochi, C. G. (2010). Brief report: Maternal kangaroo care for neonatal pain relief: a systematic narrative review. [Research Support, Non-U.S. Gov't Review]. *J Pediatr Psychol*, 35(9), 975-984. doi: 10.1093/jpepsy/jsp123
- West, Z. (2008). *Acupuncture in Pregnancy and Childbirth* (2nd ed.). Edinburgh; New York: Churchill Livingstone/Elsevier.
- Williams, W. N., & Waldron, C. M. (1985). Assessment of lingual function when ankyloglossia (tongue-tie) is suspected. *Journal of the American Dental Association*, 110(3), 353-356. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3858347>
- Wilson-Smith, E. M. (2011). Procedural Pain Management in Neonates, Infants and Children. *Rev Pain*, 5(3), 4-12. doi: 10.1177/204946371100500303
- Witt, N., Coynor, S., Edwards, C., & Bradshaw, H. (2016). A Guide to Pain Assessment and Management in the Neonate. [Review]. *Curr Emerg Hosp Med Rep*, 4, 1-10. doi: 10.1007/s40138-016-0089-y
- Wolf, L. S., & Glass, R. P. (1992). *Feeding and swallowing disorders in infancy: assessment and management*. Tucson, Ariz.: Therapy Skill Builders.
- Wong, J. W., Gallant-Behm, C., Wiebe, C., Mak, K., Hart, D. A., Larjava, H., & Hakkinen, L. (2009). Wound healing in oral mucosa results in reduced scar formation as compared with skin: evidence from the red Duroc pig model and humans. [Comparative Study Research Support, Non-U.S. Gov't Validation Studies]. *Wound Repair Regen*, 17(5), 717-729. doi: 10.1111/j.1524-475X.2009.00531.x

- Yang, C., Liu, F., Zhou, L., Shen, Q., Jia, H., Xu, P., Stephanie, L. (2019). [Clinical significance of oral motor intervention on the prognosis of early premature infant]. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue*, 31(2), 150-154. doi:10.3760/cma.j.issn.2095-4352.2019.02.006
- Yawman, D., Howard, C. R., Auinger, P., Garfunkel, L. C., Allan, M., & Weitzman, M. (2006). Pain relief for neonatal circumcision: a follow-up of residency training practices. [Research Support, Non-U.S. Gov't]. *Ambul Pediatr*, 6(4), 210-214. doi: 10.1016/j.ambp.2006.04.008
- Yin, T., Yang, L., Lee, T. Y., Li, C. C., Hua, Y. M., & Liaw, J. J. (2015). Development of atraumatic heel-stick procedures by combined treatment with non-nutritive sucking, oral sucrose, and facilitated tucking: a randomised, controlled trial. [Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *International journal of nursing studies*, 52(8), 1288-1299. doi: 10.1016/j.ijnurstu.2015.04.012
- Younesian, S., Yadegari, F., & Soleimani, F. (2015). Impact of Oral Sensory Motor Stimulation on Feeding Performance, Length of Hospital Stay, and Weight Gain of Preterm Infants in NICU. *Iran Red Crescent Med J*, 17(7), e13515. doi:10.5812/ircmj.17(5)2015.13515
- Zaghi, S., Valcu-Pinkerton, S., Jabara, M., Norouz-Knutsen, L., Govardhan, C., Moeller, J., Liu, S. Y. (2019). Lingual frenuloplasty with myofunctional therapy: Exploring safety and efficacy in 348 cases. *Laryngoscope Investig Otolaryngol*, 4(5), 489-496. doi:10.1002/lio2.297
- Zarem, C., Kidokoro, H., Neil, J., Wallendorf, M., Inder, T., & Pineda, R. (2013). Psychometrics of the neonatal oral motor assessment scale. *Dev Med Child Neurol*, 55(12), 1115-1120. doi:10.1111/dmcn.12202